

A Demographic Model of Oyster Populations in the Chesapeake Bay to Evaluate Proposed Oyster-Restoration Alternatives

4 October 2007

Jon H. Vølstad^{1,3}, Mary C. Christman^{2,4}, Danny Lewis², Edward D. Weber¹, and Jodi Dew¹

From the chaos in which I found the business, so far as regards statistical information, I have tried to evolve some facts and figures by which, by showing the importance of the trade, may cause a more careful study to be made of the means to arrest the present depletion of the beds and provide ways for increasing the natural supply of oysters. (Edmonds 1887)

¹Versar, Inc., 9200 Rumsey Road, Columbia, MD 21045

²University of Maryland, College Park, MD 20742

³Present address: Institute of Marine Research, Bergen, Norway

⁴Present address: University of Florida, IFAS Statistics Department,
Gainesville, FL, 32611

Preface

This manuscript describes a demographic model for the Eastern oyster (*Crassostrea virginica*) in the Chesapeake Bay that was developed using existing data from many researchers and sources. The authors developed the model structure, parameterized most of the input variables, and implemented the model in the Java™ programming language; however, the model includes several components provided by other scientists. The Maryland Department of Natural Resources (MDNR) estimated the population of oysters at model initiation, identified available substrate that oysters could occupy, and defined oyster restoration scenarios to be projected. We describe MDNR's procedures for developing those components briefly in the text; their detailed documentation of those procedures is presented in appendices to this report. The distribution of spat settlement among bars employed in the demographic model is based on output of a larval-transport model developed by E. North and colleagues at the University of Maryland Center for Environmental Science. Output of the demographic model, in terms of oyster abundance, will serve as input to a model developed by C. Cerco, U.S. Army Corps of Engineers Environmental Laboratory, to predict the potential effects of oyster restoration alternatives on water quality and SAV.

We thank the many researchers who contributed data and expert advice for use in the model. From the Maryland Department of Natural Resources, Fisheries Division: L. Barker provided the abundance and distribution of the oyster population at model initiation; E. Cambell, provided the detailed management scenarios to be modeled for native-oyster restoration; P. Genovese modeled salinity of oyster bars as a function of annual precipitation levels; K. Greenhawk developed the map of the estimated amount habitat that oysters could occupy; P. Jones reviewed the model and developed the report on native-oyster restoration alternatives. C. Judy provided survey data, expert advice, and model review; T. O'Connell provided expert advice and reviewed the model; M. Tarnowski served as a co-author for the mortality-estimation portion of model, and provided survey data, expert advice, and model review. R. Mann and J. Harding, Virginia Institute of Marine Science, provided density and growth data for oysters in the James River. From the University of Maryland Center for Environmental Science, J. Coakley provided height-frequency analysis used to estimate growth in Maryland, R. Newell provided expert opinion on oyster biology and suggestions for modeling Suminoe oyster dynamics, and K. Paynter, provided data used to estimate oyster growth. J. Wesson, Virginia Marine Resources Commission Division of Fisheries Management, provided descriptions of native-oyster restoration alternatives for Virginia, data on available oyster habitat and harvest estimates for the James River. S. Ford, Rutgers University, provided expert advice and review of the mortality estimation methods. We also thank the EIS independent oyster advisory panel consisting of J. Anderson, University of Rhode Island, M. Berrigan, Florida Department of Agriculture and Consumer Services, M. Heral, French Research Institute for Exploitation of the Sea, R. Mann, Virginia Institute of Marine Science, E. Powell, Rutgers University, M. Roman, University of Maryland, and B. Rothschild, Dartmouth University, for reviewing the model.

Table of Contents

| | |
|--|-----|
| Preface | ii |
| Table of Contents | iii |
| List of Tables | iv |
| List of Figures | v |
| List of Appendices | vii |
| 1.0 Introduction..... | 1 |
| 2.0 Data Sources and Parameter Estimates..... | 2 |
| 2.1 Maryland Department of Natural Resources Annual Fall Survey | 2 |
| 2.2 Available Habitat | 3 |
| 2.3 Starting Population..... | 4 |
| 2.4 Flow Regime..... | 5 |
| 2.4.1 Weather years..... | 5 |
| 2.4.2 Freshets | 6 |
| 2.5 Salinity | 6 |
| 2.6 Recruitment..... | 7 |
| 2.6.1 Recruitment Data Sources..... | 7 |
| 2.6.2 Recruitment Model | 7 |
| 2.7 Larval Transport..... | 11 |
| 2.8 Growth | 11 |
| 2.8.1 Growth Data Sources | 11 |
| 2.8.2 Growth Model | 13 |
| 2.9 Disease Intensity | 14 |
| 2.10 Natural Mortality..... | 15 |
| 2.11 Harvest | 17 |
| 2.12 Habitat Improvements | 18 |
| 3.0 Model Structure | 19 |
| 4.0 Description of Proposed Action and Alternatives | 20 |
| 5.0 Model Performance..... | 22 |
| 6.0 Alternative 1-3 Results | 24 |
| 7.0 Model Sensitivity..... | 26 |
| 8.0 Data and Uncertainty | 28 |
| 9.0 Conclusions..... | 30 |
| 10.0 Literature Cited..... | 31 |
| 11.0 Tables..... | 35 |
| 12.0 Figures | 39 |
| 1.0 Step 1 - Delineation of Maryland Basins..... | 3 |
| 2.0 Step 3 – Estimation of Oyster Density in High and Low Quality Habitat..... | 6 |

List of Tables

| Table | Page |
|---|-------------|
| Table 1. Number of oysters in the starting population at model initiation by state, salinity zone, and aggregated to the entire Bay. Estimates are based on abundance for 2004, a wet year. Note that the area of salinity zones (low < 11 ppt, medium 11-15 ppt, or high \geq 15 ppt) fluctuates with annual precipitation. Thus, oysters on individual bars may change salinity zones during different years in the model. | 35 |
| Table 2. Estimated proportions of female oysters in the population (sex ratio) by size class. Data are based on field experiments reported by Kennedy (1983). Proportions are denoted as R in Equation 3, where $r = (0-40), (41-80), \dots, (>160)$ | 35 |
| Table 3. Probability of selecting disease tier 1, 2, or 3 based on the current weather year for individual bars in the demographic model. Probabilities are based on mean disease intensities measured at disease bars in the MD DNR fall survey for the years 1990-2005. | 35 |
| Table 4. Average annual natural mortality rates based on recent box counts for eastern oysters by salinity class and disease tier. Average time-since-death of recent boxes was assumed to be between 7 and 14 days. The relative standard error, $RSE = SE/Mean$; LCL and UCL are the upper and lower 95% confidence limits, respectively. | 36 |
| Table 5. Sensitivity of demographic model to variation in underlying parameters. All sensitivity runs were compared to the run for alternative 1, no action, with a 40% harvest rate (Figure 11) unless otherwise specified. Results are reported as the percent change in median oyster abundance at year ten of the model. | 37 |

List of Figures

| Figure | Page |
|--|------|
| Figure 1. Estimated density and distribution of oysters in the starting population (spat, small, and market sized) for 8,480 bars included in the demographic model, based on year 2004. | 39 |
| Figure 2. Mean annual discharge into the Chesapeake Bay between 1937 and 2005 (Source U.S. Geological Survey, Chesapeake Bay Activities, http://md.water.usgs.gov/monthly/bay.html). | 40 |
| Figure 3. Salinity classes (low < 11 ppt, medium 11-15 ppt, or high ≥ 15 ppt) of oyster bars in the demographic model during 1995-1999. | 41 |
| Figure 4. Comparison of sampling efficiency of dredging versus diver harvest reported for replicated studies at Hog Island and Walter White reefs in the Chesapeake Bay by Chai et al. (1992). Efficiency of the dredge was low for both reefs (A). Numbers of spat captured relative to small and market-size oysters were similar, as indicated by the same data plotted on different scales (B). Estimates from the dredge survey are scaled by 40.6 times on the secondary axis in B. Vertical bars indicate approximate 95% confidence intervals. | 42 |
| Figure 5. Relationship between number of standardized female spawners (to 77 mm) and spat based on the DNR annual dredge survey, 1990-2006. Gray line indicates a ratio of 3.0 spat per spawner. Observations with higher ratios were considered outliers and used to model episodic large spat sets. Spat were modeled according to salinity as indicated by the dark lines in normal recruitment years. | 43 |
| Figure 6. Height-at-age distributions used to model oyster growth in the Chesapeake Bay. A. Data from two recent studies and a literature review that were applied to Maryland and the Potomac River. B. Data from a survey of the James River that were applied to Virginia except the Potomac River. Lines indicate the 5th, 50th, and 95th percentiles of each model. The mean maximum size of oysters H_{∞} was assumed to be 177 mm in both models based on literature (Rothschild et al. 1994) because data were not available for older oysters. The median line for Maryland is $H_{\infty} = 177 * (1 - \exp(-0.1834 * (\text{age} + 0.7232)))$. The median line for Virginia is $H_{\infty} = 177 * (1 - \exp(-0.1078 * (\text{age} + 0.7414)))$ | 44 |
| Figure 7. Mean annual natural mortality rates by disease level (tier 1 = high dermo intensity; tier 2 = medium dermo intensity; and tier 3 = low dermo intensity) and salinity class for small and market-size oysters over all years (1991-2005). Error bars represent 95% confidence intervals. We assumed an average TSD of 1-year for total boxes, and 1-2 -weeks for recent boxes. | 45 |
| Figure 8. The function: Shell condition = $1.593 + [0.939 \times \log_e(\text{age})]$ describes a decline in the quality of planted shell for oyster recruitment as planted shell is covered with sediment (Smith 2005; left axis). Condition-class 1 indicates clean shell with no sediment. Condition-class 3 represents shells that have received some sedimentation but are similar to unreplenished bars. Mean spat settlement was observed to be 4.5 times greater on newly replenished bars than on unreplenished bars (MDDNR, unpublished data). Thus, a scaling factor of 4.5 (right axis) was applied to replenished bars in the demographic model for the first year, and the scaling factor was decreased through time following the same rate of decline as the shell-condition model. | 46 |

Figure 9. Flow diagram of oyster demographic model. 47

Figure 10. Abundance of oysters in the James River, Virginia, as predicted by 1,000 runs of the oyster demographic model (solid line), and field samples (dashed line). The number of spat < 30 mm was input into the model each year for run A, and estimated by the model for run B. Model estimates are the median value of all runs. Vertical bars indicate 90% confidence intervals, as estimated by the 5th and 95th percentile of all runs..... 48

Figure 11. Change in oyster abundance from the starting population to year 10 projected by 1,000 runs of the demographic model for EIS alternatives 1 (status quo management), 2A, 2B (enhanced restoration), and 3 (no harvest). Harvest rates indicate winter exploitation rates of 20 to 80% on legally harvestable bars (71% of the total), after natural mortality has occurred. Boxes indicate the range of the 25th to 75th percentile of all runs. Vertical bars extend to the 5th and 95th percentile of all runs. Results are presented for the entire Bay (A), Maryland and Virginia (B), and by salinity zone (D). Results for Maryland and Virginia are also presented as change in predicted total abundance rather than percent change from the starting population (C). Note that scales differ among plots for spat, small, and market-size oysters on some panels. 49

Figures 12-17. Oyster abundance projected by state in 1,000 runs of the demographic model for EIS alternatives 1 (status quo management), 2A, 2B (enhanced restoration), and 3 (no harvest). Results are reported for harvest rates (winter exploitation rates) of 20 to 80% on legally harvestable bars (71% of the total), after natural mortality has occurred, and categorized by the entire Bay, states, and salinity zones. Tables indicate percent changes in abundance from the starting population at the 5th, 25th, 50th (median), 75th, and 95th percentile for each year of the simulation. Data are depicted as boxplots to the right of each table to illustrate trends. Square symbols indicate the starting population (0% change). Scales differ for each plot. Note that salinity zones do not represent fixed areas. Individual bars may change salinity depending on annual precipitation, but the starting population is for a single wet year based on 2004. 54

List of Appendices

| | Page |
|---|--|
| A | Development of Habitat Layer..... A-1 |
| B | Oyster Population Estimates for the Maryland Portion of Chesapeake Bay 1994 – 2006.....B-1 |
| C | Development of Starting Population.....C-1 |
| D | Estimation of Annual Mortality Rates for Eastern Oysters in Chesapeake Bay Based on Box Counts and Application of those Rates to Project Population Growth of <i>C. virginica</i> and <i>C. ariakensis</i> D-1 |
| E | Description of Alternative Restoration StrategiesE-1 |
| F | Explanation for Effects of Harvest F-1 |

1.0 Introduction

The U.S. Army Corps of Engineers, the states of Maryland and Virginia, and several cooperating agencies are evaluating strategies for restoring depleted oyster populations in the Chesapeake Bay. Nine alternative strategies are being considered that include various combinations of restoration of the native Eastern oyster (*Crassostrea virginica*), augmentation of wild populations with aquaculture, and introduction of the non-native Suminoe oyster (*C. ariakensis*; the proposed action). Recent research indicates that Suminoe oysters are resistant to diseases that have been an important cause of the decline in native oyster populations; however, Suminoe oysters could have undesirable effects on the ecosystem. The lead agencies, therefore, are conducting an ecological risk assessment and preparing a programmatic Environmental Impact Statement (EIS) to evaluate the relative risks and benefits of each alternative management strategy. A major component of this assessment has been the development of a demographic model to predict oyster population dynamics for ten years after the start of each alternative restoration approach. The purpose of this document is to describe the structure and parameterization of the oyster demographic model.

An important consideration in modeling population dynamics of oysters is that differences in the physical and chemical characteristics of the water have biologically significant effects on the spatial structure of populations. A suitable model must account for differences in recruitment, growth, and survival rates among oyster bars distributed throughout the estuary. Water current is one characteristic that affects oyster populations. Currents mix water of different temperatures and salinities, and disperse larval oysters. The locations of bars in relation to currents causes some bars to act as sources of larvae that are exported and others to act as sinks that receive large numbers of larvae. Differences in vertical swimming behavior between Eastern and Suminoe oysters (Newell et al. 2005) also interact with current, resulting in different patterns of transport between the two species (North et al. 2006, In Press) and different population distributions. Another hydrologic characteristic that strongly affects oysters is freshwater tributary inflow. Freshwater inflow controls salinity at individual bars, which strongly affects oyster recruitment and survival. Storms that cause freshets can kill oysters in upstream bars by reducing salinity below oysters' physiological tolerance. Tributary inflow and salinity also affect recruitment of spat. Areas of the Bay with high salinity generally exhibit the greatest recruitment (Ulanowicz et al. 1980; Tarnowski et al. 2003); therefore, recruitment is greater in downstream portions of the Bay located nearer to the ocean and during dry-weather years, when freshwater inflow is reduced. Salinity also affects survival of Eastern oysters via disease prevalence. Dermo disease, caused by the parasite *Perkinsus marinus*, and MSX disease, caused by the parasite *Haplosporidium nelsoni*, are major causes of mortality for Eastern oysters (Ford and Trip 1996). Both diseases are more prevalent in high-salinity water.

To account for the heterogeneous population structure among oyster bars in the Chesapeake Bay, we developed a spatially explicit model that incorporates interactions between hydrologic conditions and oyster populations. The demographic model accounts for variation

among bars in several ways. First, larval oysters are distributed among bars by linking the demographic model with predictions from the larval-transport model developed by North et al. (2006; In Press). The larval-transport model combines hydrodynamic modeling of the Bay during different weather conditions (inflow, wind, tides, etc) and observed characteristics about the vertical swimming behavior of larvae to forecast the distribution of larvae among bars. A second source of spatial variation included in the demographic model is salinity, which is predicted for each bar based on location and weather and used to adjust recruitment, disease prevalence, and natural mortality rates. Finally, variation in growth is modeled spatially at the level of state (Maryland or Virginia portions of the Bay) to match empirical data, although growth rates did not appear strongly related to inflow or salinity.

A large set of variables related to larval dispersion patterns, growth, recruitment, survival, hydrologic conditions, available habitat, and starting population size were parameterized for inclusion in the model. For clarity, we describe the specific parameterization of individual components first, then implementation and performance of the model in general. First we list the data sources for estimating input parameters and how their mean trends and variability were calculated for inclusion in the model (Section 2). We then describe how these components were combined to predict oyster abundance in the demographic model in Section 3, Model Structure. We follow the description of the model with a list of specific management scenarios simulated (Section 4), and an evaluation of model performance using a data set from the James River, Virginia (Section 5). Finally, we provide results for model runs involving the restoration alternatives that relate to Eastern oysters only (EIS alternatives 1-3; Section 6). Finally, we describe additional model runs to quantify model sensitivity (Section 7).

2.0 Data Sources and Parameter Estimates

2.1 Maryland Department of Natural Resources Annual Fall Survey

The Maryland Department of Natural Resources' (DNR) annual fall dredge survey served as an essential source of data for the demographic model. Data from the survey were used to parameterize growth, recruitment, and survival. Because these data were included in many parameterizations, we describe the survey in general here and refer to specific components as appropriate in the following sections. It is important to note that this survey was not designed for the purpose of developing model parameters, but provided the only comprehensive source of data useful for that purpose. As described for individual parameter estimation procedures, this required that the data be manipulated in a number of different ways. Additional details of DNR's fall survey are described by Tarnowski et al. (2003) and Coakley (2004).

Data collected from representative oyster bars throughout the Bay from 1980 to 2006 as part of the annual survey were used in the demographic model. The survey was conducted in October or early November each year using a standard 36-in wide (91.4-cm) oyster dredge to collect live oysters and shells of dead oysters. Approximately 200 to 400 representative bars

throughout the Maryland portion of the Bay were sampled each year. Bars were divided into several classifications that received varying levels of sampling effort, as described below.

Forty-three bars known as “disease bars” were sampled from 1990 to 2006 to estimate disease prevalence. Disease bars were sampled by combining two ½-bushel subsamples from replicate dredge tows into a composite sample of one Maryland bushel (0.046 m³). All oysters in the sample were counted and measured to the nearest 5-mm shell height (defined as the distance between the shell hinge and the farthest point from the hinge where the two shell halves meet). Spat were also identified morphologically. Spat have one valve that is thinner and narrower than the other (the lower valve if they are oriented parallel to the substrate). The number of articulated shells of dead oysters (known as boxes) was recorded for all bars sampled. A subsample of 30 oysters that were 40 mm or larger from each bar were diagnosed for the presence of Dermo and MSX diseases. We used these data to estimate mortality rates in the demographic model. The data were also included in modeling growth rates and the stock-recruitment relationship.

Fifty-three bars known as “key bars” were sampled each year and used to calculate an annual index of spatfall. Thirty-one of the key bars were also disease bars in the period from 1990 to 2006. Key bars were sampled using the same methods as disease bars, except that no disease sampling was conducted, and shell heights were measured to the nearest 5 mm only at bars that were also disease bars. For the remainder of the key bars, adult oysters captured were measured to the nearest size category as small (41-75 mm) or market-size (≥ 76 mm)¹. Spat were identified morphologically at all bars as described above. These data were used, in part, to model growth rates and the stock-recruitment relationship in the demographic model.

At an additional 95 to 266 bars sampled per year (mean 202), adult oysters were measured to the nearest size category as small (41-75 mm) or market-size (≥ 76 mm), and spat were identified morphologically. These included managed bars where shell or seed had been planted and natural bars where there had been no management. Some of these bars were “seed production” areas that were established to produce seed oysters when a large spat set occurred. If a large spat set occurred, some spat were relocated to bars in other areas with lower disease pressure. Seed-production bars were sampled by pooling five, 0.2-bushel subsamples from replicate tows. All other bars were sampled by collecting one, 0.5-bushel (0.0223 m³) sample. These data were included in the demographic model to estimate the starting population and stock-recruitment relationship.

2.2 Available Habitat

Procedures used to develop the oyster habitat layer are described in detail in a DNR internal report by K. Greenhawk dated December 2005 (Appendix 1).

¹ The size categories of spat, small and market have traditionally been used in oyster surveys in the Bay but do not have consistent biological meaning; while the category of spat always consists of a single yearclass, the small and market categories may include multiple yearclasses of oysters.

The total amount of habitat available for oyster settlement in the Bay was estimated based on historic survey data, records of leased-bottom culture areas, and known oyster restoration sites. Major sources of data included the Maryland Bay Bottom Survey conducted from 1975 to 1983, and an earlier study conducted by Yates from 1906 to 1911. These data were combined in a geographic information system to produce a series of polygons indicating areas that generally contained cultch or hard-bottom substrate suitable for oyster settlement. The amount of habitat has greatly decreased since much of the survey work was conducted in the 1970s and 1980s due to sedimentation. To account for the estimated reduction in habitat, polygons were reduced proportionally but retained the same center location (most were reduced to 29.17% of their original area; Smith et al. 2005).

A total of 8,480 polygons were identified and included as individual oyster bars in the demographic model. Polygons varied greatly in their size, relief, and abundance of oysters. Some polygons consisted of smaller areas of shell that were aggregated but had some unsuitable habitat between them. Although not all polygons resembled classical three-dimensional reef structures, we refer to them consistently as “bars” throughout this manuscript.

2.3 Starting Population

Procedures used to estimate the starting population in Maryland are described in detail in two DNR internal reports: one by K. Greenhawk and T. O’Connell dated June 2007 (Appendix 2), and one by L. Barker dated 14 March 2007 (Appendix 3).

The abundance, distribution, and size structure of the initial population of oysters at the start of all scenarios was based on data provided by MDNR for Maryland, and by R. Mann (Virginia Institute of Marine Science) for Virginia, for the year 2004. In Maryland, the densities of small and market-size oysters were estimated for each basin from bars sampled in DNR’s fall dredge survey (Section 2.1), as described by Greenhawk and O’Connell (2007; Appendix 2). Size-specific data obtained from the nearest disease bars sampled in the survey were used to estimate the size structure (by 5-mm size class) and proportion of spat, by National Oceanic and Atmospheric Administration (NOAA) regions. After size frequencies and densities were calculated for each NOAA region, total populations were calculated by expanding densities for each 5-mm size class by the estimated habitat available (Section 2.2).

An important limitation of the DNR survey is that the area swept by dredge tows was not recorded historically because the survey was not designed specifically to estimate population size. The area swept is needed to estimate relative or absolute density. The DNR conducted a study in 2001 and 2002 to estimate the mean tow area needed to collect one bushel of oysters (Appendix 2). Based on the study, DNR calculated densities assuming mean areas swept of 78 m²/bushel and 10% dredge catch efficiency for all dredge tows in all years. We considered these data to be suitable for modeling the starting population in Maryland with unestimated variance. However, this limitation makes the data unsuitable for analysis as a time series of historical abundance, and the data could not be used for model validation.

The starting population for Virginia was estimated based on surveys and data synthesis conducted by the Virginia Institute of Marine Science, and published by the Chesapeake Bay Program (Virginia Institute of Marine Science 2004 Chesapeake Bay Oyster Population Estimation; <http://www.vims.edu/mollusc/cbope/vabasin.htm>; and R. Mann and J. Harding, pers. comm.). Estimates were made for each basin based on a combination of fishery-independent survey data using patent tongs, fishery-dependent data, surveys of restoration efforts, and aquaculture data. The estimated total number of oysters was allocated to bars in the habitat layer (Section 2.2) in proportion to bar size. The size distributions (5-mm classes) were estimated differently depending on the basin because the amount of data varied. For the James River, we used the observed size-distribution data for 2004. For the Great Wicomico, Rappahannock, and Piankatank rivers, we used the average size distribution off all three rivers in 2004 because individual sample sizes were relatively low. For the Potomac River, Eastern Shore, and Tangier areas of Virginia, we used the average size distribution from the Maryland portions of these basins because few samples were taken in Virginia. For the York/Mobjack, Lynhaven, and Poquoson/Back basins, we used a size distribution based on unpublished data provided by R. Mann and J. Harding. Similarly to the Maryland data, we considered the Virginia data to be suitable for use as a starting population but inappropriate for use in additional model validation. This is because the precision and accuracy of the data from some of the input sources could not be estimated.

The starting population was estimated to be larger in Virginia than in Maryland (Table 1). The number of market size oysters was only about 35% greater in Virginia, but the number of small oysters was estimated to be nearly ten times greater, and the number of spat was estimated to be more than 45 times greater. The starting population is depicted graphically in Figure 1, and described in greater detail in Appendix 2 (Maryland), and at <http://www.vims.edu/mollusc/cbope/vabasin.htm> (Virginia).

2.4 Flow Regime

2.4.1 *Weather years*

The annual flow regime was entered into the model as a categorical variable called “weather year,” with three levels: dry, average, or wet. Weather years were used to adjust salinity (Section 2.5), intensity of disease (Section 2.9), and freshets (Section 2.4.2) on individual bars. Weather years also had cascading effects via salinity in the model, because salinity affected recruitment (Section 2.6), and natural mortality rates (Section 2.10).

Weather years were classified based on annual discharge into the Bay from 1937 to 2005, as reported by the U.S. Geological Survey (USGS) Chesapeake Bay Activities (Figure 2; <http://md.water.usgs.gov/monthly/bay.html>). Years when annual inflow was in the 25th to 75th percentile were defined as average. Years when inflow exceeded or failed to reach this range were classified as wet or dry years, respectively. Weather-years were simulated in the demographic model by randomly selecting a ten-year block from the historical data. A starting

year from 1937 to 1996 was selected randomly, and the type of weather for that year and the next nine were used for years 1 through 10 in the model.

Years were selected in blocks because the climate historically has tended to shift between wet and dry conditions over several years. That is, wet or dry years tended to occur in clusters through time; however, rainfall patterns have shifted between wet and dry years more randomly during the last ten years. These unpredictable changes in climate may become more prevalent as global temperatures increase (U.S. Global Change Research Program 2000). An additional set of model runs was conducted to quantify sensitivity to weather patterns in which the type of weather was selected randomly for each year, and the probability of selection was equal to the proportion of each type of weather-year in the historical data (25.37% dry, 52.24% average, and 22.39% wet). These additional runs are reported in Section 7, Model Sensitivity.

2.4.2 Freshets

Freshets severe enough to kill oysters were simulated for individual bars in the demographic model. Freshets occurred only during wet years on bars that were predicted to have low salinity (Section 2.7). During a wet year, summer freshets occurred at random with an 8.3% (1/12) chance on low-salinity bars, and winter freshets occurred with a 16.7% (1/6) chance. If a winter freshet occurred, 5% of the oysters in all size classes were removed from the affected bar for the current year. If a summer freshet occurred, 65% of all oysters were removed. Because a complete historical record of freshet occurrences is not available for all portions of the Bay, these rates were estimated based on the data and recollection of a group of regional researchers at a working group meeting held 25 October 2005 at Horn Point Laboratory, Maryland. We note that freshets have the same average effect on all alternatives simulated; therefore, freshets are not expected to introduce a bias into the comparison of alternatives, even if their relative frequency of occurrence was not estimated precisely.

2.5 Salinity

The salinity submodel was provided by P. Genovese MDNR in the form of spreadsheet files.

Salinity was assigned to each bar for a model year based on extrapolation of measurements taken by the Chesapeake Bay Program at monitoring stations throughout the Bay during 1995 to 1999 (Figure 3). These included two dry years (1995 and 1999), two wet years (1996 and 1998), and one average year. The MDNR used this range of weather-years to predict the effects of freshwater inflow on temporal variation in salinity at individual bars. A salinity class was assigned to each bar in the model for each weather-year based on the nearest monitoring station, or by interpolating between stations by kriging. Bars were assigned to one of three salinity classes as low (< 11 ppt), medium (11-15 ppt), or high (\geq 15 ppt). These ranges coincide with biologically important thresholds for changes in oyster survival and recruitment rates reported in the scientific literature (e.g., Calvo et al. 1999, 2001; Grabowski et al. 2004; Jordan et al. 2002). Eastern oysters generally do not live in areas where salinity is less than 5 ppt

for extended periods (Shumway 1996). We assigned salinities to bars for a simulated year in the demographic model by randomly selecting the 1995 or 1999 salinity pattern if the year was dry, and the 1996 or 1998 salinity submodel if the year was wet. During average weather-years, bars were assigned salinity classes based on the 1997 salinity submodel.

2.6 Recruitment

2.6.1 Recruitment Data Sources

Recruitment of spat onto oyster bars was modeled using the empirical relationship between the relative numbers of spat counted in DNR's annual fall survey (Section 2.1) and the estimated number of sexually mature female spawners from counts of market-size oysters. The relationship was modeled using data collected between 1980 and 2006, after yearly reproduction and spatfall. To avoid bias, the data were limited to bars that were not seeded with spat from elsewhere or planted with shell during the last five years to augment production. Within this set, oyster heights were measured with different precision depending on the kind of bar sampled, as described in Section 2.1. Data were available for 41 to 62 bars per year (mean 51) where oyster heights were measured to the nearest 5-mm height class (disease bars from 1990-2004), and 95 to 266 bars per year (mean 202) where height was measured to the nearest size category (spat 0-40 mm, small 41-75 mm, or market ≥ 76 mm).

Two additional relationships obtained from the literature were used to model the stock-recruitment relationship. First, the ratio of female to male oysters was assumed to increase with mean size of oysters as described by Kennedy (1983) and presented in Table 2. This relationship reflects the fact that oysters are hermaphroditic and begin changing from males to females as they grow (Kennedy 1983). Second, the number of eggs attributable to female oysters of different size classes was based on the relationship

$$(1) F_{ind} = 39.06 \times (0.000423 \times H^{1.7475})^{2.36},$$

where F_{ind} is the fecundity (number of eggs produced) of individual females in millions, and H is shell height in millimeters (Mann and Evans 1998).

2.6.2 Recruitment Model

Stock-recruitment relationships in fisheries typically project numbers of recruits as a function of numbers or biomass per unit of area (e.g., Quinn and Deriso 1999). For this model, absolute abundance or biomass of oysters by size could not be estimated from the survey data (Section 2.3); thus, a typical stock-recruitment approach could not be employed. Because of this data limitation, we modeled recruitment based on the ratio of spawners to spat measured in the survey. First we verified that relative numbers of spat and adult oysters could reasonably be estimated from dredge surveys based on the scientific literature. Chai et al. (1992) compared the sampling efficiency of dredge surveys and diver-harvest surveys using a replicated study in the Chesapeake Bay. They found that density estimates from dredge surveys were only 2-32% of the

diver estimates (Figure 4A). However, the relative numbers of spat, small, and market-size oysters were similar between the two survey methods (Figure 4B). The site-by-site comparison was not strictly valid because the dredge survey sampled a significantly larger area from each bar than the diver-harvest survey. Despite this limitation, the study suggests that there is no consistent bias in the ratios of spat to small or market-size oysters derived from dredged samples. That is, spat do not appear to be captured disproportionately more or less frequently than larger oysters in a dredge. We proceeded to model using the ratio of spawners to spat in fall survey data, which was the only information available at a broad spatial scale with which to model the stock-recruitment relationship.

The ratio of spat to female spawners measured in DNR's survey was not directly comparable across bars because fecundity increases non-linearly with oyster size (Mann and Evans 1998). Therefore, the number of spat per spawner depends on the size structure of the population. One method of dealing with this problem is to calculate a stock-recruitment relationship between the biomass of the spawning stock and corresponding number of recruits (Quinn and Deriso 1999). This method incorporates information about the size structure of the population because biomass is a function of the number of animals present and their sizes. Although biomass was not measured directly in the DNR survey, it could be estimated using an empirical relationship between oyster height and weight. The relationship listed in Equation 1 includes a conversion from height to weight, and an additional conversion from weight to estimated fecundity for an oyster of a given size. We assumed that the size-specific fecundity relationship, as determined from laboratory experiments (Mann and Evans 1998), would provide a better estimate of an oyster's contribution to the number of recruits than an estimate based on biomass alone. A similar fecundity relationship will also likely be the only information available about the stock-recruitment relationship for Suminoe oysters. Our strategy for modeling the stock-recruitment relationship was to adjust the estimated number of female spawners using the egg-size relationship in Equation 1 so that they were expressed as the number of females of a standard size that would produce the equivalent number of eggs. Numbers of females were converted to equivalent numbers of standard-size females using size-specific estimates of fecundity as a common currency. In this way, the egg-size relationship was used to account for differences in sex ratios and size distribution among bars and years, although eggs themselves were never actually quantified. The number of spat enumerated in the survey was then modeled as a function of the number of standardized female oysters to predict spat production in the demographic model.

Oyster recruitment is also strongly affected by, and generally positively related to salinity (Ulanowicz et al 1980; Tarnowski et al. 2005). Therefore, we developed the recruitment submodel and estimated recruitment in the demographic model by salinity class. Numbers of sexually mature female oysters were standardized for each year and salinity class (low, medium, or high) using the following procedure. First, means were calculated for the number of eggs produced by market-size and small oysters for the subset of bars where 5-mm height classes

were sampled (disease bars). The number of female oysters present was estimated for each bar and height class by multiplying the total number of oysters counted in the sample by the appropriate sex ratio listed in Table 2:

$$(2) Nf_{bc} = N_{bc} \times R_r,$$

where Nf is the number of sexually mature female oysters; N is the total number of oysters; b is bar; c is 5-mm size class; R is the sex ratio; and r is the class listed in Table 2.

The number of eggs produced was calculated by bar and 5-mm size class using Equation 1; the midpoint of the height class was entered as the size of all oysters in the class (e.g., 42.5 was entered as the size of oysters in the 40-45 mm height class):

$$(3) F_{bc} = Nf_{bc} \times F_{ind\ bc},$$

where F_{bc} is the total fecundity for a bar and 5-mm size class.

The mean number of eggs produced by a small (\bar{F}_{small}) or market-size (\bar{F}_{market}) oysters for each year and salinity class was calculated as

$$(4) \bar{F}_{small} = \frac{\sum_{b=1}^B \sum_{c=42.5, 47.5, \dots}^{72.5} F_{bc}}{\sum_{b=1}^B \sum_{c=42.5, 47.5, \dots}^{72.5} Nf_{bc}}, \quad \text{and} \quad (5) \bar{F}_{market} = \frac{\sum_{b=1}^B \sum_{c=42.5, 47.5, \dots}^{72.5} F_{bc}}{\sum_{b=1}^B \sum_{c=77.5, 82.5, \dots}^{90} Nf_{bc}}$$

For years when no data were recorded by height class (1980-1989), average numbers were calculated for market-size and small oysters using the same procedure, except that numbers were also summed across years. Because the subset of bars sampled by 5-mm height classes was spread throughout the Bay, these numbers were approximately representative of the size distribution of oysters within the larger categories of small or market-size for a year and salinity class.

After mean fecundities were calculated, Equations 4 and 5 were used to extrapolate the total number of eggs produced for all bars sampled in a year and salinity class, not just those where shell heights were measured to 5-mm height classes. This was done to obtain mean fecundities for a year and salinity class that better reflected the size structure for the entire Bay, because many more bars were sampled where measurements were recorded only to the level of small or market sizes. First the numbers of small and mature female oysters were calculated for all bars sampled using Equation 2, except that we denote the results as Nf_{small} and Nf_{market} , respectively. The sums of these numbers were multiplied by the mean fecundities for small and market-size oysters and added to obtain total fecundity for a year and salinity class (F_{tot}).

$$(6) F_{tot} = \bar{F}_{small} \times Nf_{small} + \bar{F}_{market} \times Nf_{market}$$

The total number of eggs produced by market-size and small oysters was then divided by the number of eggs produced by a 77-mm oyster (Equation 1) to estimate the equivalent number of standardized female oysters present in a year and salinity class. We selected 77 mm as a standard size because the 75-mm to 80-mm size class was observed most frequently in the data set when data for all years were pooled:

$$(7) N_s = \frac{F_{tot}}{25.7023},$$

note that $39.06 \times (0.000423 \times 77^{1.7475})^{2.36} = 25.7023$, Equation 1.

It is well documented that episodic large oyster spat sets periodically occur in the Bay (e.g., Ulanowicz et al. 1980; Tarnowski et al. 2005). This phenomenon was accounted for in the recruitment submodel in the following manner. The number of spat produced was calculated as the number of standardized females multiplied by a ratio of spat to standardized female spawners (N_s). The specific ratio used was selected randomly from a submodel that reflected either a typical recruitment year or an episodic large spat set. Unusually large spat sets were identified in the survey data by calculating which spat-to-spawner ratios exceeded the 75th percentile of all of the data by more than 1.5 times the interquartile range (i.e., outliers on a boxplot; Emerson et al. 1983). This threshold was 3.0 spat per female spawner (Figure 5). Of the 78 total observations, 58 were in medium or high salinity, and 10 of these were outliers with spat-to-spawners ratios that exceeded 3.0. No outliers occurred in low salinity. Therefore we modeled episodic large spat sets for a year in the model with a probability of 10/57. If a large spat set occurs, the number of spat on bars located in medium or high salinity is calculated by multiplying the number of standardized females by a ratio that is randomly selected from one of the ten extreme values (3.30, 3.50, 3.60, 3.70, 4.00, 5.08, 5.11, 7.30, 7.93, and 8.90). Large spat sets never occur at low salinity bars in the model because they were never observed in the survey.

If a large spat set is not predicted to occur for a year (probability 68/78), numbers of spat are predicted based on spat-to-spawner ratios with the outliers removed. These ratios were modeled as functions of salinity with unequal variance among salinity classes. The resulting mean ratios from the categorical model were 1.44 (SE = 0.195) spat per female spawner in high salinity, 0.47 (SE = 0.082) in medium salinity, and 0.08 (SE = 0.018) in low salinity. All three estimates were statistically significant with $P < 0.001$. Ratios of spat to spawners are randomly selected for normal recruitment years in the model based on these relationships. A random residual is selected from the standard normal curve for each basin and year, and used to select a spat-to-spawner ratio for each bar based on the bars salinity:

$$(8) N_{spat} = N_s \times (Ratio_s + SE_s \times Residual),$$

where N_{spat} = the number of spat produced on a bar in the model, $Ratio$ is the mean ratio of spat to female spawners predicted for salinity class s (1.44, 0.47, or 0.08), SE_s is the appropriate standard error for salinity class s , and $Residual$ is the randomly selected residual from a standard

normal curve. Newly recruited spat were assigned to 5-mm height classes in proportion to the occurrence of spat in normal distributions of age-0 oyster sizes, as modeled for the growth portion of the demographic model (Section 2.7).

2.7 Larval Transport

Spat were distributed among some of the modeled bars according to the larval-transport model reported by North et al. (2006; *In press*). The larval-transport model couples a hydrodynamic circulation model with information about the vertical swimming behavior of Eastern and Suminoe oysters to predict the proportions of spat that will set on each bar. The transport model run was for conditions that occurred in the Chesapeake Bay between 1995 and 1999. These years were selected to include a broad range of physical conditions such as wind, tributary inflow, and annual precipitation (Section 2.5). Separate runs were made for Eastern oysters and Suminoe oysters to reflect the fact that Eastern oyster larvae tend to swim up in the water column in response to salinity gradients, but Suminoe oyster larvae tend to swim down (Newell et al. 2005). The larval-transport model included predictions for 2,776 of 8,480 bars in the demographic model. However, that subset of bars accounted for 64% of the total habitat acreage in the habitat layer. The modeled predictions were incorporated into the demographic model as a $2,776 \times 2,776$ matrix of bars for each species and year simulated. Cells in the matrix indicated the number of simulated particles (representing oyster larvae) that were transferred from bars listed in columns of the matrix to bars listed in rows. For bars that occurred in the demographic model but not the larval-transport model, spat were assumed to be distributed evenly within the basin where they were located, in proportion to the bar's area. Larval-transport data were integrated into the demographic model by converting numbers in the matrix to proportions and allocating the estimated number of spat produced by each bar in the demographic model according to these proportions. For an average weather-year in the demographic model, the matrix for year 1997 was used to allocate spat. For dry or wet years in the demographic model, one of the two matrices for a corresponding weather-year was selected randomly with equal probability (1995 and 1999 dry, 1996 and 1998 wet).

2.8 Growth

Oyster growth rates were modeled in yearly increments as changes in shell height. The general strategy for modeling growth was to calculate a series of von Bertalanffy growth functions (von Bertalanffy 1938) and to select one of the growth functions for each simulation, with a probability based on an empirically measured mean annual growth rate and variability about that mean.

2.8.1 Growth Data Sources

Growth of Eastern oysters was modeled based on four sets of empirical data. Three of the sets were used to develop a model for use in Maryland waters and the Potomac River, and one set was used to create a model for Virginia waters, except for the Potomac River. This was

because the empirically-derived growth rates for the upstream portion of the Bay in Maryland were greater than the empirically-derived growth rates for Virginia. We were informed that growth rates for basins other than the Potomac River in Virginia were similar to the rates in the James River (R. Mann, unpublished data and pers. comm.). Although some laboratory studies indicate that Eastern oyster growth rates are higher in high salinities than in lower salinities, growth rates in this model were not adjusted for salinity. This was because no significant differences were found among salinity classes in the empirical data from which the growth functions were developed. As a result, the trend in growth versus salinity employed here was opposite that found by other researchers; mean growth was slightly greater in low salinity than in high salinity. Such a discrepancy could be accounted for by site-specific environmental factors other than salinity.

The data sources were

(1) Coakley data – 29 bars in Maryland sampled as part of DNR’s annual survey from 1990 to 2001 were analyzed by Coakley (2004). Heights-at-age were determined for these data using maximum-likelihood modal analysis on height frequencies. Mortality rates of larger oysters were relatively high because these data included many bars with high disease prevalence and some that were heavily fished (Coakley 2004). This caused fitted models of growth to be much lower than actual growth when ages 3 and greater were included in the data because the larger oysters of these ages were no longer represented; therefore, we restricted this data set to ages 0 to 2 in our model.

(2) Paynter data – 27 bars in Maryland that have been managed as a restoration project from 1998 to the present for the U.S. Army Corps of Engineers have been monitored by the University of Maryland Center for Environmental Science. Data were available for the period 1998 to 2004 (courtesy of K. Paynter, UMCES). Disease prevalence was low on these bars during the time period (although it was high elsewhere in the Bay), and no fishing occurred. Oysters were planted at known dates and measured four times per year to obtain growth estimates.

(3) Rothschild data – These data were obtained from seven studies where oysters of known age were planted in the Bay, primarily in Maryland and the central portion of the Bay. The studies were reported in the literature between 1950 and 1966 and summarized by Rothschild et al. (1994). Specifically, they were Beaven (1950 and 1953), Butler (1953), McHugh and Andrews (1955), Shaw (1966a and b), and Shaw and Merrill (1966). The data were restricted to studies that occurred before disease was prevalent in the Bay and where the authors reported no unusual mortality; consequently, growth rates reported in these studies are unlikely to be biased by differential mortality among size groups.

(4) Mann data – The Virginia model was based on a single data set from 17 bars in the James River that were monitored by the Virginia Institute of Marine Science from 1995 to 2003 (courtesy of R. Mann, VIMS). Age classes were determined through visual identification of height-frequency modes. Disease prevalence was relatively low, and the bars received little or no fishing pressure; therefore, these data likely reflect the true growth rate of oysters in the James River and similar areas of the lower Bay.

2.8.2 *Growth Model*

Each data set provided measurements of mean height-at-age for a particular bar and year (Figure 6) that were assumed to come from a random sample of size-at-age data from all possible bars and years in Maryland or Virginia. Height-at-age was assumed to be normally distributed with an estimated mean equal to the mean height for that age and state, and standard deviation equal to the standard deviation of the same data. Our strategy was to develop a series of growth functions that reflected the spatial and temporal variability observed by fitting functions through a range of percentiles of these normal distributions at age, and selecting one of the resulting growth functions randomly (with replacement) for each bar and year in the demographic model.

For each state, growth was modeled as a function of age using the von Bertalanffy growth function:

$$(9) H_t = H_\infty(1 - \exp(-K(t - t_0))),$$

where H_t is the shell height at time t (a particular age); H_∞ is the asymptotic or maximum height that an average oyster is expected to achieve; K is the Brody growth coefficient (cf, Quinn and Deriso 1999); and t_0 is the time of oyster settling.

First, the mean height and standard deviation were estimated for each age 0 to 30 (the estimated maximum age an oyster could live; Kennedy et al. 1996). The mean height was calculated for ages 0 to 4, and the von Bertalanffy function was fit to these means using the non-linear, least-squares procedure of the ‘nlme’ library (Pinheiro et al. 2006) in the statistical software package R (Version 2.4.0; R Development Core Team 2006). Data for ages 5 and 6 were not used because too few points were available to estimate variance reliably. The mean heights at ages 5 to 30 then were estimated as the fitted values from the resulting function. Mean H_∞ was fixed at 177 mm for both the Maryland and Virginia models. This was the mean of all values reported in the studies summarized by Rothschild et al. (1994). We did not attempt to estimate H_∞ from the recent data sets because they did not include any very large, older oysters that would be needed to make the estimate without extrapolating well beyond the range of the data. The standard deviation of heights at ages 5 to 30 and H_∞ were estimated from the coefficients of variation for ages 0 to 4. The mean coefficient of variation for ages 0 to 4 (22% for Maryland and 17% for Virginia) was multiplied by the mean height at ages 5 to 30, and by H_∞ , to obtain their estimated standard deviations. This approach assumes that variance increases

in proportion to height, a pattern that commonly occurs for growth of fish and shellfish (Quinn and Deriso 1999) and fits the observed data well (Figure 6).

After the set of means-at-age and associated standard deviations was calculated, a set of 999 growth functions was generated for each state by fitting the von Bertalanffy function through the 0.1th to 100th percentiles (by 0.1) of each normal curve. For example, the growth function for the 50th percentile was calculated by fitting a model through the mean at ages 0 to 30 and H_{∞} of 177. Growth then was simulated by randomly selecting one of the 999 functions, with replacement, for each basin and year and applying the curve to all bars in the basin.

An additional adjustment of the Paynter data set was required before those data could be integrated with the other data for Maryland. The other Maryland data were based on surveys taken during the fall; therefore, all heights-at-age could be combined to estimate a single normal distribution. That is, all oysters of a particular age were measured at about the same time – at the end of the annual growing season. In the Paynter data set, height measurements were taken at different times throughout the year. To standardize Paynter’s data for use in the normal distributions, the von Bertalanffy function was fit to the data, and the modeled values at rounded age increments were used in the distributions (e.g., at ages 0, 1, 2, etc. at the end of the growing season).

Growth was simulated in the demographic model by randomly selecting a growth curve for a year and basin as described above and adding the difference in shell height that oysters in a 5-mm size class were calculated to have one year after growth along the curve, at the midpoint of the height class. This was necessary because the demographic model is not cohort based. Oysters are accounted for as the number on each bar by 5-mm height class. Thus, all oysters in a height class grow in lock step. A maximum size class of 320 to 325 mm was imposed to reflect the largest size that any oyster would be likely to attain (Kennedy et al. 1996; this differs from H_{∞} [177] in that H_{∞} is a mean maximum size with an associated distribution).

Biomass was estimated from the number of oysters in each size class using the conversion

$$(10) \text{ Biomass (g carbon)} = 0.0002115 \times H^{1.7475}$$

This relationship was obtained by dividing the conversion from shell height to dry weight reported by Mann & Evans (1998) by two to estimate grams of carbon (C. Cerco, U.S. Army Corps of Engineers, pers. comm.).

2.9 Disease Intensity

Individual oyster bars were ranked according to mean intensity of Dermo disease based on the observed frequency of diseased Eastern oysters at ‘disease bars’ surveyed by DNR annually from 1990 to 2005. Subsamples of 30 oysters from each of these bars were examined for Dermo and MSX diseases by the Sarbanes Cooperative Oxford Laboratory, Oxford, MD. Each bar was ranked as fitting into one of three ‘tiers’ of Dermo intensity. Tier 1 indicates the

greatest intensity, and Tier 3 the least. Tiers reflect concentrations of pathogen in hemolymph or solid tissue (Gieseke 2001) of greater than 2.85 (Tier 1), 2.0 to 2.85 (Tier 2), or less than 2.0 (Tier 3). These boundary values indicate statistically significant break points in disease prevalence during the period, as determined using a multiple comparison based on Friedmans's Rank Sum Test (Tarnowski et al. 2003).

In the demographic model, a disease tier was assigned to a bar as a random function with probability equal to the observed distribution of mean disease intensity in DNR's survey (Table 3). Dermo was more likely to occur at high intensity (Tier 1) in dry years than in wet years. Disease tiers were assigned to bars independently of the previous year's tier. The effects of Dermo were included in the model by increasing natural mortality rates as Dermo intensity increased to match mortality rates calculated using DNR's survey data (Section 2.1).

The effects of MSX disease on eastern oysters were included in the model only if two or more dry years occurred in a row (based on data from M. Tarnowski, MDNR, personal communication). If two dry years in a row were encountered during simulation, MSX events were assigned to bars at random with a probability of 0.38 for low salinity bars, 0.71 for medium salinity bars, and 1.00 for high salinity bars. A bar's status in the previous year did not affect the probability of being affected by MSX. These probabilities were derived from a study of the Choptank River, Maryland, as described by Vølstad et al. (In Press). The effects of MSX were modeled by increasing natural mortality rates as described in Section 2.10.

2.10 Natural Mortality

Procedures used to estimate natural mortality are described in greater detail in Appendix 4, an article by J. Vølstad, J. Dew, and M. Tarnowski that has been accepted for publication in the Journal of Shellfish Management.

We estimated natural mortality rates using data from DNR's fall oyster survey from 1991 to 2006 (200-400 bars per year; Section 2.1). Estimates were based on ratios of live oysters to articulated shells of oysters that had died recently (i.e., boxes; Southworth et al. 2005; Ford et al. 2006). Boxes were categorized as 'recent' if they contained tissue, or if there was no fouling or sedimentation on the inner surfaces of the valves. The general strategy for estimating mortality was to pool data across bars within the same size class (small or market), salinity zone, and measured disease intensity to achieve sufficient sample sizes, and then use recent box counts to estimate annual proportional mortality rates (Ricker 1975).

A key parameter of this model is the average time that boxes are assumed to remain in the recent category, which serves as an estimate of the time since death (TSD). This is because TSD is used to calculate proportional annual mortality. A shorter assumed TSD would correspond to a greater estimated mortality rate because recent boxes would be interpreted to have died in a shorter period. Vølstad et al. (In Press; Appendix 4) estimated the mean time during which a box would be classified as recent in the Chesapeake Bay to be one to two weeks

after death. They obtained this estimate by comparing mortality rates calculated using different sedimentation times with interannual patterns in salinity and disease intensity that are known to affect mortality rates. That is, estimated mortality rates increased in years when salinity and disease intensity were measured to be higher than average in the Bay, and vice-versa. To account for the uncertainty in time that oysters were categorized as recent, we calculated pooled mortality estimates for one to two weeks TSD. Pooled mortality rates were calculated from one-week and two-week estimates by combining bootstrap resamples from each type of estimate, as described below.

We calculated mean annual mortality rates for each size category (small or market-size), salinity zone j , and disease tier k , based on counts of live oysters and boxes from all bars sampled in the group over time ($i = 1, \dots, n_{jk}$), using the following equations:

$$(11) \quad s_{jk} = \frac{\sum_i live_{ijk}}{\sum_i (live_{ijk} + newbox_{ijk})},$$

$$(12) \quad m_{jk} = -\log(s_{jk}),$$

$$(13) \quad totm_{jk} = 1 - \exp^{-m_{jk} \times T}$$

where s_{jk} is the 1- or 2 -week survival rate; $live_{jk}$ is the number of live oysters; $newbox_{jk}$ is the number of recent boxes; m_{jk} is the instantaneous (1- or 2 -week) mortality; T is the expansion factor to the total number of weeks (20) during which natural mortality occurs ($T = 20$ for 1-week mortality, and $T = 10$ for 2-week mortality); and $totm_{jk}$ is the mean annual mortality rate.

These calculations correspond to instantaneous mortality rates (Equation 12) and annual proportional mortality (Equation 13) rates, as described by Ricker (1975). The ratio estimator (Equations 11) provides weighted-mean mortality across bars in which weights are proportional to the number of live and dead oysters at each bar. This was necessary because the distribution of mortality estimates for individual bars was highly skewed by catches with no boxes (i.e., mortality estimates of zero), so pooling was necessary to obtain means over all bars in a salinity class and disease tier. We assumed that the instantaneous mortality rate was constant from June through October and that most oyster mortality occurred during this period; therefore, the expansion factor, T , was set to calculate annual mortality for a 20-week period, approximately from June 1 to October 31. This is because most predation and disease mortality occurs in summer, and overwinter mortality rates are believed to be negligible in comparison (cf., Vølstad et al. In press, and references therein).

We calculated mortality rates from Equations 11 and 13 and associated variances by bootstrapping (Efron and Tibshirani 1993). We ran 1,000 bootstrap resamples with one-week

TSD, and 1,000 resamples with two-weeks TSD. Each resample consisted of n_{jk} randomly selected observations (with replacement) where n_{jk} was the number of samples collected across all bars over the time series of fall surveys, in the respective salinity zone and disease tier. We then pooled the two bootstrap distributions to obtain 2,000 resamples for each estimate of mean mortality. The standard errors for the annual mean mortality rates were estimated directly from the distribution of the bootstrap estimates and are reported in Table 4. These results matched the expected trends in mortality with changes in disease intensity and salinity, indicating that the estimates were credible (Figure 7; Vølstad et al. in press).

Mortality rates for Eastern oysters were selected by bar and size category by selecting the appropriate rates for the bar's salinity class and disease tier (Table 4). Variation in mortality rates was simulated by adding a random residual selected for each basin and year to the estimate. This was calculated by selecting a number from the standard normal distribution and multiplying by the standard error associated with the estimate.

Spat were assigned the same mortality rates as small oysters on all bars. This is because estimates were made for fall, after most of the mortality for spat was expected to occur. The next period when appreciable mortality would likely occur would be the following spring through fall, when most spat from the previous year would be reaching small sizes.

2.11 Harvest

Eastern oyster harvest rates were estimated by dividing statewide reported landings in Maryland (courtesy T. O'Connell, DNR) by the statewide population estimates (Section 2.2) for years 1994 to 2004. On average, 47% of all market-size oysters were estimated to be harvested during this period. Confidence intervals could not be calculated for the population estimates, and reported landings sometimes exceeded population estimates for individual tributaries; therefore the estimated average harvest rate of 47% probably has a large, but unknown, variance. To bound the effects of harvest, we applied a range of harvest mortalities. We conducted model runs with harvest rates of 20%, 40%, 60%, and 80% of all market-size oysters on bars that were not in sanctuaries or protected areas. Harvest rates were modeled as the proportion of the market-size population present at the end of the fall that is caught during the winter harvest season. This corresponds to the exploitation rate (F ; Ricker 1975) for the winter, after natural mortality for the calendar year is assumed to occur (Section 2.10). For runs that included harvest, some small oysters also were removed to reflect incidental harvest. The number of small oysters removed was equal to 10% of the total catch (Powell 2005). We assumed bars that could not be harvested legally (e.g., sanctuaries and closed areas) were respected completely (i.e., the model does not account for illegal harvest). Sanctuaries and closed areas composed 29% of all bars in the model, 23% of the total area of bars modeled, and 22% of the starting population of market size oysters.

2.12 Habitat Improvements

EIS alternatives 1 and 3 include restoration activity at the current level. Alternative 2 includes enhanced restoration, as specified by MDNR (details are presented in Appendix E). The model simulates the effects of restoration in two ways. First, spat planted on bars are added to the population after natural recruitment and larval transport are calculated. Planted spat are assumed to experience 50% mortality between the time they are planted and the end of the growing season for that year (C. Judy, DNR, pers. comm.). Their size distribution is assumed to be the same as that of their natural counterparts. The second method of habitat improvement incorporated into the model is shell planting, which improves recruitment of spat by creating additional substrate for settlement. This improvement is gradually lost as planted shell becomes covered with sediment. We modeled this relationship based on empirical data provided by DNR (C. Judy, pers. comm.) and published data describing the rate at which planted shell is inundated with sediment in the Chesapeake Bay (Smith et al. 2005).

The initial improvement in recruitment after planting shells was estimated from MDNR's data based on 155 samples of spat density (spat/bushel from standardized tows as an index to density) collected from 1994 to 2004 on bars that had been replenished with shells earlier in the same season. As a control, the density of spat was sampled at one or more bars that were located within 1.6 km (1 mile) of each replenished bar and that had not been replenished within the last five years. First the mean densities of adjacent unreplenished control bars were averaged to provide a single number for paired comparisons with each replenished bar. Then overall mean densities were calculated for replenished bars and unreplenished bars (i.e., the mean for unreplenished bars was a mean of means). A ratio of the resulting means was calculated to estimate the improvement in spat settlement that was associated with repletion of bars, and this ratio was used as a scaling factor for recruitment. On average, the index of spat density was 4.5 times greater on bars that had been replenished (219.3 oysters/bushel) than on comparable bars that had not been replenished within five years (48.6 oysters/bushel). We verified that the increase was statistically significant using a paired *t*-test ($t = 10.4$, $df = 154$, $p < 0.01$; data were transformed to $\log_e + 1$ to meet the assumption of equal variance among treatments).

The decline in condition of replenished bars over time was modeled using the function described by Smith et al. (2005):

$$(14) \textit{Shell condition} = 1.593 + [0.939 \times \log_e(\textit{age})],$$

where shell condition was categorized into five levels. Condition-class 1 was the best condition, indicating that no sediment was present on the shells. Condition-class 5 was worst, indicating that shells were buried with sediment completely (after about 35 years). We assumed that condition-class 1 corresponded to newly replenished bars and that condition class-3 (i.e., shell cups and interstitial spaces sedimented, but entire shell outlines visible) corresponded to the condition of average unreplenished bars in the model (cf., Table 2 of Smith et al. 2005). We

therefore used the function to model the effect of shell planting for shell conditions 1, 2, and 3 only.

We applied the rate of decay in Equation 14 to MDNR's data to model how the expected improvement in spat settlement at replenished sites (a scaling factor in the model) degrades back to the baseline level (i.e., a scaling factor of unity) at unreplenished sites over time. Replenished sites that follow this function degrade non-linearly from about condition-class 1 to condition-class 3 by year 5 (Figure 8). The same rate of decay in habitat improvement was applied to DNR's data to estimate the recruitment scaling factor for each year after replenishment. Replenished bars were assigned a scaling factor of 4.5 for the first year to match the observed data, and modeled scaling factors of 3.0, 2.1, 1.5, and 1.0 for years 2 through 5 following the rate of decline in the above function.

The scaling factors were used to adjust the number of spat recruited to replenished bars in the model until replenishment no longer improved spat set above the level of a natural bar. That is, the number of spat predicted from modeling for an unreplenished bar would be multiplied by 4.5 for replenished bars the first year, by 3 in the second year, etc. Replenishment attracted no more spat than natural bars in year 5, as indicated by a scaling factor of 1.

3.0 Model Structure

The demographic model is spatially explicit. It projects abundance of oysters internally by 5-mm size class and individual bar, although results are aggregated to larger spatial scales for output. The model starts with the current state of the population in the fall, either as starting values at the beginning of a simulation or output from the previous year. The steps in modeling oyster dynamics for an individual bar j , year i , and population P (i.e., Eastern oysters or Suminoe oysters) are depicted in Figure 9.

First the type of weather-year is assigned for year i (Section 2.4.1), and salinity is assigned to each bar based on its location and the type of weather-year (Section 2.5). If the simulation includes harvest (Alternatives 1 and 2 for Eastern oysters, Section 4), and the bar may be legally harvested (i.e., it is not located in a sanctuary), a portion of the market-size oysters equal to the estimated harvest rate is removed from the population. A smaller portion of the small oysters also is removed. The portion is calculated based on the harvest rate so that small oysters constitute 10% of the total catch (Section 2.11). All harvest is assumed to occur in the winter before the following growing season. Freshet mortality is then applied to selected bars by removing a portion of all oysters on the bar (Section 2.4.2). Although freshets may occur in the winter or spring-summer, all freshet mortality is applied at the same time in the model. This does not change the model result because all freshet mortality occurs before recruitment is estimated, and freshet mortality is independent of size.

Next a growth model is selected based on the state in which the bar is located, and a growth curve is randomly selected from the model as described in Section 2.8. For each 5-mm

size class, oyster sizes are incremented by the appropriate amount based on the selected growth curve. Growth is assumed to occur during the summer months before spawning.

Disease tier is selected by weather-year (Section 2.9). After disease tiers are assigned to bars, and natural mortality rates are selected, oysters that succumb to natural mortality are removed. Natural mortality is assumed to occur during summer, at the same time as growth (Section 2.10).

Recruitment is calculated after annual growth and mortality are simulated (Section 2.6). If the bar has been planted with shell to improve habitat during the last four years, the estimated number of spat is increased as described in Section 2.12. The final number of spat present on bar *j* is adjusted according to the larval-transport model by moving spat to and from other bars in proportion to the larval-transport matrix (Section 2.7). If the bar is also planted with spat, planted spat are added to the bar but are assumed to experience 50% mortality (Section 2.12).

The population on a bar is then ready for output. Numbers or estimated biomass of oysters is summed by size class and over bars to produce an estimate for a larger area, such as a state or the entire Bay, for one year. The model begins another iteration for the next year, until it has completed year 10, when it exits.

Model predictions are based on 1,000 simulations. We report the median value of the 1,000 runs as the estimated result, and the 5th and 95th percentiles as approximate 95% confidence intervals.

4.0 Description of Proposed Action and Alternatives

The purpose of the actions being evaluated is to establish an oyster population that reaches a level of abundance in Chesapeake Bay that would support sustainable harvests comparable to harvest levels recorded during the period 1920 to 1970. The proposed action is to establish a naturalized, reproducing, and self-sustaining population of *C. ariakensis* in the tidal waters of Maryland and Virginia through introductions while continuing efforts to restore *C. virginica* using best available restoration strategies and stock assessment techniques. The following specific alternative strategies also are being considered (FR 2004):

- **Alternative 1 - No Action:** Continue Maryland's present Oyster Restoration and Repletion programs, and Virginia's Oyster Restoration Program under current program and resource management policies and available funding using the best available restoration strategies and stock assessment techniques.
- **Alternative 2 - Expand Native Oyster Restoration Programs:** Expand, improve, and accelerate Maryland's Oyster Restoration and Repletion programs, and Virginia's Oyster Restoration Program in collaboration with federal and private partners. This work would include, but would not be limited to, an assessment of cultch limitations and long-term solutions for this problem and the development, production, and

deployment of large quantities of disease resistant strain(s) of *C. virginica* (Eastern oyster) for broodstock enhancement.

- **Alternative 3 - Harvest Moratorium:** Implement a temporary harvest moratorium on native oysters and an oyster-industry compensation (buy-out) program in Maryland and Virginia, or a program that offers displaced oystermen on-water work in a restoration program.
- **Alternative 4 - Aquaculture:** Establish and/or expand state-assisted, managed or regulated aquaculture operations in Maryland and Virginia using the native oyster species.
- **Alternative 5 - Aquaculture:** Establish state-assisted, managed or regulated aquaculture operations in Maryland and Virginia using a suitable triploid, non-native oyster species.
- **Alternative 6 - Introduce and Propagate an Alternative Oyster Species (other than *C. ariakensis*) or an Alternative Strain of *C. ariakensis*:** Introduce and propagate in the state-sponsored, managed or regulated oyster restoration programs in Maryland and Virginia, a disease resistant oyster species other than *C. ariakensis*, or an alternative strain of *C. ariakensis*, from waters outside the U.S. in accordance with the ICES 2003 Code of Practices on the Introductions and Transfers of Marine Organisms.
- **Alternative 7** – Establish a naturalized, reproducing and self-sustaining population of *C. ariakensis* in the tidal waters of Maryland and Virginia through introductions beginning in 2005 (or when EIS is completed) but discontinue *C. virginica* restoration efforts.
- **Alternative 8 - Combination of Alternatives**

The demographic model will be used to project oyster abundance for Alternatives 1, 2, 3, and potentially for Alternative 8. These alternatives are modeled by selecting appropriate input parameters for harvest levels, shell planting, and spat planting, as previously described. The specific allocation of spat and cultch within the Bay for alternatives 1 and 2 are listed in Appendix E (an undated report by P. Jones, MDNR). Two plans have been developed for Alternative 2, which we modeled individually as alternatives 2a and 2b. The plans employ the same numbers of planted spat and cultch overall but differ in the numbers and locations of managed sanctuaries. Specifically, seven more bars would be stocked under alternative 2b (total 39) than alternative 2a. Six fewer bars located in low salinity waters would be stocked, but 13 more bars located medium- or high-salinity waters would be stocked (Appendix E). For Alternative 3, all factors except the harvest level were assumed to be the same as for Alternative 1.

The oyster demographic model was developed to support an Ecological Risk Assessment and an Environmental Impact Statement. The underlying concept was that vital parameters of a model developed using the extensive data available for the Eastern oyster could be modified to reflect different vital parameters of the Suminoe oyster that would be established based on findings of extensive research conducted on that species over the past four to five years. The model could then be run to project potential outcomes of the proposed action. A draft model documentation report that included suggested approaches to modeling the Suminoe oyster was reviewed by the Oyster Advisory Panel (OAP). The view of the OAP was that, given the substantial uncertainty in the model projections for the Eastern oyster and the limited information available on many of the factors that could affect Suminoe oyster outcomes in the Bay, application of the model to the proposed action would yield outcomes with extremely high levels of uncertainty. After consideration of the OAP comments, the demographic modeling team concurred with the OAP that application of the model to the Suminoe oyster would not be appropriate. Thus, alternatives involving Suminoe oysters will not be modeled. However, the total number of triploid Suminoe oysters likely to be reared in aquaculture operations throughout the Bay has been estimated by economic analysis (D. Lipton, University of Maryland) and these data will be combined with model output for natural production of Eastern oysters to evaluate alternatives 4 and 5.

5.0 Model Performance

The validity of any model is normally assessed by comparing model output to an independent data set representing the modeled outcome. In the case of this Bay-wide oyster demographic model, no long-term, Bay-wide, spatially specific oyster abundance data sets are available that could serve as a basis for model validation. Although the MDNR survey used to develop many of the model parameters represents the most spatially and temporally extensive survey data available, these data are only available for Maryland and absolute or relative densities of oysters cannot be estimated correctly from the survey data, as we have described above. Conversely, Virginia conducts intense sampling of some localized areas, but with less spatial coverage than the Maryland survey.

Because validation of the demographic model was not possible, we reviewed all existing data identified in the course of model development to assess whether any data sets were available that would allow for some level of checking of model function. Data from an oyster survey in the James River, conducted by the Virginia Institute of Marine Science (data provided courtesy of R. Mann, Virginia Institute of Marine Science), could serve that purpose. This data set was long term, provided rigorous annual population estimates (i.e., the survey was designed to estimate population size), and provided annual population composition information. Density was estimated in this survey using patent tongs at randomly selected 1-m² areas of bottom. All oysters captured were counted and measured to the nearest 5-mm class. Total numbers of oysters were estimated by multiplying mean densities at a bar by the bar's area.

Although these data provide a suitable time series for checking model functionality, they are not suitable for validating the model. The data represent oyster population behavior in a single small tributary of the Bay, but the model employs population vital parameters that are in essence Bay-wide averages of those rates. For example, James River survival and recruitment rates are likely to be substantially different than the Bay-wide average. Thus, we used the James River data to test model performance broadly by comparing predicted values from a modified version of the model with survey estimates. Modeled trends that diverged from the James River sampled abundance by a large amount would suggest that the model parameterization was incompatible with a known starting population or that there was an error in model specification.

In order to conduct this model check using James River data, the demographic model had to be modified in several ways. The model was run for nine years rather than ten, 1994-2003. Weather years, salinities, disease tiers, and MSX outbreaks were input from historical data rather than modeled. Harvest levels were likewise based on historic data on oyster landings (courtesy J. Wesson, Virginia Marine Resource Commission). This was done by removing a constant fraction of the population from each bar in the simulation to obtain a total harvest equal to the reported landings for the year. The model was also modified to accept fall survey estimates of spat abundance < 30 mm instead of estimating recruitment each year (most spat were <30 mm in the James River because of their slow growth; Section 2.8). These runs were used primarily to evaluate the growth and mortality portions of the model. When recruitment was estimated, spat were distributed in proportion to the area of each bar within salinity zones so that no larvae were imported or exported to other basins. We conducted 1,000 runs of the demographic model for the James River in which historical spat abundance was input, and another 1,000 runs in which recruitment was modeled normally.

The model produced estimates similar to the survey for abundance of small oysters, but projected that more market-size oysters would be present for runs in which spat < 30 mm were input (Figure 10A). The greater number of market sized oysters projected by the model suggests that James River mortality rates were actually greater than those used in the model. A possible explanation for this result is that oysters in the James River are exposed to disease and predation for a longer period at a given size because of their relatively slow growth rate (Section 2.8), resulting in a greater natural mortality than was modeled for Bay conditions on average. Trends in abundance were similar between model projections and survey estimates. We judged model performance for mortality and growth parameterization to be reasonable based on these results.

Differences between model projections and survey estimates were greater when recruitment was also modeled (Figure 10B). Abundance of market-size oysters was projected to be greater than survey estimates, as above. Conversely, abundance of spat was projected to be less than the survey estimates. This resulted in fewer small oysters present in model projections than in survey estimates during years 2-6. In the actual data set, an exceptionally large spat set occurred during 2001, which was not matched by the model projection. Such rare large spat sets occur randomly in the demographic model, as described in Section 2.6. They are partly reflected by

the wide confidence intervals in years 5-8. The predicted model trend, however, does not exhibit such large episodic events because it is the median of many runs. This is an important difference between model projections and the actual survey data. The survey represents a single realized time series; it is unknown if the series was a statistically rare event. The large spat set was quickly reduced as a result of high natural mortality. This resulted in similar estimates of abundance between the model and survey the next year. The final modeled numbers of oysters were similar to the survey for all size classes in year 9, and most of the field values were within a factor of 1 to 4 of the predicted values. These results suggest that the demographic model projects reasonable mean abundance estimates given the large uncertainty in parameter estimates, and that projections are suitable for comparing alternative restoration strategies for the EIS.

6.0 Alternative 1-3 Results

Abundance of market-size Eastern oysters was predicted to remain near present low levels in the simulation for EIS alternative 1, no action (Figures 11A, 12) and declined slightly for small oysters and spat (recall that the no action alternative includes restoration activity at current levels). High natural mortality rates strongly controlled the population, resulting in variation in simulated abundances of only about 25% for harvest rates ranging from 20 to 80% on harvestable bars (71% of all bars simulated). A detailed explanation for the limited effect of harvest when natural mortalities are high is presented in Appendix F. Abundance was projected to increase in range of 250-450% by year 10 under EIS alternatives 2A and 2B, enhanced restoration. However, the large percentage increase is from a very small starting population. Variation in these enhanced restoration simulations was much greater than in the no-action alternative. Some differences between the 10th and 90th percentiles ranged more than 500%, resulting in some overlap with alternatives 1 and 3. Fishing had a greater effect on abundance for alternatives 2A and 2B than it did for alternative 1, indicating that harvest had a larger proportional effect as abundance increased. Winter harvest rates of 20-80% resulted in differences in median predicted abundance greater than 100% for alternatives 2A and 2B. Small oysters and spat were predicted to increase less than market-size oysters for alternatives 2A and 2B, in the range of 100% and 200% respectively. The occurrence of episodic large spat sets resulted in these distributions tending to be skewed positively, particularly for spat. Projections for EIS alternatives 2A and 2A were similar, indicating that the reduction in seed planting of 6 bars in low salinity waters was largely offset by additional seed planting at 13 bars in medium- or high-salinity waters. Predictions for alternative 3, no harvest, resulted in a 56% increase in median abundance from the starting population, indicating that high natural mortality rates strongly controlled even unharvested populations. The increase was much less than those predicted for alternatives 2A and 2B.

Abundance of all size classes was predicted to increase at least somewhat in Maryland under all alternatives (Figures 11B, 13), and to decrease in Virginia (Figures 11B and 14), because a greater proportion of bars in Maryland were located in low salinity waters where natural

mortality rates were relatively low. Although the proportional increase in spat and small oysters was predicted to be much greater in Maryland for alternatives 2A and 2B (Figure 11B), the differences in actual abundance were much less (Figure 11C). This is because many more spat and small oysters in the starting population were located in Virginia (Table 1). Distributions tended to be skewed positively in Virginia because episodic high-recruitment events occurred more frequently in higher salinity waters. Conversely, distributions of small and market-size oysters tended to be skewed slightly negatively for Maryland because lower salinity conditions resulted in some high-mortality events in dry years, and some freshet mortality occurred during wet years (but few large spat sets occurred).

Bars located in low salinity were predicted to have the greatest increases in abundance of market-size oysters under each scenario, but the poorest recruitment of spat (Figure 11D, 15-17). Bars located in high salinity exhibited much greater and more variable recruitment of spat and small oysters, but they were quickly overwhelmed by high natural mortality rates as they reached market sizes. Confidence intervals were also asymmetric for bars in low salinity (Figures 15-17) because salinity changed with annual precipitation in the model (Section 2.5). During dry years, abundance was projected to be very low because few bars were located in low salinity, and those bars exhibited characteristically low recruitment (Section 2.6). During wet years, more bars were located in low salinity, but recruitment was still low in these areas.

Trends in abundance were related to specific management scenarios. For alternative 2A, abundance of spat increased with seed planting in Maryland until year 7, when planting reached its maximum, and remained relatively constant thereafter (Figure 13). Abundance of small oysters also remained constant during the final years of the simulation for Maryland, suggesting that the population would reach a new equilibrium near the abundance levels projected for year ten if this level of management activity were maintained. Spat abundance was projected to decline during years 9 and 10 in Virginia because habitat improvements were reduced in years 8 and 10, and the effect of previous habitat improvements was reduced with time (Section 2.12). Temporal trends were similar between alternatives 2A and 2B overall, by state (Figures 13-14) and by salinity zone (Figures 15-17).

The goal specified in the EIS is to establish an oyster population that reaches a level of abundance in Chesapeake Bay that would support sustainable harvests comparable to harvest levels recorded during the period 1920 to 1970. Assuming the best point estimate of current harvest (47%) is sustainable, the population needed to support the mean harvest level during 1920 to 1970 would be 1.16×10^{10} market-size oysters². Alternatives 2A and 2B were projected to reach, even at the 95th percentile, only about 1/6 of this level. None of simulations conducted projected a population increase that could reach the goal of the EIS in ten years with 90% confidence).

² Bay-wide annual harvests (expressed in Maryland bushels) for the years 1920 to 1970 were averaged. Bushels were converted to number of market-sized oysters using the equation 1 oyster = 0.0009 bushels. The resulting number of oysters was divided by 0.47 to derive the numerical goal.

We note that projected abundances of spat, small, and market-size oysters were similar and relatively constant (Figure 13) despite the fact that many young oysters must be present to produce older oysters when mortality rates are high. This pattern is a result of the small and market classes representing more than one cohort. An average oyster remains in the small size class between 2 (mean size 70 mm) and 3 (mean size 88 mm) years in Maryland, and for about 4 years (mean size 71 mm) in Virginia (Figure 6). The market size class includes larger oysters from all cohorts. Furthermore, spat are counted in the fall in the demographic model, after the period in which they experience the greatest mortality. Most have nearly reached the small size class at this time.

7.0 Model Sensitivity

Additional model runs were conducted to quantify the effects of model parameterization on overall predictions of abundance. Parameters generally were varied by conducting four additional runs for each parameter; runs were conducted with increases of 10% and 25%, and decreases of 10% and 25%. Except for the parameter that was varied, each run was conducted using the same parameterization as alternative 1, no action, and a 40% harvest rate. Sensitivity was quantified by comparing the median value in model-year 10 from each run to the median value in year 10 of the run for alternative 1, 40% harvest (e.g., 3.4⁸ market-size oysters; Figure 10). We analyzed only the change in median values for these runs. No attempt was made to adjust a parameters variance. To avoid reporting an inordinate number of simulations with all combinations of alternatives and variables, we limited analysis of higher order interactions in which more than one parameter was changed to the most extreme combinations of parameters (all +25% or -25%) for recruitment, growth, and mortality only. Results of all runs are summarized in Table 5 as the percent change in estimated abundance.

Varying growth parameters affected predictions in a way that would be biologically important. An increase or decrease of 25% in K resulted in a corresponding change in predicted abundance of about 25% for market-size oysters. The model was more sensitive to changes in the H_{∞} parameter than K . Changes of 25% in H_{∞} resulted in greater than 30% changes in market-size oyster abundance. Model projections changed by more than 50% for market-size oysters when both parameters were varied simultaneously. The effect was greater for a 25% increase in K and H_{∞} (87%) than for a decrease (-54%). The sensitivity of the H_{∞} parameter creates uncertainty in the model that cannot be quantified easily given the limitations in available data. The estimate is based on historical data because few oysters now survive long enough to reach H_{∞} (Section 2.8). Varying recruitment parameters had relatively small effects on predicted abundance. Changes of 10% or 25% in the spat to spawner ratios all changed abundance of market-size oysters less than 10%. Reducing the spat to spawner ratio by 10% did not change the predicted population of small oysters in the expected direction (a 1% increase resulted). This is likely because the effect of the parameter increase was small relative to the random variation in medians resulting from 1,000 runs. Eliminating episodic large spat sets had a larger effect than decreasing the spat to spawner ratio even by 25%. This result supports the idea that

infrequent large spat sets are important in supporting the population. Varying egg-production coefficients that were used in estimating recruitment also had small and inconsistent effects on model predictions.

The mortality rates entered into the model had important effects on predicted abundance. A 25% decrease in mortality rates resulted in a 94% increase in abundance of market-size oysters, and an increase of more than 100% in spat and small oysters. This result demonstrates not only the importance of obtaining reliable data with which to estimate mortality, but also that oyster populations would likely increase if natural mortality rates could be reduced.

The model was most sensitive to increases in multiple parameters that interacted. Increases in the ratio of spat to spawners interacted non-linearly with growth and mortality, resulting in large increases in population size when 25% adjustments were made. The largest population increases occurred when growth and the spat to spawner ratio were both increased by 25%. This model caused more than a ten-fold increase in the abundance of market-size oysters. When a reduction in mortality rates of 25% was added to the interaction, the increase was nearly thirty-fold.

Use of the larval transport model did not have a large effect on model results; simulations in which spat settled within the basin that they were produced rather than according to the larval transport model produced results within 5% of the corresponding simulations that used the transport model. This was expected because most larvae remain within the same basin even when the larval transport model is used (>90% for most basins; North et al 2006). The larval transport model would be important for predicting dispersal of oysters among basins if we had modeled an establishing population of Suminoe oysters. The model was likewise relatively insensitive to freshets.

Modeling random weather years instead of selecting blocks of weather years from the historical data, and simulating illegal harvest on protected bars, had moderate effects on projections. These results suggest that the effects of climate change and illegal harvest may have to be considered in designing an oyster recovery program.

The model was relatively insensitive to 10 or 25% changes in the starting population (predicted abundance changed <10%), but more sensitive to very large increases in the starting population. We conducted additional sensitivity analyses where the starting population was multiplied or divided by 100 because of the great uncertainty associated with this input parameter. The model was relatively insensitive to reducing the starting population by dividing it by 100. However, increasing the starting population by 100× resulted in 1,767% increase in market-size oysters under alternative 1. Alternatives were affected differently by such a large increase. Restoration had a much smaller relative effect when the population was 100× larger; market-size oysters were predicted to increase by 471% for alternative 2A. Fishing had a much

greater relative effect; market-size oysters were predicted to increase 3,853% for alternative 3, no harvest.

8.0 Data and Uncertainty

The demographic model integrates data about population dynamics and environmental conditions to produce abundance predictions that are suitable for comparing alternative management strategies in the EIS. The model produces results that appear credible given the data currently available for oysters in the Bay. We emphasize that the model should be viewed as a tool to provide an integrated view of a complicated multivariate problem, not a predictor of absolute abundance of oysters in the Bay. This is because the model produces expected trends in abundance, but a single realized event that involves random probabilities actually occurs. The starting population provides a demonstration. For example, the number of spat in the starting population is fewer than the number of small or market-size oysters in Maryland (Table 1). This is unsustainable for a long period and, thus, cannot represent an ‘average’ condition. If the result is not due to sampling error, then it represents recruitment failure, an extreme event. The small number of spat causes a mismatch between the starting population and model parameterization that leads to rapid changes in the projected population for the first several years of the simulation (Figures 13.1-13.4, and 13.13). The number of market-size oysters then becomes low relative to the number of spat and small oysters given the modeled mortality rates for market-size oysters. This causes the population of market-size oysters to increase for several years after the number of spat and small oysters reaches a relatively constant level. Similarly, the starting population is based on a wet year, 2004. The starting population is much greater than the modeled predictions for year 1 in low salinity (Figure 15) because many bars are included in low salinity that, on average, would be classified as having a greater salinity. The opposite occurs for high salinity (Figure 17). The demographic model is designed to provide a weight-of-evidence approach to evaluating alternatives, not to predict a specific outcome with certainty.

Confidence intervals estimated from the results of 1,000 model runs were relatively wide for all restoration scenarios. These results reflect the great uncertainty associated with model parameterization, and stochastic events that determine oyster population-vital rates. Nevertheless, these results provide an integrated view of what is known about oyster population dynamics in the Chesapeake Bay. They are intended to serve as probabilistic descriptions of different scenarios rather than hypothesis tests. For example, model results indicate the 95th percentile for alternative 1 with a 40% harvest rate is a 97% increase in market-size oysters. This is the greatest increase that is likely to be attainable (Figures 11A and 12). The lower 5th percentile for alternative 2A with a 40% harvest rate indicates a 92% increase. Although there is some overlap between these alternatives, nearly all of the runs for alternative 2A indicated an increase of 100% or more. Therefore, alternative 2A is much more likely to result in an increase in oyster abundance of 100% or more than alternative 1.

In practice, the uncertainty associated with implementing a restoration strategy is probably even greater than the confidence limits indicate because many real-world processes that affect oyster populations are not modeled. For example, harvest rates may interact with disease resistance, and available habitat may change during the ten-year period, as described below. The variation reported here only reflects the propagation of uncertainty in modeled parameters. The quality and spatial coverage of data sets used to develop model parameters also increases the uncertainty associated with the model. Data used in the modeling effort generally were collected for other research projects and later contributed for model development. The MDNR data used to develop the starting population and several other parameters were collected using a survey where area swept was not recorded because the purpose was simply to collect oysters for disease sampling. The effect of an error in the starting population is not expected to be large if the error is within 25%, but would be important if the starting population is badly underestimated (i.e., 100×; Table 5). Another important limitation is that growth data for the James River were applied to oysters throughout most of Virginia. Although these rates are believed to be generally representative of growth rates in Virginia as a whole (Section 2.8), the model would be biased to an unknown extent if large differences exist. Model results are likely to be valid only within the spatial and temporal scales at which we present the data. Despite the fact that oyster dynamics were modeled on individual bars, they are aggregated to salinity zones, states, or the entire Bay. This is because many of the input parameters of the model were averages over many bars. Therefore, the demographic model is expected to perform well over larger spatial scales but not specifically for individual bars. It would also be inappropriate to run the model for more than the 10-year period programmed in the current implementation. A major reason for this is that the model uses a fixed habitat layer. That is, the model does not include any changes to habitat availability, either as a change in the size of individual bars or creation of new bars. The amount of habitat could change as sedimentation reduces the area of suitable bars and shells deteriorate (Smith et al. 2005), or as oyster populations increase and create additional habitat. The model also does not simulate any long-term changes in environmental conditions such as water quality and disease intensity that would affect oyster population dynamics. An unorthodox approach was required to model recruitment because of the unique nature of the data available. The data differed from those typically used to develop a stock-recruitment relationship in several ways: 1) only the ratios of spat and spawners were available, not absolute or relative density estimates; 2) unpredictable large episodic spat sets were known to be important in sustaining the population; 3) no weight measurements were available; and 4) no observations were available at high stock levels where density-dependent interactions were likely to occur. The approach taken here produced reasonable results because the magnitude of normal and episodic large spat set increased with standardized stock size (Figure 5). The recruitment submodel will almost certainly overestimate recruitment at some very large stock sizes because density-dependent reductions in recruitment would occur. We had no data with which to estimate the magnitude of density dependence or the stock size at which it would begin to occur; however, we assume that the stock size would have to be many orders of magnitude greater before density-dependent

effects became important because the Bay has historically supported much larger populations. Another limitation of the recruitment submodel is that it is difficult to determine which spawners contributed to recruitment on a particular bar because oyster larvae may be transported by wind and current from throughout the Bay. We assumed that calculating an average Bay-wide recruitment by salinity zone and redistributing some spat according to the larval-transport model (Section 2.7) would provide the best overall estimates, although localized predictions based on this model would not be appropriate. Sensitivity analyses indicated that some inaccuracy in the estimation of recruitment alone did not have large effects on model outcome, but the interaction between recruitment and growth or mortality did have large effects. Data from surveys currently under development by the MDNR should be useful for future validation and refinement of this important part of the model.

9.0 Conclusions

Under current conditions, the model indicated that high natural mortality rates strongly controlled the population. Harvest did not have a large effect for alternative 1, but became increasingly important as populations increased in alternatives 2A and 2B. This is because most large oysters died quickly whether they were harvested or not, and the cohort just reaching market size tended to dominate. We provide a mathematical explanation for this unintuitive result in Appendix F. The large fraction of bars that were not harvested in the model (sanctuaries and closed areas) also acted to stabilize populations because oysters from these bars were available to spawn under any harvest rate. If an appreciable amount of illegal harvest occurs, the model projections may not accurately reflect the importance of harvest (Table 5). We do not suggest that managing harvest is unimportant for oysters in the Bay under current conditions, only that it did not have a large effect on absolute numbers in alternative 1 because populations are very low and natural mortality rates very high. If a portion of oysters were modeled to develop disease resistance, for example, the rate at which they were harvested would have important effects.

The model indicated that alternatives 2A and 2B were most likely to increase oyster abundance. However, none of the alternatives modeled were likely to reach the EIS goal within ten years. This is partly because the ten-year period established for recovery is too short for a long-lived species such as the Eastern oyster to increase its population to that degree. Even if Dermo disease level was low (tier 3), no MSX events occurred, and no harvest was conducted, an exploratory simulation projected that the population would likely reach about half of the goal in ten years (Table 5).

The data sets used to develop the demographic model are not comprehensive, nor were they collected in an ideal manner for model development. Some limitations of the data create uncertainty in model results that cannot be quantified. Nevertheless, the demographic model provides a structured evaluation of the relative effect of different management scenarios using the best data currently available. It provides the only realistic way to integrate the large number

of parameters that control oyster abundance in the Bay to visualize the likely effects of different management alternatives for Eastern oysters.

10.0 Literature Cited

- Barker, L. S. 2007. Spatially distributed estimate of the 2004 Maryland Chesapeake Bay population of *Crassostrea virginica*. Maryland Department of Natural Resources Fisheries Service. Internal report dated 14 March 2007. 12pp.
- Beaven, G. F., 1950. Growth observations of oysters held on trays at Solomons Island, Maryland. Proceedings of the National Shellfish Association 1949:43-49.
- Beaven, G. F., 1953. Some observations on rate of growth of oysters in the Maryland area. Proceedings of the National Shellfish Association 43:90-98.
- Butler, P. A., 1953. Shell growth versus meat yield in the oyster *Crassostrea virginica*. Proceedings of the National Shellfish Association 43:157-162.
- Cahn, A. 1950. Oyster culture in Japan. United States Fish and Wildlife Service Fishery Leaflet No. 383. 80 pp.
- Calvo, G. W., M. W. Luckenbach, S. K. Allen Jr, and E. M. Burreson. 1999. Comparative field study of *Crassostrea gigas* (Thunberg, 1773) and *Crassostrea virginica* (Gmelin, 1791) in relation to salinity in Virginia. Journal of Shellfish Research 18 : 465-473.
- Calvo, G. W., M. W. Luckenbach, S. K. Allen Jr., and E. M Burreson. 2001. A comparative field study of *Crassostrea ariakensis* (Fujita 1913) and *Crassostrea virginica* (Gmelin 1791) in relation to salinity in Virginia. Journal of Shellfish Research 20:221-229.
- Chai, A., M. Homer, C. Tsai, and P. Gouletquer. 1992. Evaluation of oyster sampling efficiency of patent tongs and an oyster dredge. North American Journal of Fisheries Management 12:825-832.
- Coakley, G. M. 2004. Growth of Eastern Oyster, *Crassostrea virginica*, in Chesapeake Bay. M.S. University of Maryland, College Park.
- Edmonds, R. H. 1887. Review of the oyster industry – the oyster interests of Maryland. Page 430 in G. B. Goode, editor. The fisheries and fishery industry of the United States. Section III. A geographical review of the fisheries industries and fishing communities for the year 1880. U.S. Bureau of Fisheries. Washington D.C., U.S.A.
- Efron, B., and R. J. Tibshirani. 1993. An introduction to the bootstrap. Chapman and Hall/CRC Press. New York, New York, U.S.A.

- Emerson, J. D., and J. Strenio. 1983. Boxplots and batch comparison. Pages 58-96 in Hoaglin, D. C., F. Mosteller, and J. W. Tukey, eds. Understanding robust and exploratory data analysis. Wiley and Sons, New York, New York, U.S.A.
- Federal Register. 2004. Intent [to prepare a programmatic environmental impact statement] for a proposed introduction of the oyster species *Crassostrea ariakensis* into the tidal waters of Maryland and Virginia to establish a naturalized, reproducing, and self-sustaining population of this oyster species. Volume 69:2, January 5, 2004.
- Ford, S. E., M. J. Cummings, and E. N. Powell. 2006. Estimating mortality in natural assemblages of oysters. *Estuaries and Coasts* 29:361-374.
- Ford, S. E. & M. R. Tripp. 1996. Disease and Defense Mechanisms. In: V. S. Kennedy, R. I. E. Newell & A. F. Eble, editors. The Eastern Oyster *Crassostrea virginica*. Maryland Sea Grant College, University of Maryland, College Park, MD. pp. 581-642.
- Gieseke, C. M., 2001. Year 2000 Maryland oyster disease status report. Maryland Department of Natural Resources Fisheries Service. Sarbanes Cooperative Oxford Laboratory Report FS-SCOL-01-1. Oxford, Maryland, U.S.A.
- Grabowski, J. H. 2004. Habitat complexity disrupts predator-prey interactions but not the trophic cascade on oyster reefs. *Ecology* 85:995-1004.
- Greenhawk, K. 2005. Development of a potential habitat layer for Chesapeake Bay Oyster Bottom. Maryland Department of Natural Resources. Annapolis, Maryland, U.S.A.
- Greenhawk, K. and O'Connell, T. 2007. Oyster population estimates for the Maryland Portion of Chesapeake Bay 1994 – 2006. Maryland Department of Natural Resources. Annapolis, Maryland, U.S.A.
- Jordan, S. J., K. N. Greenhawk, C. B. McCollough, J. Vanisko, and M. L. Homer. 2002. Oyster biomass, abundance and harvest in northern Chesapeake Bay: trends and forecasts. *Journal of Shellfish Research* 21:733-741.
- Kennedy, V.S. 1996. Biology of Larvae and Spat. In: V. S. Kennedy, R. I. E. Newell & A. F. Eble, editors. The Eastern Oyster *Crassostrea virginica*. Maryland Sea Grant College, University of Maryland, College Park, MD, U.S.A. pp. 371-421.
- Kennedy, V. S., R. I. E. Newell & A. F. Eble, editors. 1983. The Eastern Oyster *Crassostrea virginica*. Maryland Sea Grant College, University of Maryland, College Park, MD, U.S.A.
- Kennedy, V. S. 1983. Sex ratios in oysters, emphasizing *Crassostrea virginica* from Chesapeake Bay, Maryland. *The Veliger* 25:329-338.

- Mann, R., and D. A. Evans. 1998. Estimation of oyster *Crassostrea virginica*, standing stock, larval production and advective loss in relation to observed recruitment in the James River, Virginia.
- McHugh, J. L., and J. D. Andrews. 1955. Computation of oyster yields in Virginia. *Proceedings of the National Shellfish Association* 45:217-239.
- Merritt, D., A. Padeletti, S. K. Allen Jr., and H. Harwell. 2006. Evaluation of gametogenesis and spawning cues in diploid *Crassostrea ariakensis* for the estimations of environmental risk and establishment of diploid brood stock populations. Final report for NOAA award NA04 NMF457 0428, University of Maryland Center for Environmental Science.
- Newell, R. I. E., V. S. Kennedy, J. L. Manuel, and D. Meritt. 2005. Behavioral responses of *Crassostrea ariakensis* and *Crassostrea virginica* larvae to environmental change under spatially realistic conditions. Final report to Maryland Department of Natural Resources, Annapolis, MD, 28 pp.
- North, E. W., Z. Schlag, R. R. Hood, L. Zhong, T. and M. Li. 2006. Modeling dispersal of *Crassostrea ariakensis* oyster larvae in Chesapeake Bay. Final report to Maryland Department of Natural Resources. Annapolis, MD, U.S.A. 55 pp.
- North, E. W., Z. Schlag, R. R. Hood, M. Li, L. Zhong, T. Gross, and V. S. Kennedy. In Press. Vertical swimming behavior influences the dispersal of simulated oyster larvae in a coupled particle-tracking and hydrodynamic model of Chesapeake Bay. *Marine Ecology Progress Series*.
- Pinheiro, J., D. Bates, S. DebRoy, and D. Sarkar. 2006. nlme: Linear and nonlinear mixed effects models. R package version 3.1-77.
- Powell, E. N. 2005. Fisherman choice and incidental catch: size frequency of oyster landings in the New Jersey oyster fishery. *Journal of Shellfish Research* 24:469-477.
- R Development Core Team. 2006. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>.
- Quinn, T. J., and R. B. Deriso. 1999. Quantitative fish dynamics. Oxford University Press. New York, New York, U.S.A.
- Rothschild, B. J., J. S. Ault, P. Gouletquer, M. Heral. 1994. Decline of the Chesapeake Bay oyster population: a century of habitat destruction and overfishing. *Marine Ecology Progress Series* 111:29-39.
- Shaw, W. N. 1966a. Advances in the off-bottom culture of oysters. *Proceedings of the Gulf Caribbean Fisheries Institute* 19:108-115.

- Shaw, W. N. 1966b. The growth and mortality of seed oysters, *Crassostrea virginica* from Broad Creek, Chesapeake Bay, Maryland, in high- and low-salinity waters. Proceedings of the National Shellfish Association 56:59-63.
- Shaw, W. N., and A. S. Merrill. 1966. Setting and growth of the American oyster *Crassostrea virginica* on navigation buoys in the lower Chesapeake Bay. Proceedings of the National Shellfish Association 56:67-72.
- Shumway, S. E. 1996. Natural environmental factors. Pages 467-513 in V. S. Kennedy, R. I. E. Newell & A. F. Eble, eds. The eastern oyster *Crassostrea virginica*. Maryland Sea Grant College Publication, College Park, Maryland, U.S.A.
- Smith, G. F., D. G. Bruce, E. B. Roach, A. Hansen, R. T. E. Newell, A. M. McManus. 2005. Assessment of recent habitat conditions of eastern oyster *Crassostrea virginica* bars in mesohaline Chesapeake Bay. North American Journal of Fisheries Management 25:1569-1590.
- Southworth, M. J., A. Harding, and R. Mann. 2005. The status of Virginia's public oyster resource. 2004. Molluscan Ecology Program, Virginia Institute of Marine Science, Gloucester Point, Virginia, U.S.A.
- Tarnowski, M., Maryland Department of Natural Resources Shellfish Program, and Sarbanes Cooperative Oxford Laboratory. 2003. Maryland oyster population status report, 2002 fall survey. Maryland Department of Natural Resources Report. Annapolis, Maryland, U.S.A.
- Ulanowicz, R.E., W.C. Caplins, and E.A. Dunnington. 1980. The forecasting of oyster harvest in Central Chesapeake Bay. Estuarine and Coastal Marine Science 11:101-106.
- U.S. Global Change Research Program. 2000. Climate change impacts on the United States, the potential consequences of climate variability and change. Available at <http://www.usgcrp.gov/usgcrp/Library/nationalassessment>.
- Vølstad, J. H., J. Dew, and M. Tarnowski. In Press. Estimation of annual mortality rates for eastern oysters (*Crassostrea virginica*) in Chesapeake Bay based on box counts, and application of those rates for projection of population growth of *C. virginica* and *C. ariakensis*. Journal of Shellfish Management.
- vonBertalanffy, L. 1938. A quantitative theory of organic growth (Inquiries on growth laws. II). Human Biology 10:181-213.

11.0 Tables

Table 1. Number of oysters in the starting population at model initiation by state, salinity zone, and aggregated to the entire Bay. Estimates are based on abundance for 2004, a wet year. Note that the area of salinity zones (low < 11 ppt, medium 11-15 ppt, or high \geq 15 ppt) fluctuates with annual precipitation. Thus, oysters on individual bars may change salinity zones during different years in the model.

| Oyster size | State | | Salinity zone | | | Whole Bay |
|-------------|-------------|---------------|---------------|-------------|------------|---------------|
| | Maryland | Virginia | Low | Medium | High | |
| Market | 126,754,209 | 173,832,742 | 155,289,829 | 80,528,252 | 64,768,870 | 300,586,951 |
| Small | 101,298,360 | 984,801,546 | 603,194,238 | 387,756,300 | 95,149,368 | 1,086,099,906 |
| Spat | 23,558,031 | 1,099,596,029 | 649,253,828 | 413,617,154 | 60,283,078 | 1,123,154,060 |

Table 2. Estimated proportions of female oysters in the population (sex ratio) by size class. Data are based on field experiments reported by Kennedy (1983). Proportions are denoted as R in Equation 3, where $r = (0-40), (41-80), \dots, (>160)$.

| Size class (height in mm; r) | Proportion of females (R) |
|---------------------------------|-------------------------------|
| 0-40 | 0.50 |
| 41-80 | 0.44 |
| 81-120 | 0.63 |
| 121-160 | 0.67 |
| >160 | 0.83 |

Table 3. Probability of selecting disease tier 1, 2, or 3 based on the current weather year for individual bars in the demographic model. Probabilities are based on mean disease intensities measured at disease bars in the MD DNR fall survey for the years 1990-2005.

| Weather Year | Disease Tier | | |
|--------------|--------------|------|------|
| | 1 | 2 | 3 |
| Dry | 0.80 | 0.20 | 0.00 |
| Average | 0.00 | 0.75 | 0.25 |
| Wet | 0.00 | 0.17 | 0.83 |

Table 4. Average annual natural mortality rates based on recent box counts for eastern oysters by salinity class and disease tier. Average time-since-death of recent boxes was assumed to be between 7 and 14 days. The relative standard error, $RSE = SE/Mean$; LCL and UCL are the upper and lower 95% confidence limits, respectively.

| Salinity | Tier | Market-Sized Oysters | | | | Small Oysters | | | |
|----------|------|----------------------|------|------|------|---------------|------|------|------|
| | | Mean | RSE | LCL | UCL | Mean | RSE | LCL | UCL |
| High | 1 | 0.79 | 0.14 | 0.57 | 1.00 | 0.69 | 0.18 | 0.45 | 0.93 |
| High | 2 | 0.51 | 0.24 | 0.27 | 0.75 | 0.47 | 0.25 | 0.25 | 0.70 |
| High | 3 | 0.23 | 0.31 | 0.09 | 0.37 | 0.34 | 0.28 | 0.16 | 0.53 |
| Med | 1 | 0.59 | 0.21 | 0.34 | 0.83 | 0.56 | 0.22 | 0.31 | 0.80 |
| Med | 2 | 0.43 | 0.26 | 0.21 | 0.64 | 0.39 | 0.27 | 0.18 | 0.59 |
| Med | 3 | 0.13 | 0.32 | 0.05 | 0.21 | 0.16 | 0.31 | 0.06 | 0.26 |
| Low | 1 | 0.34 | 0.29 | 0.15 | 0.54 | 0.29 | 0.30 | 0.12 | 0.46 |
| Low | 2 | 0.22 | 0.31 | 0.08 | 0.35 | 0.15 | 0.33 | 0.05 | 0.25 |
| Low | 3 | 0.10 | 0.33 | 0.03 | 0.16 | 0.08 | 0.33 | 0.03 | 0.13 |

Table 5. Sensitivity of demographic model to variation in underlying parameters. All sensitivity runs were compared to the run for alternative 1, no action, with a 40% harvest rate (Figure 11) unless otherwise specified. Results are reported as the percent change in median oyster abundance at year ten of the model.

| Model tested | Percent difference between model tested and no-action model in median oyster abundance at year ten | | |
|---|--|-------|--------|
| | Spat | Small | Market |
| Recruitment | | | |
| Spat/spawner ratio +10% | 9 | 6 | -1 |
| Spat/spawner ratio -10% | -2 | 1 | -4 |
| Spat/spawner ratio +25% | 15 | 10 | 1 |
| Spat/spawner ratio -25% | -11 | -6 | -6 |
| No episodic large spat sets | -12 | -11 | -6 |
| Egg production ($F_{ind} = 39.06 \times [0.000423 \times H^{1.7475}]^{2.36}$; Eqn. 1) | | | |
| Egg production inner coefficient +25% | 1 | 2 | -1 |
| Egg production inner coefficient -25% | -2 | -1 | -1 |
| Egg production inner exponent +25% | 5 | 5 | 3 |
| Egg production inner exponent -25% | -1 | -2 | -3 |
| Egg production outer exponent +25% | 2 | 2 | 2 |
| Egg production outer exponent -25% | 2 | 0 | 0 |
| Growth (H_{∞} and K from the vonBertalanffy growth equation) | | | |
| H_{∞} +10% | 9 | 2 | 14 |
| H_{∞} -10% | -4 | -1 | -14 |
| H_{∞} +25% | 35 | 11 | 44 |
| H_{∞} -25% | -6 | -8 | -37 |
| K +10% | 6 | 3 | 13 |
| K -10% | -3 | -2 | -9 |
| K +25% | 21 | 7 | 29 |
| K -25% | -4 | -9 | -25 |
| H inf and K +25% | 71 | 35 | 87 |
| H inf and K -25% | -1 | -12 | -54 |
| Mortality | | | |
| Mortality -10% | 29 | 28 | 28 |
| Mortality +10% | -16 | -17 | -21 |
| Mortality -25% | 113 | 108 | 94 |
| Mortality +25% | -30 | -37 | -40 |
| Interactions | | | |
| H_{∞} and K +25%, mortality -25% | 544 | 384 | 400 |
| H_{∞} and K -25%, mortality +25% | -18 | -40 | -72 |
| H_{∞} , K, and spat/spawner ratio +25% | 1,965 | 1,264 | 1,457 |
| H_{∞} , K, and spat/spawner ratio -25% | -65 | -46 | -45 |
| Mortality -25%, spat/spawner ratio +25% | 890 | 902 | 523 |
| Mortality -25%, spat/spawner ratio -25% | -24 | -5 | -32 |

| | | | |
|--|-------|-------|-------|
| H ∞ , K, and spat/spawner ratio +25%, mortality - 25% | 3,792 | 2,651 | 2,980 |
| H ∞ , K, and spat/spawner ratio -25%, mortality + 25% | -24 | -5 | -32 |
| Other parameters | | | |
| Randomly-selected weather years | -4 | -7 | -11 |
| No larval transport model | 2 | 3 | 5 |
| No freshets | 0 | 2 | 6 |
| Starting population +10% | 1 | 2 | 1 |
| Starting population -10% | -1 | -2 | -4 |
| Starting population +25% | 7 | 6 | 5 |
| Starting population -25% | -4 | -5 | -7 |
| Starting population \times 100 | 1,465 | 1,433 | 1,767 |
| Starting population / 100 | -17 | -17 | -21 |
| Alternative 2A, starting population \times 100 | 445 | 391 | 471 |
| Alternative 2A, Starting population / 100 | -5 | -5 | -6 |
| Alternative 3, Starting population \times 100 | 3,478 | 3,146 | 3,853 |
| Alternative 3, Starting population / 100 | -31 | -30 | -39 |
| 10% Illegal harvest in closed areas | -2 | 0 | -13 |
| 25% Illegal harvest in closed areas | -6 | -3 | -16 |
| Tier 3 mortality (low disease), no MSX, no harvest | 2,983 | 2,629 | 1,641 |

12.0 Figures




Figure 1. Estimated density and distribution of oysters in the starting population (spat, small, and market sized) for 8,480 bars included in the demographic model, based on year 2004.




Figure 2. Mean annual discharge into the Chesapeake Bay between 1937 and 2005 (Source U.S. Geological Survey, Chesapeake Bay Activities, <http://md.water.usgs.gov/monthly/bay.html>).





Figure 3. Salinity classes (low < 11 ppt, medium 11-15 ppt, or high \geq 15 ppt) of oyster bars in the demographic model during 1995-1999.

A. Density estimates from Diver and Dredge Surveys, Same Scale



B. Density estimates from Diver and Dredge Surveys, Dredge data rescaled on Secondary Y Axis




Figure 4. Comparison of sampling efficiency of dredging versus diver harvest reported for replicated studies at Hog Island and Walter White reefs in the Chesapeake Bay by Chai et al. (1992). Efficiency of the dredge was low for both reefs (A). Numbers of spat captured relative to small and market-size oysters were similar, as indicated by the same data plotted on different scales (B). Estimates from the dredge survey are scaled by 40.6 times on the secondary axis in B. Vertical bars indicate approximate 95% confidence intervals.




Figure 5. Relationship between number of standardized female spawners (to 77 mm) and spat based on the DNR annual dredge survey, 1990-2006. Gray line indicates a ratio of 3.0 spat per spawner. Observations with higher ratios were considered outliers and used to model episodic large spat sets. Spat were modeled according to salinity as indicated by the dark lines in normal recruitment years




Figure 6. Height-at-age distributions used to model oyster growth in the Chesapeake Bay. A. Data from two recent studies and a literature review that were applied to Maryland and the Potomac River. B. Data from a survey of the James River that were applied to Virginia except the Potomac River. Lines indicate the 5th, 50th, and 95th percentiles of each model. The mean maximum size of oysters H_{∞} was assumed to be 177 mm in both models based on literature (Rothschild et al. 1994) because data were not available for older oysters. The median line for Maryland is $H_{\infty} = 177 * (1 - \exp(-0.1834 * (\text{age} + 0.7232)))$. The median line for Virginia is $H_{\infty} = 177 * (1 - \exp(-0.1078 * (\text{age} + 0.7414)))$.





Figure 7. Mean annual natural mortality rates by disease level (tier 1 = high dermo intensity; tier 2 = medium dermo intensity; and tier 3 = low dermo intensity) and salinity class for small and market-size oysters over all years (1991-2005). Error bars represent 95% confidence intervals. We assumed an average TSD of 1-year for total boxes, and 1-2 -weeks for recent boxes.




Figure 8. The function: $\text{Shell condition} = 1.593 + [0.939 \times \log_e(\text{age})]$ describes a decline in the quality of planted shell for oyster recruitment as planted shell is covered with sediment (Smith 2005; left axis). Condition-class 1 indicates clean shell with no sediment. Condition-class 3 represents shells that have received some sedimentation but are similar to unreplenished bars. Mean spat settlement was observed to be 4.5 times greater on newly replenished bars than on unreplenished bars (MDDNR, unpublished data). Thus, a scaling factor of 4.5 (right axis) was applied to replenished bars in the demographic model for the first year, and the scaling factor was decreased through time following the same rate of decline as the shell-condition model.




Figure 9. Flow diagram of oyster demographic model.






Figure 10. Abundance of oysters in the James River, Virginia, as predicted by 1,000 runs of the oyster demographic model (solid line), and field samples (dashed line). The number of spat < 30 mm was input into the model each year for run A, and estimated by the model for run B. Model estimates are the median value of all runs. Vertical bars indicate 90% confidence intervals, as estimated by the 5th and 95th percentile of all runs.

Figure 11. Change in oyster abundance from the starting population to year 10 projected by 1,000 runs of the demographic model for EIS alternatives 1 (status quo management), 2A, 2B (enhanced restoration), and 3 (no harvest). Harvest rates indicate winter exploitation rates of 20 to 80% on legally harvestable bars (71% of the total), after natural mortality has occurred. Boxes indicate the range of the 25th to 75th percentile of all runs. Vertical bars extend to the 5th and 95th percentile of all runs. Results are presented for the entire Bay (A), Maryland and Virginia (B), and by salinity zone (D). Results for Maryland and Virginia are also presented as change in predicted total abundance rather than percent change from the starting population (C). Note that scales differ among plots for spat, small, and market-size oysters on some panels.


A. Whole Bay




B. By States



C. Estimated **abundance** by State



D. By salinity zone



Figures 12-15. Oyster abundance projected by state in 1,000 runs of the demographic model for EIS alternatives 1 (status quo management), 2A, 2B (enhanced restoration), and 3 (no harvest). Results are reported for harvest rates (winter exploitation rates) of 20 to 80% on legally harvestable bars (71% of the total), after natural mortality has occurred, and categorized by the entire Bay, states, and salinity zones. Tables indicate percent changes in abundance from the starting population at the 5th, 25th, 50th (median), 75th, and 95th percentile for each year of the simulation. Data are depicted as boxplots to the right of each table to illustrate trends. Square symbols indicate the starting population (0% change). Scales differ for each plot. Note that salinity zones do not represent fixed areas. Individual bars may change salinity depending on annual precipitation, but the starting population is for a single wet year based on 2004.

Figure 12 . Whole Bay















Figure 12 . Whole Bay (continued)
















Figure 12 . Whole Bay (continued)


Market 12.9 Alternative 2B, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 43 | 80 | 108 | 108 | 138 | 182 | 256 | 358 | 484 | 604 |
| P75 | 27 | 60 | 84 | 82 | 87 | 120 | 175 | 287 | 374 | 487 |
| P50 | 16 | 25 | 30 | 28 | 53 | 82 | 130 | 208 | 297 | 375 |
| P25 | 2 | -11 | -7 | -7 | 13 | 38 | 79 | 129 | 186 | 252 |
| P5 | -42 | -48 | -48 | -50 | -40 | -29 | -9 | 18 | 51 | 100 |



Small 12.9 Alternative 2B, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -10 | -18 | 2 | 24 | 82 | 87 | 124 | 183 | 195 | 224 |
| P75 | -19 | -31 | -25 | -14 | 16 | 40 | 72 | 106 | 128 | 141 |
| P50 | -28 | -42 | -38 | -29 | -4 | 17 | 48 | 81 | 98 | 104 |
| P25 | -38 | -54 | -62 | -45 | -28 | -5 | 22 | 46 | 50 | 59 |
| P5 | -57 | -70 | -68 | -68 | -58 | -43 | -23 | -7 | 2 | 8 |




Spat 12.9 Alternative 2B, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 117 | 145 | 215 | 262 | 274 | 330 | 415 | 421 | 441 | 445 |
| P75 | 3 | 23 | 81 | 115 | 137 | 178 | 217 | 237 | 227 | 208 |
| P50 | -4 | 9 | 89 | 89 | 105 | 141 | 181 | 195 | 181 | 183 |
| P25 | -15 | -7 | 40 | 65 | 84 | 118 | 158 | 168 | 152 | 133 |
| P5 | -38 | -28 | 19 | 32 | 55 | 85 | 116 | 124 | 105 | 82 |




Market 12.11 Alternative 2B, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 17 | 38 | 47 | 39 | 72 | 111 | 182 | 275 | 387 | 491 |
| P75 | 2 | 12 | 18 | 12 | 39 | 71 | 128 | 212 | 299 | 382 |
| P50 | -8 | -8 | -7 | -9 | 15 | 43 | 83 | 158 | 232 | 291 |
| P25 | -22 | -35 | -35 | -34 | -14 | 11 | 47 | 92 | 142 | 187 |
| P5 | -54 | -63 | -62 | -64 | -55 | -42 | -28 | -3 | 43 | 66 |




Small 12.11 Alternative 2B, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -11 | -20 | -1 | 13 | 47 | 66 | 101 | 146 | 178 | 202 |
| P75 | -21 | -34 | -29 | -20 | 7 | 32 | 84 | 97 | 119 | 131 |
| P50 | -30 | -44 | -41 | -34 | -9 | 13 | 41 | 71 | 88 | 99 |
| P25 | -41 | -56 | -65 | -49 | -31 | -12 | 15 | 33 | 48 | 59 |
| P5 | -59 | -72 | -71 | -68 | -59 | -44 | -28 | -11 | 5 | 6 |




Spat 12.11 Alternative 2B, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 83 | 114 | 184 | 190 | 214 | 279 | 347 | 387 | 401 | 402 |
| P75 | -2 | 16 | 86 | 85 | 115 | 154 | 185 | 219 | 208 | 180 |
| P50 | -9 | 1 | 47 | 74 | 93 | 129 | 188 | 184 | 170 | 150 |
| P25 | -20 | -14 | 31 | 65 | 74 | 111 | 149 | 161 | 145 | 124 |
| P5 | -41 | -33 | 14 | 28 | 45 | 81 | 113 | 121 | 108 | 91 |




Market 12.10 Alternative 2B, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 29 | 54 | 88 | 82 | 94 | 137 | 208 | 319 | 417 | 534 |
| P75 | 14 | 28 | 35 | 30 | 54 | 90 | 147 | 235 | 335 | 428 |
| P50 | 4 | 5 | 6 | 4 | 28 | 59 | 109 | 180 | 264 | 336 |
| P25 | -11 | -27 | -27 | -24 | -4 | 18 | 59 | 109 | 185 | 225 |
| P5 | -49 | -59 | -58 | -58 | -47 | -38 | -22 | 13 | 52 | 78 |




Small 12.10 Alternative 2B, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -11 | -21 | -5 | 18 | 47 | 81 | 113 | 159 | 184 | 219 |
| P75 | -21 | -34 | -29 | -19 | 10 | 37 | 70 | 103 | 123 | 138 |
| P50 | -29 | -44 | -41 | -32 | -6 | 15 | 48 | 78 | 84 | 103 |
| P25 | -39 | -56 | -55 | -47 | -28 | -8 | 20 | 42 | 52 | 61 |
| P5 | -59 | -72 | -69 | -68 | -55 | -44 | -23 | -3 | 2 | 4 |




Spat 12.10 Alternative 2B, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 99 | 114 | 190 | 211 | 239 | 299 | 354 | 431 | 429 | 413 |
| P75 | 1 | 18 | 71 | 108 | 123 | 182 | 211 | 228 | 224 | 198 |
| P50 | -7 | 4 | 51 | 78 | 97 | 135 | 173 | 188 | 179 | 155 |
| P25 | -18 | -11 | 34 | 59 | 78 | 115 | 151 | 165 | 150 | 130 |
| P5 | -39 | -32 | 15 | 27 | 51 | 88 | 119 | 125 | 106 | 89 |




Market 12.12 Alternative 2B, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 3 | 21 | 29 | 20 | 57 | 90 | 157 | 258 | 399 | 448 |
| P75 | -10 | -1 | 4 | -1 | 29 | 60 | 117 | 195 | 282 | 353 |
| P50 | -19 | -17 | -17 | -18 | 7 | 35 | 81 | 145 | 214 | 289 |
| P25 | -29 | -40 | -39 | -39 | -20 | 3 | 38 | 87 | 133 | 173 |
| P5 | -59 | -64 | -65 | -66 | -54 | -48 | -27 | 2 | 29 | 47 |



Small 12.12 Alternative 2B, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -12 | -23 | -13 | -2 | 35 | 62 | 94 | 133 | 169 | 185 |
| P75 | -21 | -38 | -32 | -24 | 8 | 29 | 59 | 93 | 113 | 125 |
| P50 | -30 | -45 | -43 | -35 | -12 | 12 | 39 | 69 | 88 | 95 |
| P25 | -40 | -55 | -64 | -48 | -31 | -13 | 11 | 38 | 49 | 54 |
| P5 | -59 | -69 | -71 | -69 | -58 | -47 | -27 | -6 | -4 | 2 |



Spat 12.12 Alternative 2B, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 69 | 78 | 147 | 172 | 192 | 247 | 307 | 354 | 402 | 377 |
| P75 | -5 | 8 | 58 | 88 | 109 | 148 | 180 | 211 | 208 | 181 |
| P50 | -11 | -3 | 43 | 69 | 89 | 128 | 164 | 179 | 188 | 147 |
| P25 | -20 | -15 | 31 | 54 | 74 | 108 | 146 | 160 | 144 | 124 |
| P5 | -41 | -33 | 13 | 27 | 47 | 79 | 115 | 122 | 103 | 84 |





Figure 12 . Whole Bay (continued)


Market 12.13 Alternative 3 (no harvest)

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | 58 | 110 | 150 | 149 | 187 | 181 | 192 | 205 | 218 | 215 |
| P75 | 40 | 75 | 98 | 98 | 105 | 109 | 111 | 108 | 103 | 102 |
| P50 | 26 | 45 | 56 | 53 | 65 | 64 | 60 | 60 | 59 | 56 |
| P25 | 9 | 0 | 5 | 9 | 23 | 24 | 19 | 17 | 12 | 8 |
| P5 | -37 | -42 | -41 | -43 | -43 | -42 | -46 | -46 | -46 | -45 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | -9 | -15 | 8 | 8 | 15 | 17 | 19 | 15 | 21 | 24 |
| P75 | -18 | -30 | -30 | -34 | -37 | -36 | -38 | -38 | -37 | -40 |
| P50 | -27 | -41 | -43 | -48 | -52 | -54 | -54 | -54 | -55 | -56 |
| P25 | -39 | -53 | -57 | -61 | -64 | -66 | -66 | -66 | -68 | -67 |
| P5 | -58 | -69 | -72 | -78 | -82 | -83 | -82 | -83 | -82 | -81 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | 129 | 125 | 137 | 165 | 168 | 149 | 133 | 137 | 129 | 107 |
| P75 | 1 | 6 | 2 | 14 | 17 | 15 | 13 | 11 | -2 | -10 |
| P50 | -7 | -11 | -18 | -18 | -18 | -17 | -19 | -20 | -31 | -53 |
| P25 | -18 | -28 | -35 | -39 | -38 | -37 | -37 | -38 | -47 | -58 |
| P5 | -42 | -50 | -53 | -61 | -60 | -58 | -58 | -57 | -65 | -66 |




Market 12.14 No harvest and Alt 2B

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | 58 | 109 | 146 | 158 | 204 | 255 | 340 | 472 | 622 | 782 |
| P75 | 40 | 75 | 98 | 108 | 138 | 174 | 238 | 339 | 457 | 588 |
| P50 | 27 | 47 | 55 | 60 | 87 | 123 | 177 | 258 | 388 | 487 |
| P25 | 12 | 1 | 6 | 13 | 39 | 69 | 111 | 173 | 245 | 327 |
| P5 | -39 | -40 | -40 | -42 | -31 | -22 | 6 | 39 | 89 | 132 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | -9 | -16 | 5 | 25 | 72 | 108 | 161 | 206 | 259 | 271 |
| P75 | -17 | -30 | -24 | -10 | 23 | 52 | 84 | 124 | 147 | 159 |
| P50 | -27 | -41 | -37 | -25 | 2 | 28 | 55 | 90 | 109 | 119 |
| P25 | -37 | -53 | -51 | -42 | -22 | -2 | 28 | 58 | 85 | 72 |
| P5 | -59 | -68 | -68 | -66 | -58 | -42 | -16 | 0 | 8 | 14 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | 98 | 159 | 247 | 311 | 356 | 417 | 508 | 574 | 529 | 513 |
| P75 | 5 | 31 | 89 | 138 | 181 | 205 | 251 | 274 | 281 | 238 |
| P50 | -2 | 18 | 87 | 100 | 123 | 159 | 200 | 212 | 201 | 179 |
| P25 | -11 | -2 | 44 | 72 | 92 | 128 | 168 | 180 | 186 | 144 |
| P5 | -39 | -28 | 21 | 33 | 53 | 88 | 127 | 127 | 118 | 102 |




Market 12.14 No harvest and Alt 2A

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | 58 | 108 | 148 | 154 | 199 | 258 | 361 | 479 | 653 | 787 |
| P75 | 39 | 72 | 93 | 105 | 138 | 180 | 247 | 348 | 483 | 612 |
| P50 | 27 | 47 | 56 | 63 | 95 | 124 | 179 | 271 | 383 | 486 |
| P25 | 15 | 0 | 11 | 19 | 41 | 69 | 115 | 173 | 254 | 343 |
| P5 | -37 | -43 | -42 | -42 | -31 | -15 | 4 | 49 | 98 | 161 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | -9 | -14 | 7 | 32 | 80 | 123 | 189 | 222 | 240 | 259 |
| P75 | -18 | -30 | -23 | -9 | 25 | 53 | 89 | 120 | 144 | 152 |
| P50 | -27 | -40 | -35 | -24 | 4 | 26 | 59 | 91 | 108 | 115 |
| P25 | -36 | -53 | -49 | -41 | -23 | -3 | 28 | 54 | 68 | 71 |
| P5 | -57 | -69 | -68 | -64 | -55 | -44 | -23 | 2 | 12 | 18 |



Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | 109 | 184 | 275 | 370 | 407 | 505 | 471 | 521 | 519 | 489 |
| P75 | 7 | 32 | 95 | 145 | 189 | 206 | 251 | 283 | 258 | 227 |
| P50 | -2 | 18 | 69 | 105 | 127 | 157 | 198 | 207 | 195 | 188 |
| P25 | -10 | -2 | 47 | 75 | 95 | 127 | 165 | 175 | 158 | 134 |
| P5 | -36 | -27 | 18 | 35 | 54 | 83 | 120 | 128 | 111 | 95 |




Figure 13 . Maryland





Figure 13 . Maryland (continued)


Market 13.5 Alternative 2A, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|------|------|
| P95 | 20 | -4 | 12 | 85 | 167 | 291 | 455 | 698 | 1000 | 1304 |
| P75 | 10 | -16 | -13 | 39 | 117 | 231 | 385 | 595 | 867 | 1136 |
| P50 | 3 | -24 | -26 | 18 | 87 | 189 | 318 | 506 | 736 | 971 |
| P25 | -5 | -47 | -47 | -13 | 42 | 120 | 228 | 372 | 540 | 889 |
| P5 | -44 | -68 | -69 | -49 | -20 | 25 | 91 | 181 | 283 | 385 |




Market 13.6 Alternative 2A, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|------|
| P95 | 6 | -29 | -18 | 45 | 126 | 247 | 402 | 628 | 887 | 1119 |
| P75 | -3 | -36 | -34 | 15 | 93 | 200 | 342 | 532 | 770 | 985 |
| P50 | -10 | -42 | -44 | -1 | 67 | 165 | 290 | 457 | 654 | 840 |
| P25 | -20 | -61 | -59 | -25 | 28 | 98 | 194 | 318 | 487 | 617 |
| P5 | -55 | -77 | -78 | -55 | -31 | 13 | 72 | 154 | 245 | 323 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|------|------|------|------|
| P95 | -61 | 144 | 278 | 384 | 637 | 747 | 1062 | 1425 | 1814 | 1712 |
| P75 | -64 | 23 | 149 | 265 | 423 | 616 | 907 | 1251 | 1423 | 1483 |
| P50 | -68 | 14 | 130 | 240 | 392 | 572 | 851 | 1174 | 1327 | 1375 |
| P25 | -69 | 2 | 109 | 208 | 362 | 511 | 779 | 1084 | 1181 | 1214 |
| P5 | -79 | -31 | 41 | 98 | 199 | 339 | 518 | 730 | 813 | 843 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|------|------|------|
| P95 | -82 | 124 | 241 | 333 | 494 | 688 | 992 | 1340 | 1522 | 1812 |
| P75 | -64 | 20 | 140 | 256 | 409 | 603 | 867 | 1213 | 1377 | 1438 |
| P50 | -67 | 11 | 122 | 231 | 382 | 568 | 835 | 1142 | 1292 | 1344 |
| P25 | -70 | 1 | 101 | 200 | 341 | 516 | 748 | 1028 | 1118 | 1194 |
| P5 | -81 | -34 | 35 | 89 | 187 | 328 | 514 | 700 | 781 | 808 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| P95 | 1558 | 1931 | 2184 | 2983 | 3954 | 5038 | 6770 | 7332 | 7587 | 8354 |
| P75 | 628 | 1063 | 1488 | 2168 | 2932 | 4207 | 5585 | 5907 | 6142 | 6360 |
| P50 | 594 | 993 | 1403 | 2064 | 2809 | 4050 | 5368 | 5848 | 5839 | 6021 |
| P25 | 583 | 948 | 1353 | 1994 | 2723 | 3952 | 5208 | 5480 | 5641 | 5814 |
| P5 | 621 | 890 | 1291 | 1919 | 2613 | 3829 | 5056 | 5268 | 5398 | 5506 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| P95 | 1497 | 1698 | 1935 | 2673 | 3514 | 4892 | 6059 | 6598 | 7116 | 7482 |
| P75 | 600 | 997 | 1437 | 2099 | 2854 | 4138 | 5473 | 5783 | 5956 | 6104 |
| P50 | 571 | 949 | 1365 | 2023 | 2749 | 4003 | 5311 | 5558 | 5722 | 5878 |
| P25 | 540 | 914 | 1322 | 1988 | 2684 | 3921 | 5184 | 5399 | 5553 | 5671 |
| P5 | 504 | 882 | 1272 | 1900 | 2610 | 3802 | 5020 | 5228 | 5344 | 5401 |




Market 13.7 Alternative 2A, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|------|
| P95 | -11 | -48 | -34 | 27 | 100 | 212 | 399 | 661 | 803 | 1018 |
| P75 | -18 | -62 | -48 | 0 | 72 | 176 | 308 | 490 | 703 | 863 |
| P50 | -23 | -67 | -58 | -14 | 52 | 141 | 284 | 421 | 600 | 751 |
| P25 | -33 | -71 | -68 | -36 | 11 | 84 | 180 | 298 | 440 | 549 |
| P5 | -59 | -82 | -81 | -61 | -34 | 5 | 71 | 162 | 267 | 337 |




Market 13.8 Alternative 2A, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | -26 | -61 | -45 | 11 | 85 | 193 | 340 | 533 | 752 | 933 |
| P75 | -32 | -65 | -58 | -8 | 62 | 160 | 296 | 470 | 653 | 808 |
| P50 | -38 | -68 | -62 | -20 | 42 | 131 | 251 | 399 | 568 | 680 |
| P25 | -41 | -77 | -71 | -40 | 10 | 78 | 168 | 272 | 401 | 478 |
| P5 | -66 | -86 | -82 | -62 | -33 | -1 | 45 | 116 | 187 | 263 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|------|------|------|
| P95 | -62 | 116 | 203 | 293 | 472 | 664 | 957 | 1307 | 1512 | 1826 |
| P75 | -65 | 15 | 130 | 243 | 399 | 590 | 875 | 1197 | 1355 | 1414 |
| P50 | -67 | 7 | 113 | 222 | 373 | 555 | 825 | 1138 | 1278 | 1319 |
| P25 | -71 | -3 | 92 | 187 | 335 | 497 | 747 | 1040 | 1143 | 1170 |
| P5 | -80 | -38 | 25 | 86 | 188 | 307 | 512 | 731 | 811 | 836 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|------|------|------|
| P95 | -63 | 92 | 176 | 272 | 441 | 650 | 940 | 1283 | 1458 | 1622 |
| P75 | -65 | 11 | 122 | 238 | 394 | 582 | 862 | 1188 | 1335 | 1389 |
| P50 | -68 | 3 | 109 | 217 | 368 | 551 | 813 | 1117 | 1259 | 1302 |
| P25 | -70 | -5 | 89 | 185 | 335 | 496 | 736 | 1008 | 1110 | 1134 |
| P5 | -81 | -38 | 24 | 87 | 198 | 302 | 487 | 697 | 782 | 802 |



Spat

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| P95 | 1407 | 1437 | 1691 | 2587 | 3295 | 4555 | 5931 | 6325 | 7022 | 7397 |
| P75 | 568 | 957 | 1389 | 2059 | 2800 | 4084 | 5413 | 5691 | 5903 | 6050 |
| P50 | 543 | 912 | 1333 | 1989 | 2722 | 3973 | 5247 | 5514 | 5687 | 5808 |
| P25 | 518 | 883 | 1298 | 1939 | 2659 | 3890 | 5137 | 5378 | 5504 | 5626 |
| P5 | 488 | 843 | 1258 | 1858 | 2598 | 3790 | 4998 | 5202 | 5297 | 5387 |



Spat

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| P95 | 1236 | 1276 | 1594 | 2352 | 3168 | 4411 | 5854 | 6326 | 6741 | 7318 |
| P75 | 535 | 928 | 1358 | 2023 | 2777 | 4029 | 5380 | 5668 | 5843 | 5980 |
| P50 | 513 | 887 | 1318 | 1988 | 2899 | 3939 | 5232 | 5477 | 5624 | 5739 |
| P25 | 498 | 865 | 1286 | 1928 | 2846 | 3871 | 5118 | 5348 | 5471 | 5571 |
| P5 | 470 | 832 | 1247 | 1881 | 2581 | 3787 | 4987 | 5185 | 5266 | 5359 |





Figure 13 . Maryland (continued)


Market 13.9 Alternative 2B, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|------|
| P95 | 20 | -5 | 10 | 88 | 159 | 298 | 428 | 657 | 942 | 1219 |
| P75 | 11 | -16 | -14 | 33 | 107 | 211 | 358 | 552 | 792 | 1048 |
| P50 | 4 | -23 | -26 | 13 | 77 | 168 | 292 | 463 | 661 | 855 |
| P25 | -6 | -48 | -48 | -17 | 30 | 100 | 188 | 318 | 457 | 612 |
| P5 | -48 | -68 | -70 | -54 | -31 | 9 | 80 | 133 | 224 | 324 |




Market 13.10 Alternative 2B, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|------|
| P95 | 6 | -28 | -19 | 43 | 114 | 221 | 373 | 584 | 827 | 1073 |
| P75 | -3 | -37 | -38 | 10 | 81 | 180 | 318 | 497 | 717 | 919 |
| P50 | -10 | -42 | -46 | -5 | 56 | 146 | 267 | 423 | 611 | 783 |
| P25 | -19 | -62 | -61 | -31 | 16 | 82 | 182 | 297 | 426 | 581 |
| P5 | -54 | -78 | -78 | -61 | -34 | 0 | 51 | 127 | 226 | 282 |




Small 13.9 Alternative 2B, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|------|------|------|------|
| P95 | -81 | 155 | 288 | 345 | 518 | 729 | 1089 | 1481 | 1988 | 1811 |
| P75 | -64 | 22 | 143 | 235 | 408 | 597 | 875 | 1200 | 1374 | 1449 |
| P50 | -68 | 13 | 124 | 232 | 378 | 554 | 818 | 1119 | 1274 | 1332 |
| P25 | -69 | 3 | 103 | 200 | 328 | 489 | 740 | 1005 | 1082 | 1148 |
| P5 | -81 | -35 | 28 | 83 | 178 | 298 | 481 | 662 | 749 | 788 |



Small 13.10 Alternative 2B, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|------|------|------|------|
| P95 | -81 | 118 | 233 | 314 | 488 | 685 | 1002 | 1380 | 1592 | 1745 |
| P75 | -64 | 18 | 132 | 245 | 398 | 584 | 868 | 1187 | 1345 | 1422 |
| P50 | -67 | 8 | 114 | 222 | 368 | 543 | 810 | 1112 | 1254 | 1313 |
| P25 | -70 | -3 | 83 | 189 | 325 | 487 | 733 | 993 | 1072 | 1163 |
| P5 | -81 | -37 | 27 | 88 | 178 | 295 | 487 | 677 | 761 | 778 |




Spot 13.9 Alternative 2B, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|-------|
| P95 | 1734 | 1763 | 2135 | 2916 | 3854 | 5530 | 7283 | 7812 | 8881 | 10133 |
| P75 | 829 | 1041 | 1485 | 2182 | 2930 | 4213 | 5834 | 5989 | 6383 | 6679 |
| P50 | 594 | 887 | 1408 | 2068 | 2794 | 4062 | 5390 | 5728 | 5893 | 6243 |
| P25 | 568 | 948 | 1358 | 1993 | 2720 | 3949 | 5239 | 5562 | 5730 | 5912 |
| P5 | 523 | 882 | 1292 | 1918 | 2818 | 3828 | 5081 | 5298 | 5455 | 5617 |




Spot 13.10 Alternative 2B, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| P95 | 1472 | 1543 | 1998 | 2673 | 3578 | 5134 | 6935 | 7665 | 8667 | 9670 |
| P75 | 598 | 988 | 1421 | 2090 | 2858 | 4122 | 5527 | 5898 | 6224 | 6484 |
| P50 | 568 | 944 | 1383 | 2009 | 2758 | 3988 | 5342 | 5681 | 5888 | 6095 |
| P25 | 539 | 910 | 1321 | 1958 | 2688 | 3910 | 5188 | 5481 | 5680 | 5831 |
| P5 | 502 | 880 | 1287 | 1895 | 2592 | 3804 | 5058 | 5278 | 5386 | 5488 |




Market 13.11 Alternative 2B, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | -10 | -48 | -33 | 22 | 93 | 193 | 340 | 532 | 781 | 971 |
| P75 | -17 | -53 | -49 | -2 | 63 | 157 | 289 | 455 | 657 | 830 |
| P50 | -23 | -57 | -58 | -18 | 42 | 127 | 244 | 378 | 549 | 685 |
| P25 | -31 | -70 | -67 | -37 | 6 | 71 | 155 | 264 | 374 | 485 |
| P5 | -59 | -82 | -81 | -64 | -42 | -7 | 38 | 108 | 211 | 264 |




Market 13.12 Alternative 2B, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | -26 | -61 | -46 | 8 | 73 | 171 | 320 | 512 | 718 | 877 |
| P75 | -31 | -65 | -67 | -11 | 51 | 143 | 271 | 437 | 618 | 789 |
| P50 | -38 | -67 | -62 | -22 | 32 | 118 | 227 | 380 | 518 | 644 |
| P25 | -42 | -77 | -71 | -42 | 0 | 64 | 146 | 249 | 363 | 441 |
| P5 | -66 | -88 | -83 | -67 | -41 | -14 | 41 | 105 | 177 | 218 |




Small 13.11 Alternative 2B, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|------|------|------|
| P95 | -62 | 105 | 193 | 283 | 447 | 655 | 964 | 1296 | 1527 | 1679 |
| P75 | -65 | 14 | 129 | 235 | 385 | 578 | 849 | 1157 | 1327 | 1393 |
| P50 | -67 | 5 | 111 | 214 | 369 | 537 | 793 | 1089 | 1225 | 1292 |
| P25 | -70 | -6 | 89 | 180 | 318 | 477 | 719 | 932 | 1082 | 1143 |
| P5 | -81 | -38 | 22 | 77 | 178 | 297 | 473 | 662 | 788 | 749 |




Small 13.12 Alternative 2B, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|------|------|------|
| P95 | -63 | 77 | 164 | 260 | 425 | 633 | 922 | 1289 | 1500 | 1623 |
| P75 | -65 | 9 | 118 | 229 | 379 | 568 | 843 | 1155 | 1313 | 1384 |
| P50 | -67 | 3 | 104 | 208 | 354 | 533 | 789 | 1081 | 1222 | 1287 |
| P25 | -71 | -6 | 83 | 178 | 316 | 481 | 708 | 968 | 1089 | 1100 |
| P5 | -81 | -40 | 19 | 75 | 178 | 284 | 469 | 685 | 732 | 780 |



Spot 13.11 Alternative 2B, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| P95 | 1315 | 1397 | 1706 | 2507 | 3387 | 4952 | 6373 | 7521 | 8244 | 8828 |
| P75 | 588 | 961 | 1385 | 2050 | 2813 | 4081 | 5452 | 5829 | 6055 | 6356 |
| P50 | 542 | 911 | 1330 | 1988 | 2724 | 3984 | 5288 | 5817 | 5799 | 5988 |
| P25 | 518 | 885 | 1298 | 1937 | 2680 | 3884 | 5164 | 5447 | 5602 | 5751 |
| P5 | 485 | 838 | 1257 | 1884 | 2685 | 3781 | 5032 | 5238 | 5342 | 5448 |



Spot 13.12 Alternative 2B, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| P95 | 1128 | 1185 | 1826 | 2396 | 3254 | 4654 | 6283 | 7089 | 7720 | 8280 |
| P75 | 533 | 911 | 1380 | 2029 | 2782 | 4050 | 5424 | 5783 | 6044 | 6252 |
| P50 | 514 | 883 | 1314 | 1972 | 2703 | 3947 | 5281 | 5570 | 5778 | 5920 |
| P25 | 486 | 883 | 1284 | 1929 | 2650 | 3889 | 5157 | 5409 | 5557 | 5692 |
| P5 | 470 | 831 | 1249 | 1881 | 2576 | 3779 | 5001 | 5230 | 5340 | 5401 |





Figure 13 . Maryland (continued)


Market 13.13 Alternative 3 (no harvest)

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 35 | 24 | 48 | 111 | 183 | 199 | 235 | 284 | 298 | 348 |
| P75 | 26 | 10 | 18 | 68 | 95 | 127 | 158 | 175 | 191 | 202 |
| P50 | 17 | -1 | -1 | 32 | 64 | 84 | 104 | 120 | 135 | 144 |
| P25 | 5 | -33 | -31 | -3 | 22 | 40 | 50 | 59 | 69 | 74 |
| P5 | -36 | -80 | -60 | -50 | -41 | -32 | -22 | -16 | -15 | -7 |




Market 13.14 No harvest and Alt 2A

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|------|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 35 | 23 | 40 | 124 | 224 | 369 | 569 | 825 | 1187 | 1630 |
| P75 | 25 | 9 | 16 | 71 | 159 | 283 | 452 | 688 | 989 | 1299 |
| P50 | 17 | -2 | -1 | 47 | 123 | 225 | 364 | 569 | 823 | 1069 |
| P25 | 8 | -33 | -31 | 8 | 81 | 135 | 249 | 390 | 589 | 797 |
| P5 | -34 | -60 | -59 | -42 | -15 | 27 | 90 | 199 | 317 | 452 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -60 | 173 | 230 | 242 | 264 | 266 | 252 | 299 | 383 | 375 |
| P75 | -63 | 31 | 101 | 110 | 117 | 127 | 132 | 134 | 140 | 149 |
| P50 | -68 | 20 | 81 | 84 | 87 | 92 | 97 | 99 | 101 | 102 |
| P25 | -69 | 7 | 57 | 58 | 62 | 65 | 69 | 66 | 71 | 70 |
| P5 | -80 | -28 | 3 | 7 | 5 | 10 | 14 | 16 | 16 | 17 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|------|------|------|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -60 | 159 | 298 | 432 | 621 | 814 | 1163 | 1510 | 1776 | 1907 |
| P75 | -63 | 29 | 159 | 280 | 444 | 637 | 932 | 1272 | 1445 | 1518 |
| P50 | -65 | 18 | 140 | 252 | 405 | 588 | 865 | 1191 | 1342 | 1397 |
| P25 | -68 | 3 | 115 | 219 | 351 | 505 | 782 | 1057 | 1175 | 1220 |
| P5 | -79 | -33 | 40 | 100 | 203 | 327 | 507 | 724 | 816 | 864 |




Spot

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 1923 | 1631 | 1730 | 1748 | 1797 | 1762 | 1847 | 2284 | 2177 | 2426 |
| P75 | 666 | 760 | 782 | 817 | 863 | 863 | 899 | 937 | 968 | 978 |
| P50 | 628 | 678 | 681 | 678 | 701 | 715 | 728 | 740 | 757 | 749 |
| P25 | 581 | 621 | 594 | 594 | 609 | 619 | 627 | 634 | 635 | 634 |
| P5 | 538 | 541 | 514 | 518 | 522 | 523 | 527 | 535 | 537 | 531 |




Spot

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 1724 | 2084 | 2644 | 3610 | 4293 | 6125 | 7049 | 8195 | 9121 | 9629 |
| P75 | 660 | 1115 | 1560 | 2275 | 3074 | 4338 | 5745 | 6083 | 6388 | 6672 |
| P50 | 628 | 1042 | 1462 | 2133 | 2881 | 4124 | 5483 | 5765 | 5985 | 6214 |
| P25 | 588 | 989 | 1394 | 2041 | 2767 | 4002 | 5285 | 5539 | 5734 | 5927 |
| P5 | 542 | 913 | 1309 | 1936 | 2645 | 3649 | 5085 | 5309 | 5423 | 5558 |




Market 13.14 No harvest and Alt 2B

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|------|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 36 | 23 | 39 | 118 | 217 | 341 | 625 | 787 | 1116 | 1478 |
| P75 | 26 | 11 | 18 | 71 | 152 | 268 | 428 | 647 | 928 | 1234 |
| P50 | 18 | 0 | -1 | 43 | 118 | 218 | 355 | 541 | 783 | 1015 |
| P25 | 8 | -33 | -33 | 3 | 61 | 138 | 236 | 374 | 573 | 746 |
| P5 | -40 | -61 | -62 | -47 | -23 | 16 | 88 | 169 | 298 | 399 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|------|------|------|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -60 | 157 | 285 | 400 | 575 | 834 | 1177 | 1575 | 1845 | 2058 |
| P75 | -63 | 26 | 154 | 271 | 428 | 620 | 909 | 1244 | 1434 | 1510 |
| P50 | -65 | 18 | 133 | 243 | 398 | 574 | 844 | 1149 | 1312 | 1390 |
| P25 | -69 | 4 | 107 | 207 | 343 | 509 | 757 | 1029 | 1164 | 1188 |
| P5 | -80 | -33 | 32 | 95 | 181 | 297 | 519 | 699 | 784 | 822 |



Spot

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|-------|-------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 1661 | 2035 | 2328 | 3447 | 4484 | 5928 | 8061 | 8686 | 10946 | 11441 |
| P75 | 657 | 1108 | 1556 | 2251 | 3075 | 4368 | 5787 | 6294 | 6636 | 7186 |
| P50 | 620 | 1039 | 1481 | 2121 | 2881 | 4171 | 5521 | 5894 | 6172 | 6504 |
| P25 | 579 | 987 | 1396 | 2036 | 2770 | 4017 | 5346 | 5643 | 5880 | 6108 |
| P5 | 535 | 904 | 1309 | 1929 | 2649 | 3853 | 5121 | 5370 | 5502 | 5715 |





Figure 14 . Virginia

Market 14.1 Alternative 1, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 64 | 147 | 181 | 140 | 136 | 123 | 111 | 98 | 84 | 66 |
| P75 | 42 | 100 | 118 | 83 | 73 | 58 | 33 | 14 | 1 | -9 |
| P50 | 24 | 61 | 73 | 40 | 34 | 11 | -6 | -21 | -30 | -37 |
| P25 | 2 | 18 | 16 | -7 | -10 | -21 | -38 | -46 | -55 | -62 |
| P5 | -43 | -37 | -33 | -52 | -54 | -64 | -73 | -80 | -85 | -87 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -4 | -24 | -24 | -17 | -14 | -21 | -23 | -24 | -27 | -33 |
| P75 | -14 | -38 | -46 | -51 | -57 | -60 | -61 | -63 | -64 | -65 |
| P50 | -24 | -47 | -57 | -63 | -68 | -73 | -74 | -75 | -75 | -75 |
| P25 | -35 | -60 | -69 | -75 | -79 | -83 | -82 | -82 | -84 | -84 |
| P5 | -55 | -75 | -82 | -87 | -91 | -94 | -94 | -94 | -94 | -94 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 64 | 78 | 83 | 73 | 71 | 62 | 62 | 55 | 45 | 18 |
| P75 | -14 | -15 | -20 | -19 | -21 | -21 | -23 | -22 | -38 | -43 |
| P50 | -22 | -28 | -37 | -39 | -42 | -43 | -44 | -44 | -55 | -60 |
| P25 | -32 | -43 | -51 | -58 | -57 | -58 | -68 | -67 | -67 | -70 |
| P5 | -53 | -63 | -67 | -74 | -72 | -71 | -72 | -70 | -79 | -80 |



Market 14.3 Alternative 1, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 36 | 97 | 106 | 80 | 68 | 58 | 35 | 26 | 20 | 17 |
| P75 | 18 | 58 | 61 | 22 | 21 | 3 | -15 | -25 | -32 | -34 |
| P50 | 4 | 27 | 24 | -8 | -8 | -25 | -38 | -45 | -52 | -55 |
| P25 | -17 | -9 | -13 | -35 | -37 | -47 | -67 | -64 | -71 | -74 |
| P5 | -52 | -49 | -47 | -67 | -69 | -76 | -83 | -88 | -89 | -91 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -7 | -27 | -26 | -27 | -32 | -38 | -42 | -42 | -43 | -44 |
| P75 | -17 | -40 | -49 | -57 | -63 | -69 | -69 | -69 | -69 | -70 |
| P50 | -25 | -50 | -61 | -68 | -73 | -78 | -79 | -79 | -79 | -79 |
| P25 | -37 | -61 | -72 | -78 | -82 | -85 | -85 | -86 | -86 | -87 |
| P5 | -57 | -76 | -83 | -89 | -93 | -95 | -95 | -95 | -95 | -95 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 60 | 65 | 35 | 33 | 24 | 26 | 28 | 20 | -2 | -10 |
| P75 | -19 | -25 | -35 | -37 | -36 | -35 | -34 | -37 | -49 | -54 |
| P50 | -25 | -38 | -47 | -52 | -52 | -52 | -62 | -63 | -63 | -67 |
| P25 | -38 | -49 | -57 | -63 | -63 | -62 | -61 | -61 | -71 | -74 |
| P5 | -58 | -68 | -69 | -77 | -75 | -74 | -72 | -73 | -81 | -81 |



Market 14.2 Alternative 1, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 60 | 121 | 144 | 97 | 84 | 66 | 62 | 61 | 48 | 45 |
| P75 | 31 | 75 | 86 | 48 | 42 | 25 | 1 | -15 | -19 | -23 |
| P50 | 14 | 42 | 43 | 12 | 8 | -8 | -27 | -38 | -45 | -51 |
| P25 | -5 | 4 | 2 | -22 | -23 | -36 | -48 | -59 | -65 | -71 |
| P5 | -50 | -44 | -42 | -59 | -61 | -70 | -79 | -85 | -88 | -90 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -5 | -28 | -18 | -19 | -33 | -30 | -30 | -31 | -33 | -35 |
| P75 | -15 | -39 | -47 | -53 | -59 | -65 | -66 | -66 | -67 | -67 |
| P50 | -25 | -49 | -58 | -65 | -70 | -75 | -78 | -77 | -78 | -78 |
| P25 | -35 | -61 | -70 | -75 | -80 | -83 | -84 | -85 | -85 | -86 |
| P5 | -57 | -78 | -83 | -88 | -92 | -94 | -94 | -95 | -95 | -94 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 73 | 83 | 48 | 53 | 71 | 63 | 43 | 36 | 18 | 5 |
| P75 | -15 | -17 | -28 | -27 | -29 | -28 | -31 | -32 | -44 | -51 |
| P50 | -23 | -32 | -42 | -48 | -48 | -48 | -49 | -60 | -60 | -65 |
| P25 | -33 | -48 | -54 | -60 | -61 | -60 | -69 | -69 | -70 | -73 |
| P5 | -58 | -65 | -68 | -75 | -73 | -73 | -72 | -72 | -81 | -81 |



Market 14.4 Alternative 1, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 23 | 77 | 88 | 40 | 45 | 29 | 18 | 18 | 17 | 12 |
| P75 | 7 | 44 | 46 | 6 | 7 | -10 | -23 | -28 | -34 | -37 |
| P50 | -7 | 17 | 15 | -18 | -17 | -32 | -44 | -60 | -64 | -68 |
| P25 | -25 | -15 | -20 | -43 | -42 | -62 | -62 | -66 | -71 | -75 |
| P5 | -57 | -53 | -55 | -69 | -72 | -77 | -83 | -87 | -90 | -91 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -8 | -28 | -31 | -34 | -38 | -40 | -42 | -45 | -47 | -48 |
| P75 | -17 | -40 | -50 | -58 | -65 | -70 | -70 | -70 | -71 | -71 |
| P50 | -27 | -51 | -61 | -68 | -74 | -79 | -79 | -79 | -79 | -80 |
| P25 | -38 | -62 | -71 | -78 | -83 | -86 | -86 | -86 | -86 | -87 |
| P5 | -58 | -77 | -83 | -89 | -93 | -95 | -95 | -95 | -95 | -95 |



Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 34 | 48 | 36 | 31 | 22 | 12 | 9 | 7 | 7 | -7 |
| P75 | -21 | -28 | -38 | -39 | -39 | -40 | -41 | -41 | -48 | -54 |
| P50 | -28 | -39 | -48 | -53 | -54 | -64 | -64 | -63 | -62 | -67 |
| P25 | -38 | -51 | -59 | -65 | -65 | -63 | -63 | -61 | -70 | -74 |
| P5 | -58 | -67 | -69 | -77 | -75 | -74 | -73 | -72 | -81 | -81 |



Figure 14 . Virginia (continued)
















Figure 14 . Virginia (continued)


Market 14.9 Alternative 2B, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 82 | 148 | 182 | 145 | 144 | 140 | 147 | 174 | 186 | 220 |
| P75 | 42 | 100 | 122 | 80 | 73 | 68 | 63 | 70 | 76 | 81 |
| P50 | 24 | 61 | 70 | 38 | 32 | 18 | 12 | 19 | 16 | 19 |
| P25 | 2 | 15 | 19 | -3 | -10 | -18 | -26 | -25 | -29 | -36 |
| P5 | -43 | -37 | -37 | -54 | -56 | -63 | -69 | -74 | -79 | -81 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -5 | -25 | -18 | -7 | 20 | 28 | 35 | 40 | 55 | 74 |
| P75 | -15 | -38 | -44 | -41 | -25 | -15 | -9 | -4 | 0 | 6 |
| P50 | -24 | -49 | -55 | -58 | -44 | -38 | -33 | -28 | -28 | -26 |
| P25 | -35 | -60 | -68 | -70 | -62 | -57 | -64 | -54 | -54 | -53 |
| P5 | -55 | -76 | -82 | -85 | -84 | -84 | -82 | -84 | -84 | -83 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 80 | 111 | 174 | 203 | 209 | 229 | 263 | 266 | 255 | 256 |
| P75 | -9 | 2 | 52 | 71 | 77 | 89 | 101 | 115 | 94 | 68 |
| P50 | -17 | -12 | 30 | 45 | 47 | 57 | 68 | 77 | 65 | 30 |
| P25 | -28 | -28 | 12 | 23 | 28 | 35 | 46 | 50 | 27 | 4 |
| P5 | -50 | -48 | -9 | -9 | -2 | 3 | 8 | 10 | -12 | -29 |




Market 14.11 Alternative 2B, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 38 | 97 | 113 | 80 | 72 | 67 | 88 | 119 | 160 | 188 |
| P75 | 20 | 59 | 68 | 24 | 26 | 13 | 16 | 35 | 44 | 58 |
| P50 | 3 | 29 | 28 | -8 | -8 | -18 | -18 | -9 | -8 | -1 |
| P25 | -18 | -9 | -12 | -36 | -36 | -44 | -45 | -46 | -47 | -48 |
| P5 | -54 | -52 | -61 | -67 | -68 | -74 | -80 | -84 | -85 | -84 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -6 | -27 | -19 | -13 | 9 | 12 | 19 | 36 | 43 | 69 |
| P75 | -18 | -39 | -45 | -48 | -31 | -24 | -18 | -11 | -8 | 2 |
| P50 | -28 | -50 | -58 | -60 | -48 | -42 | -37 | -34 | -33 | -27 |
| P25 | -38 | -62 | -70 | -73 | -65 | -61 | -68 | -69 | -57 | -54 |
| P5 | -58 | -76 | -83 | -88 | -86 | -85 | -84 | -88 | -82 | -82 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 70 | 89 | 133 | 144 | 148 | 183 | 222 | 247 | 243 | 223 |
| P75 | -13 | -4 | 37 | 63 | 59 | 71 | 83 | 99 | 83 | 61 |
| P50 | -21 | -19 | 20 | 33 | 38 | 48 | 58 | 67 | 47 | 23 |
| P25 | -32 | -33 | 4 | 13 | 18 | 28 | 39 | 43 | 24 | 1 |
| P5 | -63 | -61 | -13 | -16 | -9 | 1 | 5 | 7 | -8 | -27 |




Market 14.10 Alternative 2B, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 49 | 117 | 139 | 91 | 82 | 81 | 111 | 149 | 151 | 180 |
| P75 | 29 | 75 | 88 | 43 | 38 | 28 | 29 | 47 | 61 | 64 |
| P50 | 13 | 40 | 42 | 9 | 4 | -7 | -10 | 1 | 3 | 7 |
| P25 | -8 | -3 | -3 | -24 | -25 | -34 | -40 | -37 | -41 | -38 |
| P5 | -49 | -47 | -46 | -62 | -63 | -69 | -78 | -77 | -60 | -63 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -6 | -27 | -22 | -12 | 8 | 23 | 32 | 43 | 61 | 70 |
| P75 | -16 | -40 | -46 | -46 | -29 | -19 | -12 | -6 | -2 | 1 |
| P50 | -25 | -50 | -57 | -58 | -45 | -40 | -34 | -32 | -28 | -25 |
| P25 | -36 | -62 | -70 | -71 | -64 | -60 | -55 | -55 | -54 | -52 |
| P5 | -57 | -77 | -83 | -86 | -84 | -83 | -83 | -82 | -82 | -83 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 65 | 84 | 169 | 161 | 159 | 198 | 228 | 272 | 260 | 221 |
| P75 | -11 | -2 | 42 | 65 | 65 | 78 | 97 | 109 | 94 | 64 |
| P50 | -19 | -18 | 23 | 38 | 40 | 51 | 61 | 71 | 54 | 28 |
| P25 | -29 | -31 | 8 | 18 | 21 | 32 | 40 | 46 | 29 | 4 |
| P5 | -51 | -52 | -12 | -14 | -4 | 4 | 9 | 12 | -11 | -31 |




Market 14.12 Alternative 2B, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 25 | 82 | 89 | 43 | 57 | 46 | 69 | 113 | 148 | 159 |
| P75 | 7 | 46 | 47 | 8 | 11 | 2 | 7 | 25 | 38 | 49 |
| P50 | -7 | 19 | 15 | -17 | -13 | -24 | -25 | -13 | -8 | -5 |
| P25 | -21 | -14 | -18 | -41 | -41 | -48 | -50 | -44 | -47 | -49 |
| P5 | -56 | -48 | -54 | -70 | -70 | -77 | -82 | -84 | -86 | -85 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -6 | -27 | -28 | -27 | 1 | 10 | 16 | 23 | 41 | 48 |
| P75 | -17 | -41 | -47 | -50 | -32 | -26 | -20 | -15 | -10 | -1 |
| P50 | -28 | -50 | -58 | -61 | -50 | -43 | -39 | -38 | -33 | -29 |
| P25 | -38 | -61 | -69 | -72 | -67 | -61 | -58 | -58 | -57 | -55 |
| P5 | -57 | -75 | -82 | -87 | -85 | -87 | -85 | -84 | -84 | -83 |



Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 45 | 58 | 119 | 126 | 131 | 153 | 183 | 216 | 238 | 211 |
| P75 | -18 | -11 | 30 | 47 | 51 | 66 | 79 | 92 | 80 | 55 |
| P50 | -22 | -22 | 15 | 28 | 33 | 44 | 55 | 63 | 48 | 22 |
| P25 | -32 | -34 | 4 | 13 | 18 | 27 | 36 | 44 | 23 | 1 |
| P5 | -53 | -52 | -14 | -14 | -9 | -2 | 8 | 10 | -13 | -32 |




Figure 14 . Virginia (continued)













Figure 15 . Low Salinity


Market 15.1 Alternative 1, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | 43 | 93 | 133 | 145 | 166 | 178 | 177 | 181 | 181 | 181 |
| P75 | -10 | 16 | 49 | 79 | 116 | 129 | 127 | 129 | 127 | 129 |
| P50 | -17 | 2 | 31 | 67 | 86 | 92 | 91 | 88 | 89 | 86 |
| P25 | -24 | -19 | 2 | 15 | 41 | 48 | 43 | 39 | 34 | 22 |
| P5 | -68 | -56 | -42 | -41 | -29 | -33 | -39 | -44 | -50 | -57 |



Small 15.1 Alternative 1, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | -7 | -12 | -19 | -33 | -44 | -47 | -60 | -51 | -60 | -53 |
| P75 | -56 | -47 | -43 | -51 | -60 | -64 | -64 | -64 | -65 | -65 |
| P50 | -58 | -51 | -49 | -56 | -65 | -68 | -68 | -68 | -69 | -69 |
| P25 | -60 | -57 | -56 | -64 | -70 | -72 | -73 | -73 | -73 | -75 |
| P5 | -76 | -77 | -80 | -86 | -90 | -93 | -93 | -94 | -94 | -94 |




Spat 15.1 Alternative 1, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | -18 | -39 | -52 | -63 | -55 | -58 | -68 | -57 | -66 | -58 |
| P75 | -51 | -58 | -69 | -69 | -69 | -70 | -69 | -70 | -70 | -70 |
| P50 | -64 | -64 | -74 | -73 | -73 | -74 | -74 | -73 | -74 | -74 |
| P25 | -56 | -68 | -76 | -77 | -76 | -76 | -77 | -77 | -77 | -77 |
| P5 | -79 | -87 | -93 | -93 | -92 | -93 | -93 | -93 | -94 | -94 |




Market 15.3 Alternative 1, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | 18 | 45 | 66 | 69 | 101 | 112 | 115 | 122 | 130 | 138 |
| P75 | -28 | -13 | 11 | 35 | 71 | 81 | 81 | 86 | 93 | 96 |
| P50 | -34 | -25 | -5 | 18 | 47 | 53 | 64 | 55 | 63 | 61 |
| P25 | -41 | -40 | -25 | -13 | 11 | 13 | 13 | 12 | 9 | 8 |
| P5 | -75 | -66 | -57 | -55 | -48 | -50 | -52 | -63 | -64 | -60 |




Small 15.3 Alternative 1, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | -9 | -15 | -26 | -40 | -52 | -55 | -67 | -56 | -66 | -56 |
| P75 | -57 | -49 | -46 | -54 | -63 | -66 | -68 | -67 | -67 | -67 |
| P50 | -59 | -53 | -51 | -59 | -67 | -70 | -70 | -70 | -70 | -70 |
| P25 | -62 | -60 | -60 | -68 | -72 | -75 | -74 | -75 | -74 | -75 |
| P5 | -77 | -78 | -81 | -87 | -92 | -94 | -93 | -94 | -94 | -94 |




Spat 15.3 Alternative 1, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | -19 | -43 | -60 | -60 | -61 | -61 | -59 | -61 | -64 | -62 |
| P75 | -62 | -68 | -72 | -72 | -72 | -72 | -71 | -72 | -72 | -72 |
| P50 | -64 | -66 | -75 | -75 | -75 | -75 | -75 | -75 | -75 | -75 |
| P25 | -57 | -70 | -78 | -78 | -77 | -78 | -77 | -78 | -78 | -78 |
| P5 | -79 | -88 | -94 | -94 | -94 | -94 | -94 | -94 | -94 | -94 |




Market 15.2 Alternative 1, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | 29 | 65 | 93 | 103 | 126 | 135 | 134 | 142 | 149 | 152 |
| P75 | -19 | -2 | 29 | 54 | 88 | 99 | 99 | 101 | 108 | 108 |
| P50 | -25 | -14 | 10 | 32 | 62 | 70 | 66 | 62 | 68 | 67 |
| P25 | -32 | -31 | -15 | 1 | 24 | 28 | 20 | 14 | 15 | 8 |
| P5 | -72 | -63 | -50 | -48 | -42 | -48 | -49 | -56 | -60 | -62 |




Small 15.2 Alternative 1, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | -6 | -15 | -20 | -36 | -48 | -63 | -60 | -60 | -62 | -53 |
| P75 | -56 | -48 | -44 | -52 | -61 | -65 | -68 | -68 | -68 | -68 |
| P50 | -59 | -52 | -60 | -57 | -68 | -69 | -70 | -69 | -69 | -70 |
| P25 | -61 | -59 | -58 | -64 | -70 | -74 | -74 | -75 | -74 | -75 |
| P5 | -77 | -79 | -80 | -86 | -91 | -94 | -94 | -94 | -94 | -94 |




Spat 15.2 Alternative 1, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | -19 | -42 | -59 | -68 | -55 | -69 | -68 | -60 | -61 | -60 |
| P75 | -61 | -69 | -71 | -70 | -71 | -71 | -71 | -71 | -72 | -72 |
| P50 | -64 | -64 | -74 | -74 | -74 | -74 | -75 | -75 | -75 | -75 |
| P25 | -66 | -70 | -77 | -77 | -77 | -77 | -77 | -78 | -78 | -78 |
| P5 | -80 | -88 | -93 | -93 | -93 | -94 | -93 | -94 | -94 | -94 |



Market 15.4 Alternative 1, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | 2 | 33 | 63 | 66 | 90 | 99 | 107 | 116 | 126 | 127 |
| P75 | -37 | -23 | 2 | 26 | 61 | 71 | 75 | 83 | 87 | 92 |
| P50 | -43 | -34 | -13 | 6 | 38 | 45 | 48 | 53 | 57 | 58 |
| P25 | -48 | -48 | -32 | -19 | 5 | 9 | 8 | 12 | 9 | 3 |
| P5 | -78 | -69 | -58 | -56 | -51 | -51 | -58 | -61 | -63 | -66 |



Small 15.4 Alternative 1, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | -11 | -16 | -25 | -42 | -52 | -66 | -66 | -67 | -67 | -58 |
| P75 | -57 | -50 | -47 | -55 | -64 | -67 | -67 | -67 | -67 | -67 |
| P50 | -60 | -54 | -52 | -59 | -68 | -71 | -70 | -70 | -70 | -70 |
| P25 | -62 | -60 | -60 | -68 | -72 | -75 | -74 | -74 | -75 | -75 |
| P5 | -77 | -79 | -80 | -87 | -91 | -93 | -94 | -94 | -94 | -94 |



Spat 15.4 Alternative 1, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| P95 | -21 | -45 | -58 | -61 | -61 | -63 | -63 | -61 | -59 | -60 |
| P75 | -62 | -60 | -72 | -72 | -72 | -73 | -72 | -73 | -72 | -72 |
| P50 | -66 | -68 | -75 | -75 | -75 | -75 | -75 | -75 | -75 | -75 |
| P25 | -67 | -70 | -78 | -78 | -77 | -78 | -78 | -78 | -77 | -78 |
| P5 | -79 | -88 | -94 | -94 | -94 | -94 | -94 | -94 | -94 | -94 |





Figure 15 . Low Salinity (continued)


Market 15.5 Alternative 2A, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 42 | 91 | 129 | 183 | 199 | 294 | 407 | 585 | 806 | 1039 |
| P75 | -11 | 16 | 48 | 85 | 154 | 233 | 338 | 493 | 701 | 897 |
| P50 | -17 | 2 | 27 | 60 | 123 | 191 | 278 | 421 | 593 | 762 |
| P25 | -23 | -19 | -2 | 19 | 85 | 113 | 180 | 277 | 389 | 474 |
| P5 | -67 | -55 | -43 | -44 | -31 | -21 | -1 | 40 | 102 | 158 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -7 | -12 | -8 | -4 | 7 | 39 | 87 | 150 | 184 | 195 |
| P75 | -58 | -49 | -38 | -24 | -8 | 19 | 86 | 119 | 144 | 153 |
| P50 | -68 | -63 | -42 | -31 | -15 | 12 | 67 | 105 | 127 | 133 |
| P25 | -61 | -58 | -60 | -40 | -23 | 0 | 43 | 86 | 98 | 97 |
| P5 | -78 | -78 | -81 | -85 | -82 | -71 | -47 | -28 | -22 | -21 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -19 | -28 | -23 | 1 | 27 | 71 | 120 | 134 | 162 | 167 |
| P75 | -63 | -47 | -42 | -18 | 7 | 52 | 96 | 108 | 112 | 118 |
| P50 | -68 | -64 | -47 | -24 | 1 | 44 | 82 | 91 | 93 | 97 |
| P25 | -68 | -69 | -61 | -28 | -4 | 37 | 74 | 81 | 81 | 84 |
| P5 | -81 | -88 | -92 | -82 | -68 | -39 | -22 | -21 | -20 | -19 |




Market 15.7 Alternative 2A, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 18 | 47 | 60 | 78 | 140 | 217 | 318 | 488 | 654 | 809 |
| P75 | -29 | -14 | 9 | 38 | 103 | 173 | 267 | 403 | 568 | 698 |
| P50 | -34 | -28 | -7 | 18 | 78 | 138 | 223 | 341 | 478 | 586 |
| P25 | -43 | -40 | -28 | -10 | 32 | 74 | 134 | 221 | 316 | 383 |
| P5 | -74 | -66 | -61 | -68 | -49 | -41 | -22 | 23 | 76 | 115 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -10 | -14 | -16 | -13 | 2 | 32 | 79 | 135 | 167 | 188 |
| P75 | -67 | -61 | -39 | -28 | -11 | 16 | 82 | 112 | 137 | 145 |
| P50 | -69 | -66 | -44 | -34 | -17 | 10 | 63 | 100 | 120 | 128 |
| P25 | -62 | -60 | -62 | -44 | -25 | -1 | 38 | 83 | 92 | 88 |
| P5 | -77 | -79 | -83 | -85 | -82 | -72 | -48 | -26 | -21 | -20 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -20 | -31 | -29 | -7 | 21 | 65 | 112 | 130 | 142 | 149 |
| P75 | -63 | -60 | -46 | -21 | 6 | 49 | 83 | 103 | 107 | 115 |
| P50 | -67 | -68 | -49 | -25 | -1 | 42 | 80 | 88 | 90 | 94 |
| P25 | -69 | -61 | -62 | -30 | -8 | 37 | 72 | 79 | 78 | 82 |
| P5 | -82 | -88 | -83 | -82 | -69 | -40 | -23 | -21 | -20 | -18 |




Market 15.6 Alternative 2A, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 29 | 63 | 90 | 105 | 178 | 259 | 362 | 517 | 718 | 897 |
| P75 | -20 | -1 | 28 | 58 | 128 | 199 | 298 | 440 | 618 | 777 |
| P50 | -26 | -13 | 9 | 37 | 95 | 162 | 248 | 375 | 516 | 661 |
| P25 | -33 | -33 | -17 | 4 | 51 | 95 | 151 | 229 | 324 | 423 |
| P5 | -73 | -63 | -52 | -54 | -45 | -32 | -17 | 33 | 73 | 125 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -8 | -15 | -14 | -10 | 10 | 32 | 79 | 139 | 188 | 181 |
| P75 | -56 | -50 | -37 | -28 | -9 | 18 | 64 | 114 | 138 | 149 |
| P50 | -69 | -64 | -43 | -32 | -15 | 11 | 64 | 101 | 121 | 130 |
| P25 | -61 | -60 | -61 | -40 | -24 | 1 | 37 | 77 | 84 | 101 |
| P5 | -77 | -80 | -81 | -86 | -83 | -72 | -48 | -26 | -23 | -21 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -19 | -29 | -27 | -3 | 28 | 66 | 118 | 134 | 133 | 149 |
| P75 | -63 | -49 | -43 | -20 | 7 | 51 | 92 | 105 | 103 | 114 |
| P50 | -68 | -65 | -48 | -25 | 0 | 43 | 81 | 88 | 90 | 98 |
| P25 | -68 | -69 | -61 | -28 | -5 | 38 | 72 | 78 | 78 | 82 |
| P5 | -82 | -88 | -82 | -82 | -69 | -39 | -24 | -23 | -22 | -19 |




Market 15.8 Alternative 2A, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 1 | 27 | 42 | 63 | 125 | 194 | 298 | 444 | 610 | 745 |
| P75 | -37 | -24 | -2 | 27 | 92 | 160 | 254 | 385 | 523 | 641 |
| P50 | -43 | -34 | -18 | 9 | 64 | 128 | 213 | 323 | 448 | 531 |
| P25 | -48 | -46 | -35 | -15 | 29 | 70 | 124 | 193 | 275 | 307 |
| P5 | -77 | -69 | -62 | -63 | -51 | -48 | -34 | 6 | 41 | 86 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -10 | -17 | -17 | -13 | -1 | 29 | 77 | 134 | 159 | 176 |
| P75 | -67 | -62 | -39 | -29 | -12 | 15 | 60 | 111 | 133 | 141 |
| P50 | -69 | -65 | -45 | -34 | -18 | 9 | 62 | 97 | 117 | 124 |
| P25 | -62 | -60 | -63 | -44 | -25 | -2 | 37 | 74 | 79 | 86 |
| P5 | -77 | -79 | -82 | -85 | -82 | -73 | -50 | -29 | -27 | -24 |



Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -20 | -32 | -32 | -7 | 18 | 64 | 114 | 135 | 132 | 146 |
| P75 | -63 | -60 | -46 | -21 | 4 | 49 | 82 | 103 | 103 | 111 |
| P50 | -67 | -68 | -49 | -28 | -2 | 41 | 79 | 88 | 89 | 92 |
| P25 | -69 | -60 | -62 | -30 | -5 | 38 | 71 | 77 | 75 | 78 |
| P5 | -81 | -88 | -83 | -82 | -69 | -40 | -24 | -24 | -22 | -19 |





Figure 15 . Low Salinity (continued)


Market 15.9 Alternative 2B, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 43 | 93 | 133 | 148 | 199 | 274 | 370 | 511 | 708 | 894 |
| P75 | -10 | 16 | 48 | 79 | 142 | 207 | 298 | 427 | 590 | 768 |
| P50 | -16 | 3 | 27 | 65 | 108 | 163 | 242 | 353 | 494 | 623 |
| P25 | -23 | -20 | -1 | 18 | 54 | 94 | 148 | 221 | 282 | 404 |
| P5 | -68 | -58 | -44 | -47 | -38 | -34 | -18 | 10 | 82 | 107 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -8 | -12 | -8 | -7 | 4 | 29 | 71 | 123 | 143 | 160 |
| P75 | -58 | -50 | -39 | -30 | -15 | 8 | 48 | 89 | 111 | 118 |
| P50 | -68 | -64 | -45 | -37 | -24 | -2 | 38 | 78 | 94 | 100 |
| P25 | -61 | -60 | -63 | -48 | -34 | -13 | 23 | 52 | 48 | 70 |
| P5 | -78 | -79 | -82 | -87 | -87 | -77 | -55 | -45 | -40 | -38 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -19 | -27 | -29 | -4 | 17 | 51 | 97 | 110 | 118 | 134 |
| P75 | -64 | -61 | -44 | -24 | 0 | 35 | 72 | 81 | 82 | 90 |
| P50 | -67 | -68 | -63 | -34 | -12 | 24 | 58 | 82 | 85 | 89 |
| P25 | -60 | -63 | -67 | -38 | -18 | 19 | 49 | 51 | 43 | 55 |
| P5 | -81 | -90 | -84 | -87 | -73 | -47 | -39 | -40 | -39 | -36 |




Market 15.11 Alternative 2B, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 17 | 47 | 67 | 75 | 134 | 194 | 291 | 417 | 579 | 722 |
| P75 | -28 | -12 | 8 | 33 | 89 | 151 | 236 | 348 | 486 | 602 |
| P50 | -34 | -28 | -7 | 15 | 83 | 118 | 191 | 285 | 400 | 491 |
| P25 | -41 | -41 | -27 | -12 | 24 | 57 | 111 | 160 | 244 | 315 |
| P5 | -74 | -65 | -58 | -60 | -53 | -51 | -40 | -11 | 35 | 67 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -8 | -14 | -16 | -13 | 1 | 23 | 65 | 111 | 137 | 154 |
| P75 | -67 | -62 | -40 | -32 | -17 | 6 | 43 | 84 | 108 | 113 |
| P50 | -69 | -68 | -47 | -39 | -27 | -4 | 33 | 71 | 87 | 95 |
| P25 | -62 | -61 | -65 | -50 | -36 | -14 | 21 | 34 | 64 | 71 |
| P5 | -77 | -80 | -83 | -87 | -88 | -77 | -54 | -43 | -39 | -40 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -19 | -30 | -33 | -11 | 13 | 50 | 96 | 107 | 117 | 130 |
| P75 | -64 | -62 | -48 | -28 | -1 | 33 | 71 | 78 | 78 | 85 |
| P50 | -68 | -68 | -65 | -35 | -13 | 22 | 64 | 80 | 82 | 85 |
| P25 | -61 | -63 | -68 | -39 | -18 | 17 | 47 | 38 | 51 | 53 |
| P5 | -82 | -90 | -84 | -88 | -74 | -47 | -40 | -41 | -38 | -36 |




Market 15.10 Alternative 2B, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 28 | 61 | 91 | 100 | 156 | 238 | 331 | 458 | 634 | 797 |
| P75 | -20 | -2 | 23 | 48 | 109 | 172 | 257 | 379 | 529 | 688 |
| P50 | -25 | -14 | 6 | 29 | 79 | 136 | 215 | 320 | 449 | 557 |
| P25 | -32 | -32 | -18 | -2 | 30 | 89 | 131 | 192 | 275 | 375 |
| P5 | -72 | -63 | -50 | -54 | -48 | -42 | -28 | 5 | 54 | 84 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -9 | -18 | -11 | -11 | -2 | 29 | 68 | 115 | 143 | 158 |
| P75 | -56 | -51 | -40 | -31 | -17 | 8 | 45 | 90 | 109 | 114 |
| P50 | -69 | -65 | -48 | -39 | -25 | -3 | 35 | 74 | 91 | 98 |
| P25 | -61 | -61 | -55 | -47 | -35 | -14 | 23 | 55 | 57 | 72 |
| P5 | -77 | -80 | -81 | -88 | -87 | -77 | -57 | -43 | -38 | -39 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -21 | -33 | -31 | -13 | 14 | 52 | 98 | 106 | 129 | 128 |
| P75 | -64 | -62 | -48 | -25 | -1 | 34 | 72 | 81 | 84 | 88 |
| P50 | -68 | -68 | -64 | -35 | -13 | 24 | 58 | 81 | 84 | 87 |
| P25 | -60 | -64 | -68 | -39 | -18 | 18 | 49 | 51 | 49 | 58 |
| P5 | -81 | -91 | -84 | -88 | -73 | -48 | -38 | -40 | -39 | -37 |




Market 15.12 Alternative 2B, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 2 | 27 | 49 | 64 | 123 | 178 | 271 | 398 | 550 | 650 |
| P75 | -37 | -23 | -2 | 23 | 81 | 135 | 221 | 332 | 457 | 568 |
| P50 | -43 | -34 | -18 | 8 | 55 | 108 | 179 | 274 | 378 | 483 |
| P25 | -47 | -46 | -34 | -21 | 14 | 52 | 96 | 164 | 233 | 287 |
| P5 | -77 | -68 | -62 | -63 | -53 | -52 | -41 | -6 | 30 | 52 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -10 | -17 | -16 | -16 | -2 | 21 | 59 | 108 | 138 | 147 |
| P75 | -67 | -62 | -41 | -33 | -18 | 3 | 42 | 83 | 103 | 111 |
| P50 | -69 | -68 | -47 | -40 | -27 | -4 | 33 | 70 | 87 | 93 |
| P25 | -62 | -61 | -64 | -49 | -35 | -14 | 19 | 51 | 63 | 65 |
| P5 | -77 | -79 | -83 | -88 | -87 | -78 | -57 | -42 | -41 | -39 |



Spat

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | -21 | -35 | -34 | -12 | 11 | 47 | 94 | 103 | 115 | 124 |
| P75 | -65 | -63 | -48 | -28 | -2 | 31 | 69 | 74 | 78 | 84 |
| P50 | -68 | -69 | -65 | -38 | -14 | 22 | 64 | 69 | 81 | 84 |
| P25 | -60 | -63 | -68 | -39 | -18 | 17 | 47 | 50 | 51 | 53 |
| P5 | -81 | -90 | -84 | -88 | -74 | -48 | -41 | -41 | -39 | -37 |




Figure 15 . Low Salinity (continued)












Figure 16 . Medium Salinity















Figure 16 . Medium Salinity (continued)















Figure 16 . Medium Salinity (continued)















Figure 16 . Medium Salinity (continued)












Figure 17 . High Salinity





Figure 17 . High Salinity (continued)


Market 17.5 Alternative 2A, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 110 | 131 | 124 | 72 | 68 | 93 | 136 | 170 | 216 | 228 |
| P75 | 85 | 88 | 31 | -1 | -10 | -9 | 10 | 45 | 64 | 71 |
| P50 | 58 | 18 | -17 | -44 | -51 | -47 | -41 | -19 | -13 | -10 |
| P25 | 14 | -19 | -49 | -71 | -80 | -80 | -78 | -75 | -68 | -70 |
| P5 | -30 | -58 | -81 | -91 | -94 | -94 | -95 | -95 | -95 | -96 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 227 | 143 | 281 | 409 | 594 | 730 | 708 | 709 | 792 | 798 |
| P75 | 178 | 70 | 90 | 150 | 277 | 337 | 355 | 386 | 395 | 428 |
| P50 | 138 | 20 | 25 | 63 | 141 | 171 | 182 | 188 | 228 | 222 |
| P25 | 47 | -38 | -27 | -8 | 20 | 28 | 41 | 40 | 66 | 41 |
| P5 | -43 | -70 | -81 | -80 | -68 | -67 | -70 | -77 | -78 | -88 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 1873 | 2484 | 3271 | 4140 | 4052 | 3750 | 4180 | 4148 | 3478 | 2899 |
| P75 | 780 | 1073 | 1747 | 1921 | 2010 | 2155 | 2340 | 2318 | 1988 | 1522 |
| P50 | 647 | 859 | 1383 | 1430 | 1452 | 1639 | 1867 | 1898 | 1420 | 974 |
| P25 | 484 | 831 | 978 | 983 | 914 | 947 | 1057 | 1015 | 900 | 820 |
| P5 | 141 | 204 | 139 | 153 | 108 | 83 | 12 | -40 | -37 | -59 |




Market 17.6 Alternative 2A, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 89 | 101 | 84 | 33 | 21 | 32 | 83 | 143 | 179 | 174 |
| P75 | 68 | 44 | 18 | -22 | -31 | -28 | -7 | 19 | 33 | 38 |
| P50 | 42 | -1 | -32 | -55 | -65 | -59 | -51 | -37 | -23 | -29 |
| P25 | 3 | -34 | -58 | -77 | -83 | -81 | -78 | -79 | -68 | -76 |
| P5 | -40 | -71 | -83 | -92 | -95 | -95 | -88 | -86 | -86 | -88 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 223 | 135 | 228 | 316 | 501 | 581 | 595 | 635 | 721 | 752 |
| P75 | 175 | 89 | 84 | 130 | 243 | 279 | 303 | 319 | 382 | 363 |
| P50 | 133 | 18 | 19 | 54 | 122 | 143 | 164 | 181 | 202 | 181 |
| P25 | 37 | -39 | -31 | -15 | 1 | 15 | 34 | 12 | 67 | 16 |
| P5 | -43 | -72 | -85 | -84 | -67 | -67 | -78 | -81 | -82 | -80 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 1719 | 1974 | 3121 | 3202 | 3057 | 3408 | 3585 | 3477 | 3219 | 2440 |
| P75 | 723 | 981 | 1838 | 1720 | 1754 | 1914 | 2077 | 2180 | 1816 | 1359 |
| P50 | 813 | 802 | 1307 | 1308 | 1275 | 1346 | 1517 | 1571 | 1311 | 845 |
| P25 | 447 | 583 | 933 | 891 | 773 | 907 | 1069 | 988 | 886 | 400 |
| P5 | 123 | 153 | 98 | 94 | 50 | 37 | -8 | -45 | -54 | -71 |




Market 17.7 Alternative 2A, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 71 | 68 | 67 | 7 | 2 | 18 | 82 | 124 | 156 | 183 |
| P75 | 49 | 23 | -2 | -37 | -44 | -38 | -20 | 8 | 22 | 28 |
| P50 | 25 | -18 | -42 | -66 | -71 | -67 | -57 | -43 | -35 | -37 |
| P25 | -13 | -44 | -65 | -81 | -86 | -86 | -85 | -81 | -78 | -80 |
| P5 | -47 | -75 | -88 | -95 | -98 | -98 | -96 | -97 | -96 | -98 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 224 | 120 | 212 | 300 | 463 | 494 | 596 | 575 | 651 | 658 |
| P75 | 171 | 81 | 70 | 113 | 214 | 260 | 290 | 298 | 340 | 360 |
| P50 | 120 | 11 | 10 | 33 | 102 | 118 | 142 | 149 | 178 | 172 |
| P25 | 28 | -43 | -37 | -29 | -7 | 7 | 11 | 9 | 46 | 17 |
| P5 | -45 | -71 | -87 | -71 | -73 | -78 | -81 | -88 | -87 | -95 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 1558 | 1897 | 2481 | 2835 | 3008 | 2968 | 3199 | 3378 | 3072 | 2708 |
| P75 | 684 | 900 | 1549 | 1870 | 1888 | 1838 | 1968 | 2048 | 1799 | 1367 |
| P50 | 573 | 724 | 1243 | 1240 | 1145 | 1280 | 1418 | 1493 | 1229 | 810 |
| P25 | 409 | 545 | 890 | 836 | 722 | 851 | 969 | 1008 | 802 | 348 |
| P5 | 81 | 122 | 69 | 68 | 32 | -12 | -33 | -69 | -82 | -85 |




Market 17.8 Alternative 2A, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 50 | 52 | 40 | -4 | -9 | 9 | 56 | 99 | 140 | 158 |
| P75 | 30 | 12 | -14 | -48 | -50 | -45 | -28 | 1 | 17 | 21 |
| P50 | 11 | -24 | -48 | -88 | -72 | -72 | -62 | -47 | -37 | -40 |
| P25 | -22 | -49 | -65 | -83 | -87 | -87 | -86 | -84 | -78 | -82 |
| P5 | -54 | -77 | -86 | -95 | -96 | -96 | -97 | -97 | -97 | -97 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 221 | 122 | 199 | 282 | 403 | 442 | 534 | 522 | 649 | 641 |
| P75 | 186 | 59 | 83 | 94 | 194 | 222 | 251 | 278 | 309 | 315 |
| P50 | 121 | 13 | 11 | 28 | 98 | 101 | 125 | 145 | 174 | 180 |
| P25 | 30 | -39 | -37 | -28 | -4 | -6 | 7 | 7 | 40 | 0 |
| P5 | -45 | -72 | -87 | -71 | -72 | -77 | -85 | -88 | -88 | -93 |



Spat


| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 1491 | 1757 | 2409 | 2385 | 2489 | 2887 | 3141 | 3175 | 2921 | 2523 |
| P75 | 848 | 854 | 1528 | 1541 | 1588 | 1763 | 1911 | 2015 | 1725 | 1318 |
| P50 | 533 | 898 | 1171 | 1147 | 1080 | 1225 | 1345 | 1493 | 1225 | 780 |
| P25 | 370 | 525 | 879 | 780 | 772 | 817 | 957 | 990 | 824 | 384 |
| P5 | 87 | 87 | 65 | 54 | 19 | -10 | -45 | -80 | -87 | -82 |



Figure 17 . High Salinity (continued)


Market 17.9 Alternative 2B, 20% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 110 | 133 | 123 | 79 | 71 | 76 | 128 | 199 | 222 | 291 |
| P75 | 84 | 85 | 42 | 3 | -12 | -8 | 15 | 50 | 77 | 95 |
| P50 | 57 | 15 | -19 | -43 | -50 | -45 | -36 | -13 | 5 | 11 |
| P25 | 14 | -22 | -60 | -71 | -78 | -76 | -75 | -69 | -63 | -65 |
| P5 | -32 | -65 | -79 | -69 | -64 | -64 | -64 | -64 | -63 | -64 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 227 | 145 | 242 | 390 | 622 | 596 | 858 | 748 | 837 | 897 |
| P75 | 178 | 74 | 95 | 147 | 276 | 309 | 345 | 410 | 464 | 470 |
| P50 | 134 | 22 | 30 | 62 | 139 | 159 | 190 | 248 | 283 | 273 |
| P25 | 32 | -39 | -26 | -9 | 18 | 36 | 58 | 76 | 105 | 76 |
| P5 | -43 | -71 | -82 | -60 | -63 | -67 | -71 | -61 | -78 | -82 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 1964 | 2242 | 3259 | 3306 | 3227 | 3722 | 4101 | 4178 | 3754 | 3247 |
| P75 | 759 | 1048 | 1737 | 1808 | 1909 | 2186 | 2368 | 2534 | 2233 | 1789 |
| P50 | 648 | 836 | 1398 | 1400 | 1333 | 1672 | 1783 | 1903 | 1815 | 1222 |
| P25 | 448 | 594 | 936 | 919 | 818 | 1036 | 1234 | 1311 | 1119 | 866 |
| P5 | 139 | 179 | 132 | 110 | 41 | 62 | 12 | -25 | -27 | -51 |




Market 17.10 Alternative 2B, 40% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 89 | 103 | 84 | 38 | 29 | 34 | 87 | 183 | 206 | 247 |
| P75 | 65 | 43 | 10 | -22 | -32 | -29 | -5 | 38 | 59 | 82 |
| P50 | 40 | -1 | -33 | -57 | -63 | -59 | -49 | -27 | -13 | 7 |
| P25 | -1 | -33 | -61 | -78 | -84 | -82 | -80 | -75 | -68 | -65 |
| P5 | -41 | -72 | -84 | -93 | -95 | -94 | -95 | -95 | -95 | -96 |




Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 221 | 132 | 214 | 334 | 533 | 515 | 677 | 688 | 772 | 838 |
| P75 | 171 | 88 | 79 | 123 | 251 | 287 | 318 | 386 | 431 | 468 |
| P50 | 130 | 19 | 16 | 50 | 124 | 133 | 181 | 221 | 258 | 281 |
| P25 | 23 | -39 | -33 | -16 | 11 | 14 | 25 | 56 | 79 | 87 |
| P5 | -44 | -70 | -68 | -69 | -68 | -78 | -61 | -87 | -67 | -69 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 1766 | 2006 | 3000 | 2922 | 2843 | 3307 | 3396 | 3779 | 3536 | 3178 |
| P75 | 721 | 988 | 1808 | 1653 | 1784 | 1969 | 2224 | 2418 | 2116 | 1742 |
| P50 | 608 | 792 | 1303 | 1244 | 1191 | 1407 | 1611 | 1775 | 1618 | 1143 |
| P25 | 427 | 597 | 899 | 868 | 734 | 918 | 1116 | 1191 | 1004 | 707 |
| P5 | 112 | 188 | 91 | 61 | 27 | 0 | -16 | -54 | -53 | -68 |




Market 17.11 Alternative 2B, 60% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 69 | 78 | 63 | 11 | 10 | 36 | 74 | 133 | 164 | 231 |
| P75 | 48 | 22 | -1 | -34 | -41 | -38 | -16 | 25 | 43 | 58 |
| P50 | 22 | -17 | -42 | -63 | -70 | -64 | -57 | -32 | -18 | -18 |
| P25 | -13 | -42 | -64 | -82 | -86 | -85 | -82 | -78 | -70 | -73 |
| P5 | -45 | -76 | -85 | -84 | -86 | -95 | -95 | -96 | -94 | -95 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 220 | 136 | 248 | 323 | 455 | 480 | 550 | 636 | 694 | 792 |
| P75 | 169 | 82 | 78 | 125 | 226 | 251 | 278 | 359 | 387 | 415 |
| P50 | 123 | 13 | 12 | 42 | 111 | 122 | 152 | 218 | 245 | 254 |
| P25 | 15 | -44 | -34 | -24 | -11 | -1 | 19 | 55 | 88 | 72 |
| P5 | -45 | -71 | -68 | -71 | -69 | -78 | -81 | -69 | -84 | -90 |




Spat

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 1564 | 1987 | 2546 | 2560 | 2549 | 2995 | 3176 | 3489 | 3408 | 2880 |
| P75 | 680 | 925 | 1666 | 1666 | 1635 | 1884 | 2084 | 2361 | 1948 | 1609 |
| P50 | 574 | 738 | 1247 | 1179 | 1134 | 1362 | 1532 | 1787 | 1417 | 1072 |
| P25 | 408 | 547 | 866 | 800 | 710 | 936 | 1081 | 1227 | 995 | 645 |
| P5 | 88 | 114 | 78 | 41 | -2 | -13 | -17 | -58 | -68 | -68 |




Market 17.12 Alternative 2B, 80% harvest

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 49 | 58 | 44 | -6 | -9 | 11 | 50 | 120 | 147 | 197 |
| P75 | 29 | 12 | -12 | -43 | -50 | -42 | -25 | 12 | 28 | 55 |
| P50 | 11 | -24 | -47 | -68 | -73 | -68 | -62 | -42 | -27 | -16 |
| P25 | -23 | -48 | -66 | -82 | -86 | -85 | -84 | -79 | -69 | -77 |
| P5 | -49 | -73 | -86 | -94 | -95 | -96 | -96 | -96 | -95 | -97 |



Small

| Percentile | Year | | | | | | | | | |
|------------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 217 | 131 | 190 | 245 | 401 | 436 | 495 | 574 | 630 | 725 |
| P75 | 167 | 83 | 65 | 101 | 197 | 225 | 272 | 334 | 371 | 408 |
| P50 | 126 | 12 | 10 | 31 | 93 | 112 | 139 | 200 | 223 | 244 |
| P25 | 31 | -42 | -34 | -33 | -12 | 3 | 10 | 54 | 74 | 45 |
| P5 | -44 | -71 | -71 | -73 | -71 | -77 | -83 | -87 | -86 | -91 |



Spat

| Percentile | Year | | | | | | | | | |
|------------|------|------|------|------|------|------|------|------|------|------|
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
| P95 | 1451 | 1633 | 2377 | 2240 | 2264 | 2610 | 3011 | 3229 | 3184 | 2642 |
| P75 | 644 | 836 | 1504 | 1513 | 1553 | 1797 | 2043 | 2264 | 1904 | 1513 |
| P50 | 544 | 688 | 1158 | 1074 | 1028 | 1305 | 1483 | 1680 | 1404 | 1039 |
| P25 | 382 | 500 | 894 | 784 | 674 | 965 | 1072 | 1240 | 1005 | 629 |
| P5 | 69 | 63 | 50 | 26 | -4 | -1 | -40 | -57 | -48 | -75 |





Figure 17 . High Salinity (continued)



12. Appendices

Appendix A. Development of Habitat Layer (Greenhawk 2005)

Development of a

Potential Habitat Layer for Chesapeake Bay Oyster Bottom

Prepared by Kelly Greenhawk
Maryland Department of Natural Resources, Fisheries Division

December 2005

The dataset described in this document was created for use by scientists, managers, and modelers involved in work related to the preparation of an environmental impact statement entitled "Development of an Environmental Impact Statement for Introducing Non-Native Oyster Species into the Chesapeake Bay, Including an Evaluation of Native Oyster Restoration Alternatives". The dataset (filename "Chesapeake Bay habitat v1.shp") is in the ESRI shapefile format, and is a spatial representation of the locations and extent of oyster habitat (including leased bottom) for both the Maryland and Virginia portions of the Chesapeake Bay. The file will be used as input for modeling efforts related to the EIS referenced above, and in the generation of spatial products related to this project such as report figures. The creation of this file was necessitated by the lack of a recent comprehensive oyster bar survey in either Maryland or Virginia and an awareness that significant oyster habitat loss has occurred in recent years. The approach in developing these data sets involved adjusting the area and the habitat quality of oyster bars throughout Chesapeake Bay as determined in historical surveys to account for this loss of habitat. For Maryland, the area and habitat quality of each oyster bar determined in a comprehensive field survey carried out in the mid 1970s through early 1980s was adjusted by the results of a 1999 – 2000 survey of the size and habitat quality of a small subset of these bars. In Virginia, adjustments to a comprehensive historical oyster bar survey were made based on recent experience in the field. The dataset described in this document was developed as input to model the distribution and abundance of oysters in Chesapeake Bay. While this data set will provide the resolution needed to model the oyster population, a re-survey of oyster habitat in Maryland and Virginia is needed prior to initiation of any site specific or regional management activities requiring accurate delineation of Chesapeake Bay oyster bars.

Created using ESRI's ArcGIS product, the file was derived from the following spatial data layers created and maintained by the Maryland Department of Natural Resources, the Virginia Institute of Marine Sciences, and the Virginia Marine Resources Commission:

Maryland datasets:

- ESRI shapefile which delineates results from the Maryland Bay Bottom Survey, an acoustic and patent tong survey conducted by the Department from 1978 to 1983 (survey will be referred to as the "MBBS" onward in this document). The results of the survey categorize Maryland's Bay bottom into seven classifications: cultch, mud with cultch, sand with cultch, mud, sand, hard bottom and leased bottom. Based upon review of recent acoustic survey work and the collective knowledge of the Department's Shellfish Program, it was determined that areas classified as mixed cultch by the MBBS ("mud with cultch" and "sand with cultch") had likely degraded to non-cultch bottom. For this reason, only areas classified as "cultch" bottom type have been included in this data product. (file name "BBSurvey.shp")
- ESRI shapefile which delineates oyster bar boundaries charted by C.C. Yates 1906-1911. Yates boundaries were only used in limited sections of the Eastern Bay, Broad Creek and Harris Creek, where data from the Maryland BBS is missing. (file name "Yatesbrs.shp")
- ESRI shapefile which delineates oyster repletion activities undertaken by the Department's Shellfish Program (file name "DNRRepletion.shp")
- ESRI shapefile which delineates oyster repletion activities undertaken by the Department and other groups within Maryland's oyster sanctuaries and reserves (file name "MdSancRes.shp")
- Dxf file prepared by the Natural Resources Police containing linear boundaries for current Maryland oyster leases (file name "Lselines.dxf")
- Dbase file prepared by the Natural Resources Police containing corner coordinates for current (2004) Maryland oyster leases (file name "Lease91.dbf")

Virginia datasets:

- ESRI shapefile which delineates potential oyster restoration sites the Virginia portion of the Chesapeake Bay. (filename "Potential.shp")
- ESRI shapefile representation of Virginia's lease boundaries (based on 2002 lease boundaries) prepared by staff at VIMS (filename "Privlease.shp")

Metadata for the source datasets can be found in Appendix 1.

Detailed Process Step:

Data was processed using a combination of the following software packages: ArcGIS software versions 9.0 and 9.1, XTools Pro version 2.0.0, MapInfo Professional versions 7.0 and 7.5, MapBasic version 7.5, Excel 2002, and Multi-Edit version 7.11. Footnotes throughout this document indicate the specific software package used for a particular process step.

Development of the Maryland habitat

- 1.) Polygons not classified as "cultch" in the Maryland Bay Bottom Survey dataset were removed from the MBBS shapefile.¹
- 2.) Because chart # 9 was never located when the MBBS was digitized, data is missing from the source file for parts of the Eastern Bay, Harris Creek and Broad Creek. Consequently, oyster bar boundaries from the Yates survey were used for habitat definitions in these areas. A total of 59 Yates bar boundaries were appended to the file.¹
- 3.) During the MBBS, areas of Rangia clam shells in the Magothy River were mapped as cultch. Based on the collective knowledge of and recent observations by Shellfish Program biologists, *today* that area is mud bottom. For this reason, "cultch" polygons in this tributary (totaling 1,963 acres) were removed from the file¹
- 4.) Based on longevity studies of shell plantings conducted using underwater video and acoustic survey gear, and consultation with Shellfish Program staff, a conservative estimate of planting longevity is five years. The five year time span is an average and was used in order to provide the best estimate of habitat for the file. Polygons representing oyster repletion efforts undertaken by Maryland DNR and other entities between 1999 and 2003 were appended to the file. (files "DNRRepletion.shp" and "MdSancRes.shp") Plantings older than 5 years were not used.¹
- 5.) In order to calculate acreages for all the polygons, the file was projected to US State Plane projection and acreages were calculated. The file was then un-projected to a geographic coordinate system (WGS84).^{1 2}
- 6.) File was translated to MapInfo for further processing.³
- 7.) Overlapping planting polygons were combined in MapInfo to ensure that habitat (hence acreages) were not duplicated³. Figure 1 show how a pair of overlapping planting polygons looked before and after the combine operation.




Figure 1. The two plantings in the figure on the left (hatched objects), were combined to create one polygon, shown on the right.

8.) Due to the contiguous nature of the bars charted by Yates, those that shared a boundary were treated as one large oyster bar, and adjacent bars were combined before being reduced. To facilitate this task, MapInfo's "Combine" operation was applied.³ Figure 2 illustrates how one cluster of Yates bars looked before and after the combine operation.



Figure 2. Five polygons in the map on the left were combined to create one large polygon shown on the right (dark outline).

9.) MapInfo file was translated back to a shapefile for continued work in ArcMap.³

10.) Concurrent with well-documented evidence that the Bay's oyster habitat has degraded over time, it was determined that adjustments to the dataset were necessary to make the layer more accurately reflect the state of today's oyster habitat. In order to account for the age of the data (The MBBS was conducted between 1978 and 1984.), reduction factors for habitat defined by the MBBS and the Yates survey were identified, based on the results of acoustic surveys performed between 1999 and 2003 by MDNR.⁷ Appendix 2 shows the summary table from which the reduction factors were taken. The summary table was extracted from the paper "Assessment of Recent Habitat Conditions of Eastern Oyster *Crassostrea virginica* Bars in Mesohaline Chesapeake Bay" (Smith et al., North American Journal of Fisheries Management 25:1569-1590, 2005). Appendix 3 contains underwater images from visual ground truth operations for the various bottom classifications used in Appendix 2. The reduction factors used were as follows:

- MBBS cultch polygons - reduction factor equal to 29.17%

This adjustment factor will result in a polygon that is 29.17% of its original area after the scale operation is applied. This value was obtained by summing the frequencies for the four acoustic survey categories "heavily sedimented shell with mud", "heavily sedimented shell with sand", "lightly sedimented shell with sand", and "clean shell" in the column labeled "Inside MBBS Bottom" ($21.64 + 5.16 + 1.21 + 1.16 = 29.17\%$).

- Yates polygons - reduction factor equal to 12.24%

This value was obtained by summing the frequencies for the same four acoustic survey categories in the column labeled "Inside Yates Bars". (8.33 + 2.96 + .33 + .62 = 12.24%)

- Plantings deployed within the last 5 years (1999 through 2003) were not reduced.

11.) Scale factors were calculated for a.) the Maryland BBS polygons and b.) the Yates polygons, and Visual Basic code obtained from ESRI was used to "scale" or shrink the polygons.¹ The formula used to calculate the scale factor is

$$S = \sqrt{(a_d / a_o)}$$

whereby S = scale factor, a_d = desired area, and a_o = original area.

The program used to perform this task is detailed in Appendix 4. Figure 3 illustrates the result of the scale operation on a cluster of habitat polygons.




Figure 3. Figure showing the effect of the scale operation. Scaled polygons (dark gray) are shown overlaying the original (MBBS) habitat polygons (light gray). Polygons resulting from the scale operation (dark gray) in this step represent areas of low *and* high quality habitat, specifically as follows:

Maryland BBS polygons -----> 26.8% of resulting polygons represent "low quality" cultch (21.64% + 5.16% from Appendix 2) and
2.37% of resulting polygons represent "high quality" cultch (1.21% + 1.16% from Appendix 2).

Therefore, it follows that 91.9 % of the MBBS polygons resulting from the scale operation can be designated low quality cultch

calculated $.268 / .2917 = .918753 = .919 = 91.9 \%$ and

8.1% of the MBBS polygons resulting from the scale operation can be designated high quality cultch

calculated $.0237 / .2917 = .081248 = .081 = 8.1 \%$.

Yates polygons -----> 11.29% of resulting polygons represent "low quality" cultch (8.33% + 2.96% from Appendix 2) and
.95% of resulting polygons represent "high quality" cultch (.33% + .62% from Appendix 2).

Therefore, 92.2 % of the Yates polygons resulting from the scale

operation can be designated low quality cultch

calculated $.1129 / .1224 = .922386 = .922 = 92.2\%$ and

7.8 % of the Yates polygons resulting from the scale operation can be designated high quality cultch

calculated $.0095 / .1224 = .077615 = .078 = 7.8\%$.

The concept of "low quality" and "high quality" cultch was derived from work performed under the interstate research project entitled "Oyster Population Estimation in Support of the Ten-year Goal for Oyster Restoration in the Chesapeake Bay Fishery". More information about the oyster population estimation work can be found on the project's website. (<http://www.vims.edu/mollusc/cbope/overview.htm>)

12.) New acreage values were calculated for the dataset, in order to confirm the reduced areas.¹

13.) File was translated back to a MapInfo format for further processing.³

14.) Overlapping polygons of different types were addressed as follows:

- In areas where planting polygons (considered "high quality" habitat) overlapped polygons from the MBBS or Yates survey, the area of overlap was "erased" from the low quality habitat to ensure that habitat was not duplicated.³
- There were no areas of overlap involving Maryland BBS polygons and Yates boundaries, since the Yates boundaries were only used in the area encompassed by chart number 9 of the MBBS.

Figure 4 shows overlapping habitat polygons before and after the erase tool was used.



Figure 4. The image on the left shows a planting (gray hatched polygon) that overlaps a Maryland BBS cultch polygon (gray). The image in the center shows the cultch polygon after the erase operation. (For illustration purposes the planting polygon has been omitted.) The right-most image shows both polygons after the erase operation, with the cultch polygon surrounding three sides of the planting, but not overlapping the planting.

15.) Dataset was mapped with layers representing NOAA's inshore water body areas and basins from the Chesapeake Bay Oyster Population Estimation (CBOPE) project (Harding 2005). Fields were added to accommodate region codes and basin names, then updated with these values.³

16.) Dataset was plotted for QA/QC operations.³

Summary statistics for Maryland habitat records / polygons:

- 2,115 polygons
 - 1949 polygons were derived from the Maryland Bay Bottom Survey
 - 123 polygons were derived from shell plantings
 - 43 polygons were derived from the Yates bars dataset
- Acres of Maryland habitat, not including leased bottom sum of field "A" = 36,144.49 (sum of field "A" acres of Maryland habitat, not including leased bottom)

Development of the Maryland leased bottom

Maryland's official lease delineations are maintained by DNR's Natural Resources Police in Autocad format. For this reason, considerable effort was required to transform the lease lines into a GIS-compatible format. The process of translating the data to a shapefile format began with the acquisition of the following datasets from NRP:

- Iselines.dxf - linear delineation of Maryland's oyster leases, stored in Autocad format
- lease91.dbf - dBase file; records for 780 locations; each record representing the location of the first corner of each lease

In addition, Maryland lease charts (prepared by MDNR, Natural Resources Police, dated 7-17-2001) were used for QA/QC operations. (Leases located in Maryland's coastal bays are not included in this file.)

- 1.) Coordinate information in the Dbase file was used to create a shapefile in ArcGIS 9. (point feature type)¹
- 2.) Dxf file was imported into TNT MIPS software by Kevin Boone (MDNR, Watershed Services Division), where various processes were applied in order to convert lines in the DXF file to a polygon shapefile.
- 3.) The 2 files resulting from the above processes were converted to MapInfo format then overlaid in MapInfo for verification and editing purposes.³
- 4.) Polygons present in the file generated by MIPS without corresponding point objects in the file of corner coordinates were deleted since they could not be verified by the NRP Dbase file. (per instructions from Williams, NRP)³
- 5.) Polygons that overlapped the shoreline were shaped to the shoreline.³
- 6.) Once all edits were applied, a series of charts were printed on transparent film and overlaid onto copies of the official charts, obtained from NRP. The leases on each film were compared to lease outlines on corresponding charts to verify lease shape, size, and relationship to the shoreline.³
- 7.) A field was added to hold a calculated acreage value for each lease. Acreages were calculated in MapInfo.³

It should be stressed that this dataset was created solely for use on this project, and should not be used for any other purposes. For legal definitions of Maryland's leased bottom, contact the Natural Resources Police.

Summary statistics for Maryland leased bottom records / polygons:

- 767 lease polygons (File does not contain leased bottom for Maryland's coastal bays)
- Total lease acreage = 7,747.81 (sum of field "A")

Although the Maryland lease boundaries are included in this habitat layer, because leased bottom is considered potential habitat only for aquaculture purposes, the Maryland leases were not used in subsequent modeling operations. (Only potential habitat currently available for larval settlement was used in the modeling scenarios.)

Development of the Virginia habitat

- 1.) The source dataset for non-lease habitat in the Virginia portion of the Bay is the shapefile "Potential.shp", which is a collection of polygons representing suitable oyster restoration sites. Obtained from the Virginia Institute of Marine Sciences, the file contains 2,711 polygon regions, totaling 12,290.32 acres of potential oyster habitat. The file was generated by staff at VIMS and VMRC and complements the "Virginia Oyster Reef Restoration Map Atlas", published by VIMS in 2002.
- 2.) Habitat polygons in the coastal bays were removed from the file. (1,063 polygons removed) ¹
- 3.) Polygons in the file without attribute data were determined to represent holes in oyster habitat (personal communication with VIMS staff). A hole is a hole in a habitat polygon; an area of non-oyster habitat that is completely surrounded by habitat. These polygons were removed from the file and saved as a separate layer, for erase operations. ¹
- 4.) Habitat polygons were erased using the holes layer. ³
- 5.) A series of maps were plotted similar in scale to those found in the map atlas and each map was compared to its corresponding map in the map atlas. ¹
- 6.) Overlapping polygons were merged into one polygon using the dissolve operation. ¹
- 7.) Using the file of NOAA water body regions, each habitat polygon was assigned a region code. ¹
- 8.) Using the CBOPE basin layer, each polygon was assigned a basin name. Pursuant to instructions from Versar modelers, Virginia polygons which fell outside of the CBOPE basins (Harding 2005) were assigned to the closest basin. ¹

Summary statistics for Virginia habitat records / polygons:

- 769 polygons
- total acreage = 11,355.13 acres (sum of field "A")

Development of the Virginia leased bottom

The source file for the Virginia leases ("Privlease.shp") is a polygon delineation of leased shellfish bottom prepared by VIMS and based on VMRC lease records.

- 1.) Shapefile converted to MapInfo format for processing in MapInfo. ³
- 2.) Lease polygons in the Virginia coastal bays were removed from the file, leaving 5,142 records. ³
- 3.) Each of 48 multi-polygon lease polygons in the file was located, and nested polygons were combined. This operation was necessary to prepare for the scale operation to follow. ³
- 4.) File was translated to a shapefile for further processing in Arcmap ³
- 5.) A scale factor was calculated for the reduction process as follows:

$$S = \sqrt{(a_d / a_o)}$$

whereby S = scale factor, a_d = desired area, and a_o = original area.

- 6.) Visual Basic code (Appendix 4) was used to "scale" or shrink the polygons to 30% of their original area. This 30% value was provided by scientific staff at VMRC, based on recent field experience and is considered the best available estimate of current habitat. ¹

7.) Results of scale operation were again visually inspected for accuracy. During this inspection, it was discovered that nested leases (a lease sharing 2 or more sides with another lease) did not shift in tangent with outer leases. An example of the resulting inconsistencies is shown in the following figures.



Figure 5. Nested lease polygons before (left) and after (right) the scale operation.

- 8.) File was translated file to Mapinfo for combine operations.³
- 9.) Nested polygons were located and the combine operation was performed.³
- 10.) File was translated back to Arcview format.³
- 11.) In order to confirm reduced areas, the file was projected to US State Plane projection (Maryland, NAD 83, meters), and acreage values were calculated for the dataset.^{1,2}
- 12.) Leases smaller than 0.1 acre were identified, and if determined to be sliver polygons, were deleted.¹
- 13.) File was un-projected back to a geographic coordinate system (WGS84).¹
- 14.) Dataset was again visually inspected for QA/QC purposes.¹ Attempts to obtain hardcopy charts of Virginia's oyster lease boundaries from VMRC for QA/QC purposes failed.

Summary statistics for Virginia leased bottom records / polygons:

- 4,829 polygons
- total acreage = 20,782.80 acres (sum of field "A")

Compilation of master habitat file

- 1.) A master habitat file was created by merging the 4 datasets described above into one shapefile.¹
- 2.) MapInfo split tool was applied to polygons which fell on the border of two basins.³
- 3.) MapInfo "check regions" feature was used to detect self-intersections, overlaps and gaps. This operation returned a table of 269 problem areas.³
- 4.) Intersections, overlaps and gaps were corrected in the file using the following rules:³

The overlap between a Maryland habitat polygon and a Maryland lease polygon was corrected by allowing the habitat polygon to take precedence, so the intersection was erased from the lease.
- 5.) Fields were added to the master file to accommodate lease ids and EIS alternative numbers, then updated accordingly.

A total of 767 records (all Maryland leases) contain lease ids and 60 records contain data in the alternative field.¹

¹ ArcGIS/ArcMap version 9.0

² XTools Pro version 2.0.0

³ MapInfo Professional versions 7.0 and 7.5

⁴ MapBasic version 7.5

⁵ Excel 2002

⁶ Multi-Edit version 7.11

⁷ Acoustic surveys were conducted by MDNR using a seabed classification system produced by Questar Tangent. See <http://marine.questertangent.com> for more information on the system used.

Attribute names, field types and definitions:

FID - ESRI internal feature identifier

Shape - (type of object); ESRI defined value (Value is "Polygon" in this file.)

ID - integer (9) unique numeric identifier; possible values 1 through 8,480

CX - Double (12) longitude value of the centroid, expressed in decimal degrees

CY - Double (12) latitude value of the centroid, expressed in decimal degrees

A - Double (19) area of habitat polygon, expressed in acres

B - String (1) Basin name from Chesapeake Bay Oyster Population Estimation project:

Maryland basins:

CHESTER
CHOPTANK
EASTERN BAY
LITTLE CHOPTANK
MD MAINSTEM
MD POTOMAC
PATUXENT
TANGIER

Virginia basins:

EASTERN SHORE / TANGIER
GREAT / LITTLE WICOMICO
JAMES
PIANKATANK
POQUOSON / BACK
RAPPAHANNOCK
VA MAINSTEM
VA POTOMAC
YORK / MOBJACK

See <http://www.vims.edu/mollusc/cbope/index.htm> for details on the above basin designations.

N - String (3) numeric NOAA water body code; possible values are as follows:

| | | |
|-----|-----|-----|
| 212 | 243 | 271 |
| 217 | 245 | 273 |
| 218 | 246 | 276 |
| 220 | 248 | 278 |
| 224 | 249 | 279 |
| 225 | 251 | 301 |
| 226 | 252 | 306 |
| 228 | 253 | 307 |
| 230 | 254 | 308 |
| 231 | 257 | 309 |
| 232 | 258 | 311 |
| 235 | 259 | 313 |
| 236 | 267 | 314 |
| 237 | 268 | 315 |
| 239 | 270 | 316 |

| | | |
|-----|-----|-----|
| 317 | 339 | 367 |
| 321 | 343 | 368 |
| 322 | 345 | 369 |
| 324 | 346 | 371 |
| 327 | 347 | 372 |
| 328 | 351 | 374 |
| 329 | 353 | 375 |
| 332 | 354 | 379 |
| 333 | 355 | 380 |
| 335 | 358 | 381 |
| 336 | 363 | 382 |
| 337 | 364 | 391 |
| 338 | 366 | 392 |

The shapefile used to assign the NOAA codes was obtained from the Chesapeake Bay Program office and is based on NOAA's water body codes, also known as "NEM" areas. "NEM" is an abbreviation for NEMFIS, which was the Northeast Marine Fisheries Information System, a State and Federal initiative to work together in collecting and managing commercial fisheries data.

S - String (2) source code for origin of habitat polygon; possible values are as follows:

MH = Maryland habitat
ML = Maryland lease
VH = Virginia habitat
VL = Virginia lease

HT - String (1) alphabetic code representing the habitat type for Maryland habitat polygons; possible values are as follows:

B = Maryland Bay Bottom Survey
P = planting
Y = Yates bar

Only Maryland habitat polygons will have a value in the HT field.

LID - String (15) lease identifier; 3 digit lease identifier (assigned by MDNR, NRP), followed by a dash, followed by an acreage value. LID field will be blank for Virginia leases.

ALT - String (2) EIS alternative number for which lease will be used; possible values are as follows:

4 - EIS alternative number 4
5 - EIS alternative number 5
45 - EIS alternative numbers 4 and 5

Sixty Maryland leases have a value in the ALT field. The remaining records are blank.

CLOS_STAT - String (50) closure status; possible values for Maryland records are "SANCTUARY", "RESTRICTED" or blank; possible values for Virginia records are "CONDEMNED", "PROHIBITED", "PROHIBITED NON-PRODUCTIVE", "CONDEMNED AND PROHIBITED" or blank.

Note: Only polygons with greater than 50% area inside a closure were flagged using this field.

CLOS_NAME - String (50) name of closure assigned by either the Maryland Department of the Environment or the Virginia Department of Health

Summary statistics for file Chesapeake Bay Habitat v1.shp:

- shapefile contains 8,480 polygons / records
 - 2,115 polygons define Maryland habitat
 - 767 polygons define Maryland leases
 - 769 polygons represent Virginia habitat
 - 4,829 polygons represent Virginia leases

- sum of acreage = 76,030.23, specifically as follows:
 - 36,144.49 acres of Maryland habitat
 - 7,747.81 acres of Maryland leases
 - 11,355.13 acres of Virginia habitat
 - 20,782.80 acres of Virginia leases

References

Harding, J.. (2005). Chesapeake Bay Oyster Population Estimation (CBOPE). Available: <http://www.vims.edu/mollusc/cbope/overview.htm>. Last accessed April 5, 2007.

Appendix 1.

Metadata for source datasets.

"BBSurvey.shp"

Identification_Information:

Citation:

Citation_Information:

Originator: Maryland Department of Natural Resources

Publication_Date: 20030204

Publication_Time: Unknown

Title: **BBSURVEY**

Geospatial_Data_Presentation_Form: vector digital data

Series_Information:

Publication_Information:

Online_Linkage: <http://dnrweb.dnr.state.md.us/gis/data/index.html>

Description:

Abstract: Polygon dataset characterizing bottom type designations determined by MDNR's Acoustic Bay Bottom Survey conducted from 1974 to 1983. Bottom type designations include cultch, mud, sand, leased bottom, hard bottom, mud with cultch and sand with cultch. Note: The data in this file is up to 30 years old and areas designated as "cultch bottom" when this survey was conducted have likely degraded. For this reason, it is very likely that many of the areas shown as cultch in the dataset are no longer valid. The data in this file should only be used as a guide to the location of oyster bars in the mid to late 1970's and early 1980's.

Purpose:

This file was created for Maryland DNR planning purposes, specifically for the purpose of managing Maryland's oyster resource. This file is the result of the digitization of the 37 mylars of the Md Bay Bottom Survey.

Supplemental_Information: Digitization was performed using MapInfo software (version 3x) between May of 1994 and May of 1996 at the Oxford Laboratory. All mylars were digitized with the exception of mylar # 9, which could not be located. Mylar # 9 covers the southern portion of the Eastern Bay.

Time_Period_of_Content:

Time_Period_Information:

Range_of_Dates/Times:

Beginning_Date: 1974

Beginning_Time: unknown

Ending_Date: 1983

Ending_Time: unknown

Currentness_Reference: ground condition

Status:

Progress: Complete

Maintenance_and_Update_Frequency: None planned

Spatial_Domain:

Bounding_Coordinates:

West_Bounding_Coordinate: -77.049563

East_Bounding_Coordinate: -75.669141

North_Bounding_Coordinate: 39.312854

South_Bounding_Coordinate: 37.903566

Keywords:

Theme:

Theme_Keyword_Thesaurus: none

Theme_Keyword: sediment

Theme_Keyword: bars

Theme_Keyword: mud

Theme_Keyword: sand

Theme_Keyword: cultch

Theme_Keyword: leased
 Theme_Keyword: hard bottom
 Place:
 Place_Keyword_Thesaurus: none
 Place_Keyword: Maryland
 Stratum:
 Stratum_Keyword_Thesaurus: none
 Stratum_Keyword: Chesapeake Bay seafloor
 Temporal:
 Temporal_Keyword_Thesaurus: none
 Temporal_Keyword: 1970's
 Access_Constraints: none
 Use_Constraints: The Department of Natural Resources makes no warranty, expressed or implied, as to the use or appropriateness of Spatial Data, and there are no warranties of merchantability or fitness for a particular purpose or use. The information contained in Spatial Data is from publicly available sources, but no representation is made as to the accuracy or completeness of Spatial Data. The Department of Natural Resources shall not be subject to liability for human error, error due to software conversion, defect, or failure of machines, or any material used in the connection with the machines, including tapes, disks, CD-ROM's or DVD-ROM's and energy. The Department of Natural Resources shall not be liable for any lost profits, consequential damages, or claims against the Department of Natural Resources by third parties. The liability of the Department of Natural Resources for damage regardless of the form of the action shall not exceed any distribution fees that may have been paid in obtaining Spatial Data.

Point_of_Contact:
 Contact_Information:
 Contact_Person_Primary:
 Contact_Person: Kelly Greenhawk
 Contact_Organization: Maryland Department of Natural Resources
 Contact_Position: DP Programmer / Analyst
 Contact_Address:
 Address_Type: mailing address
 Address: 904 South Morris Street
 City: Oxford
 State_or_Province: Maryland
 Postal_Code: 21654
 Country: USA
 Contact_Voice_Telephone: 410-226-0078
 Contact_TDD/TTY_Telephone: none
 Contact_Facsimile_Telephone: 410-226-0120
 Contact_Electronic_Mail_Address: kgreenhawk@dnr.state.md.us
 Hours_of_Service: Monday through Friday 0830 - 1700 (EST)

Data_Set_Credit: Maryland Department of Natural Resources, Sarbanes Cooperative Oxford Laboratory
 Security_Information:
 Security_Classification_System: none
 Security_Classification: Unclassified
 Security_Handling_Description: none
 Native_Data_Set_Environment: Microsoft Windows XP Version 5.1 (Build 2600) Service Pack 2; ESRI ArcCatalog 9.1.0.722

Data_Quality_Information:
 Attribute_Accuracy:
 Attribute_Accuracy_Report: Dataset was digitized from mylars of the survey which were generated by the Department's Hydrographic Operations Division. Scale of mylars is 1:20,000.
 Positional_Accuracy:
 Horizontal_Positional_Accuracy:
 Horizontal_Positional_Accuracy_Report: *****
 Vertical_Positional_Accuracy:
 Vertical_Positional_Accuracy_Report: *****
 Lineage:

Process_Step:

Process_Description: Before digitization, mylars were registered in US State Plane 1927 feet. Original MapInfo file created took on the MapInfo proprietary coordinate system and projection, Lat / Long with a proprietary datum comparable to NAD 83.

Process_Date: Unknown

Process_Contact:

Contact_Information:

Contact_Person_Primary:

Contact_Person: Kelly Greenhawk

Contact_Organization: Maryland Department of Natural Resources

Contact_Position: DP Programmer Analyst

Contact_Voice_Telephone: 410-226-0078

Contact_TDD/TTY_Telephone: none

Contact_Facsimile_Telephone: 410-226-0120

Contact_Electronic_Mail_Address: kgreenhawk@dnr.state.md.us

Hours_of_Service: Monday through Friday 0800 - 1630 EST

Process_Step:

Process_Description: Metadata imported.

Source_Used_Citation_Abbreviation: C:\arcgis\arcexe82\Metadata\Stylesheets\DNRTemplate.xml

Process_Step:

Process_Description: Metadata imported.

Source_Used_Citation_Abbreviation: Z:\All\SCOLdata\Outgoing\Shapefiles\Mdoysbrs.shp.xml

Process_Step:

Process_Description: Metadata imported.

Source_Used_Citation_Abbreviation: Z:\All\SCOLdata\Metadata\BBSURVEY.shp.xml

Process_Step:

Process_Description: Dataset copied.

Source_Used_Citation_Abbreviation:

Spatial_Data_Organization_Information:

Direct_Spatial_Reference_Method: Vector

Point_and_Vector_Object_Information:

SDTS_Terms_Description:

SDTS_Point_and_Vector_Object_Type: G-polygon

Point_and_Vector_Object_Count: 7556

Spatial_Reference_Information:

Horizontal_Coordinate_System_Definition:

Planar:

Grid_Coordinate_System:

Grid_Coordinate_System_Name: State Plane Coordinate System 1983

State_Plane_Coordinate_System:

SPCS_Zone_Identifier: 1900

Lambert_Conformal_Conic:

Standard_Parallel: 38.300000

Standard_Parallel: 39.450000

Longitude_of_Central_Meridian: -77.000000

Latitude_of_Projection_Origin: 37.666667

False_Easting: 400000.000000

False_Northing: 0.000000

Planar_Coordinate_Information:

Planar_Coordinate_Encoding_Method: coordinate pair

Coordinate_Representation:

Abcissa_Resolution: 0.000256

Ordinate_Resolution: 0.000256

Planar_Distance_Units: meters

Geodetic_Model:

Horizontal_Datum_Name: North American Datum of 1983

Ellipsoid_Name: Geodetic Reference System 80
 Semi-major_Axis: 6378137.000000
 Denominator_of_Flattening_Ratio: 298.257222
 Entity_and_Attribute_Information:
 Detailed_Description:
 Entity_Type:
 Entity_Type_Label: BBSURVEY
 Attribute:
 Attribute_Label: FID
 Attribute_Definition: Internal feature number.
 Attribute_Definition_Source: ESRI
 Attribute_Domain_Values:
 Unrepresentable_Domain: Sequential unique whole numbers that are automatically generated.
 Attribute:
 Attribute_Label: Shape
 Attribute_Definition: Feature geometry.
 Attribute_Definition_Source: ESRI
 Attribute_Domain_Values:
 Unrepresentable_Domain: Coordinates defining the features.
 Attribute:
 Attribute_Label: BOT_TYPE
 Attribute_Definition: Numeric bottom type code, 1 through 7 as follows: 1 = mud; 2 = sand; 3 = sand with cultch; 4 = mud with cultch; 5 = cultch; 6 = hard bottom; 7 = leased bottom (leased bottom was not surveyed, only charted.)
 Attribute_Definition_Source: Maryland Department of Natural Resources
 Attribute:
 Attribute_Label: BOTTOM
 Attribute_Definition: Textual description of bottom type as follows: "MUD", "SAND", "MUD WITH CULTCH", "SAND WITH CULTCH", "CULTCH", "HARD BOTTOM", AND "LEASED"
 Attribute_Definition_Source: Maryland Department of Natural Resources
 Attribute:
 Attribute_Label: BORDER
 Attribute_Definition: Field contains "I" for "incomplete" if any part of the polygon's border overlaps the mylar's border.
 Attribute_Definition_Source: Maryland Department of Natural Resources
 Attribute:
 Attribute_Label: HOLES
 Attribute_Definition: Number of "holes" in the polygon object. Holes are either caused by physical features such as islands, where the surveyors could not work, or other bottom types.
 Attribute_Definition_Source: Maryland Department of Natural Resources
 Attribute:
 Attribute_Label: Acres
 Overview_Description:
 Entity_and_Attribute_Detail_Citation: *****
 Distribution_Information:
 Resource_Description: Downloadable Data
 Standard_Order_Process:
 Digital_Form:
 Digital_Transfer_Information:
 Transfer_Size: 3.140
 Distribution_Information:
 Distributor:
 Contact_Information:
 Contact_Person_Primary:
 Contact_Person: Kelly Greenhawk
 Contact_Organization: Maryland Department of Natural Resources
 Contact_Position: DP Programmer / Analyst
 Contact_Address:

Address_Type: mailing and physical address
Address: Sarbanes Cooperative Oxford Laboratory
Address: 904 South Morris Street
City: Oxford
State_or_Province: Maryland
Postal_Code: 21654
Country: USA
Contact_Voice_Telephone: 410-226-0078
Contact_TDD/TTY_Telephone: none
Contact_Facsimile_Telephone: 410-226-0120
Contact_Electronic_Mail_Address: kgreenhawk@dnr.state.md.us
Hours_of_Service: Monday through Friday 0800 - 1700

Distribution_Liability: This data represents the results of data generated for a specific Maryland Department of Natural Resources management activity. As such, they are only valid for their intended use, content, time, and accuracy specifications. The user is responsible for any application of the data for other than it's intended purpose.

Standard_Order_Process:

Digital_Form:

Digital_Transfer_Information:

Format_Name: ARCE shapefile

Format_Information_Content: Data is normally distributed as a statewide dataset - Md State Plane NAD 83 meters

File-Decompression_Technique: no compression applied

Transfer_Size: 3.140

Digital_Transfer_Option:

Online_Option:

Online_Computer_and_Operating_System: <http://dnrweb.dnr.state.md.us/gis/data/index.html>

Metadata_Reference_Information:

Metadata_Date: 20050629

Metadata_Contact:

Contact_Information:

Contact_Person_Primary:

Contact_Person: Kelly Greenhawk

Contact_Organization: Maryland Department of Natural Resources

Contact_Position: DP Programmer / Analyst

Contact_Address:

Address_Type: mailing and physical address

Address: Sarbanes Cooperative Oxford Laboratory

Address: 904 South Morris Street

City: Oxford

State_or_Province: Maryland

Postal_Code: 21654

Country: USA

Contact_Voice_Telephone: 410-226-0078

Contact_TDD/TTY_Telephone: none

Contact_Facsimile_Telephone: 410-226-0120

Contact_Electronic_Mail_Address: kgreenhawk@dnr.state.md.us

Hours_of_Service: Monday through Friday 0830 - 1700 (EST)

Metadata_Standard_Name: FGDC Content Standards for Digital Geospatial Metadata

Metadata_Standard_Version: FGDC-STD-001-1998

Metadata_Time_Convention: local time

Metadata_Security_Information:

Metadata_Security_Classification: Unclassified

Metadata_Extensions:

Online_Linkage: <http://www.esri.com/metadata/esriprof80.html>

Profile_Name: ESRI Metadata Profile

"Yatesbrs.shp"

Identification_Information:

Citation:

Citation_Information:

Originator: Maryland Department of Natural Resources, Fisheries Service, Sarbanes Cooperative Oxford Laboratory

Publication_Date: 1911

Publication_Time: Unknown

Title: **Yatesbrs**

Geospatial_Data_Presentation_Form: vector digital data

Series_Information:

Publication_Information:

Online_Linkage: <http://dnrweb.dnr.state.md.us/gis/data/index.html>

Description:

Abstract: Polygon delineation of Maryland oyster bottom as surveyed by C.C. Yates, circa 1906.

Purpose:

This file was created for Maryland DNR planning purposes, specifically for the purpose of managing Maryland's oyster resource.

Supplemental_Information: Boundaries were generated using Dbase III+ and MapInfo software based on coordinate information found in publications by the Department of Commerce and Labor, Bureau of the Coast & Geodetic Survey and Bureau of Fisheries.

Time_Period_of_Content:

Time_Period_Information:

Range_of_Dates/Times:

Beginning_Date: 01/01/1906

Beginning_Time: unknown

Ending_Date: 12/31/1911

Ending_Time: unknown

Currentness_Reference: publication date

Status:

Progress: Complete

Maintenance_and_Update_Frequency: None planned

Spatial_Domain:

Bounding_Coordinates:

West_Bounding_Coordinate: -76.863501

East_Bounding_Coordinate: -75.186116

North_Bounding_Coordinate: 39.295060

South_Bounding_Coordinate: 37.898347

Keywords:

Theme:

Theme_Keyword_Thesaurus: none

Theme_Keyword: oysters

Theme_Keyword: oyster bars

Place:

Place_Keyword_Thesaurus: none

Place_Keyword: Maryland

Stratum:

Stratum_Keyword: Chesapeake Bay seafloor

Temporal:

Temporal_Keyword: early 1900's

Temporal_Keyword: 1906

Access_Constraints: none

Use_Constraints: The Department of Natural Resources makes no warranty, expressed or implied, as to the use or appropriateness of Spatial Data, and there are no warranties of merchantability or fitness for a particular purpose or use.

The information contained in Spatial Data is from publicly available sources, but no representation is made as to the accuracy or completeness of Spatial Data. The Department of Natural Resources shall not be subject to liability for human error, error due to software conversion, defect, or failure of machines, or any material used in the connection with the machines, including tapes, disks, CD-ROM's or DVD-ROM's and energy. The Department of Natural Resources shall not be liable for any lost profits, consequential damages, or claims against the Department of Natural Resources by third parties. The liability of the Department of Natural Resources for damage regardless of the form of the action shall not exceed any distribution fees that may have been paid in obtaining Spatial Data.

Point_of_Contact:

Contact_Information:

Contact_Person_Primary:

Contact_Person: Kelly Greenhawk

Contact_Organization: Maryland Department of Natural Resources

Contact_Position: DP Programmer / Analyst

Contact_Address:

Address_Type: mailing address

Address: 904 South Morris Street

City: Oxford

State_or_Province: Maryland

Postal_Code: 21654

Country: USA

Contact_Voice_Telephone: 410-226-0078

Contact_TDD/TTY_Telephone: none

Contact_Facsimile_Telephone: 410-226-0120

Contact_Electronic_Mail_Address: kgreenhawk@dnr.state.md.us

Hours_of_Service: Monday through Friday 0830 - 1700 (EST)

Data_Set_Credit: Maryland Department of Natural Resources, Sarbanes Cooperative Oxford Laboratory

Security_Information:

Security_Classification_System: none

Security_Classification: Unclassified

Security_Handling_Description: none

Native_Data_Set_Environment: Microsoft Windows XP Version 5.1 (Build 2600) Service Pack 2; ESRI ArcCatalog 9.1.0.722

Data_Quality_Information:

Attribute_Accuracy:

Attribute_Accuracy_Report: Dataset was derived from coordinate information in Department of Commerce & Labor publication entitled "Survey of Maryland Oyster Bars, 1906 - 1912", Parts I and II

Positional_Accuracy:

Horizontal_Positional_Accuracy:

Horizontal_Positional_Accuracy_Report: *****

Vertical_Positional_Accuracy:

Vertical_Positional_Accuracy_Report: *****

Lineage:

Process_Step:

Process_Description:

The spatial layer that this dataset is derived from was originally in the form of a MapInfo file, and consisted of a series of polylines (shoreline) and polygons (islands). MapInfo was used to join adjacent polylines until the entire Chesapeake Bay was represented by one large "donut-polygon". Islands in this secondary file were represented by holes. (DNR filename "Baygon"). MapInfo's Universal Translator was used to convert the file "Baygon" to an ArcView shapefile. A copy of the shapefile was then made in ArcMap and used as the base file. The polygons in this dataset (COMPRRegions.shp) were created by using the Geoprocessing Wizard in ArcMap version 8.2. Specifically, each polygon was created by clipping a copy of the original (baywide) polygons in a temporary dataset with hand-drawn polygons.

Process_Step:

Process_Description: Metadata imported.

Source_Used_Citation_Abbreviation: C:\arcgis\arcexe82\Metadata\Stylesheets\DNRTemplate.xml

Process_Step:

Process_Description: Metadata imported.

Source_Used_Citation_Abbreviation: Z:\All\SCOLdata\Metadata\yatesbrs.shp.xml
 Process_Step:
 Process_Description: Dataset copied.
 Source_Used_Citation_Abbreviation:
 Spatial_Data_Organization_Information:
 Direct_Spatial_Reference_Method: Vector
 Point_and_Vector_Object_Information:
 SDTS_Terms_Description:
 SDTS_Point_and_Vector_Object_Type: G-polygon
 Point_and_Vector_Object_Count: 769
 Spatial_Reference_Information:
 Horizontal_Coordinate_System_Definition:
 Planar:
 Grid_Coordinate_System:
 Grid_Coordinate_System_Name: State Plane Coordinate System 1983
 State_Plane_Coordinate_System:
 SPCS_Zone_Identifier: 1900
 Lambert_Conformal_Conic:
 Standard_Parallel: 38.300000
 Standard_Parallel: 39.450000
 Longitude_of_Central_Meridian: -77.000000
 Latitude_of_Projection_Origin: 37.666667
 False_Easting: 400000.000000
 False_Northing: 0.000000
 Planar_Coordinate_Information:
 Planar_Coordinate_Encoding_Method: coordinate pair
 Coordinate_Representation:
 Abscissa_Resolution: 0.000256
 Ordinate_Resolution: 0.000256
 Planar_Distance_Units: meters
 Geodetic_Model:
 Horizontal_Datum_Name: North American Datum of 1983
 Ellipsoid_Name: Geodetic Reference System 80
 Semi-major_Axis: 6378137.000000
 Denominator_of_Flattening_Ratio: 298.257222
 Entity_and_Attribute_Information:
 Detailed_Description:
 Entity_Type:
 Entity_Type_Label: yatesbrs
 Attribute:
 Attribute_Label: FID
 Attribute_Definition: Internal feature number.
 Attribute_Definition_Source: ESRI
 Attribute_Domain_Values:
 Unrepresentable_Domain: Sequential unique whole numbers that are automatically generated.
 Attribute:
 Attribute_Label: Shape
 Attribute_Definition: Feature geometry.
 Attribute_Definition_Source: ESRI
 Attribute_Domain_Values:
 Unrepresentable_Domain: Coordinates defining the features.
 Attribute:
 Attribute_Label: YATESBRS_I
 Attribute_Definition: Name given to oyster bar by Yates.
 Attribute_Definition_Source: Maryland Department of Natural Resources
 Attribute:

Attribute_Label: YATESNAME
Attribute_Definition: Number of corner coordinates published for oyster bar.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: NUMCORN
Attribute_Definition: Reference code - Volume number and page number coordinates were taken from
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: REFCODE
Attribute_Definition: Two letter code for county within which oyster bar lies
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: COUNTY
Attribute_Definition: Longitude of centroid value, expressed in negative decimal degrees and calculated from MapInfo
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: SQACRES
Attribute_Definition: Latitude of centroid value, expressed in negative decimal degrees and calculated from MapInfo
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: CENTROIDX
Attribute_Definition: Size of oyster bar, in acres, calculated in MapInfo
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: CENTROIDY
Attribute:
Attribute_Label: SQMETERS
Attribute:
Attribute_Label: CALCDACRES

Overview_Description:

Entity_and_Attribute_Detail_Citation: *****

Distribution_Information:

Resource_Description: Downloadable Data

Standard_Order_Process:

Digital_Form:

Digital_Transfer_Information:

Transfer_Size: 0.112

Distribution_Information:

Distributor:

Contact_Information:

Contact_Person_Primary:

Contact_Person: Kelly Greenhawk

Contact_Organization: Maryland Department of Natural Resources

Contact_Position: DP Programmer / Analyst

Contact_Address:

Address_Type: mailing and physical address

Address: Sarbanes Cooperative Oxford Laboratory

Address: 904 South Morris Street

City: Oxford

State_or_Province: Maryland

Postal_Code: 21654

Country: USA

Contact_Voice_Telephone: 410-226-0078

Contact_TDD/TTY_Telephone: none

Contact_Facsimile_Telephone: 410-226-0120

Contact_Electronic_Mail_Address: kgreenhawk@dnr.state.md.us

Hours_of_Service: Monday through Friday 0800 - 1700

Distribution_Liability: This data represents the results of data generated for a specific Maryland Department of Natural Resources management activity. As such, they are only valid for their intended use, content, time, and accuracy specifications. The user is responsible for any application of the data for other than it's intended purpose.

Standard_Order_Process:

Digital_Form:

Digital_Transfer_Information:

Format_Name: ARCE shapefile

Format_Information_Content: Data is normally distributed as a statewide dataset - Md State Plane NAD 83 meters

File-Decompression_Technique: no compression applied

Transfer_Size: 0.112

Digital_Transfer_Option:

Online_Option:

— Online_Computer_and_Operating_System: <http://dnrweb.dnr.state.md.us/gis/data/index.html>

Metadata_Reference_Information:

Metadata_Date: 20050629

Metadata_Contact:

Contact_Information:

Contact_Person_Primary:

Contact_Person: Kelly Greenhawk

Contact_Organization: Maryland Department of Natural Resources

Contact_Position: DP Programmer / Analyst

Contact_Address:

Address_Type: mailing and physical address

Address: Sarbanes Cooperative Oxford Laboratory

Address: 904 South Morris Street

City: Oxford

State_or_Province: Maryland

Postal_Code: 21654

Country: USA

Contact_Voice_Telephone: 410-226-0078

Contact_TDD/TTY_Telephone: none

Contact_Facsimile_Telephone: 410-226-0120

Contact_Electronic_Mail_Address: kgreenhawk@dnr.state.md.us

Hours_of_Service: Monday through Friday 0830 - 1700 (EST)

Metadata_Standard_Name: FGDC Content Standards for Digital Geospatial Metadata

Metadata_Standard_Version: FGDC-STD-001-1998

Metadata_Time_Convention: local time

Metadata_Security_Information:

Metadata_Security_Classification: Unclassified

Metadata_Extensions:

Online_Linkage: <http://www.esri.com/metadata/esriprof80.html>

Profile_Name: ESRI Metadata Profile

"DNRRepletion.shp"

Identification_Information:

Citation:

Citation_Information:

Originator: Maryland Department of Natural Resources, Fisheries Service, Cooperative Oxford Laboratory

Publication_Date: 20030409

Publication_Time: Unknown

Title: DNR Repletion

Geospatial_Data_Presentation_Form: vector digital data

Series_Information:

Publication_Information:

Publication_Place: Oxford, Maryland

Publisher: Maryland Department of Natural Resources

Online_Linkage: <http://dnrweb.dnr.state.md.us/gis/data/index.html>

Larger_Work_Citation:

Citation_Information:

Originator: John Hess, MDNR Shellfish Program, Deale Island Hatchery, Deale Island, Md

Publication_Date: Unpublished Material

Title: DNR Shellfish Oyster Propagation Activity summary reports

Description:

Abstract: Polygon delineation of annual oyster repletion activities undertaken by the Maryland Department of Natural Resources, Shellfish Program.

Purpose: This file was created for MDNR planning purposes, specifically for the purpose of tracking the activities of the Department's Shellfish Repletion Program.

Supplemental_Information: Boundaries were generated using MapInfo software (version 5) from coordinates taken from planting forms completed by the Department's Shellfish Program. The file was later migrated to Arcview.

Time_Period_of_Content:

Time_Period_Information:

Range_of_Dates/Times:

Beginning_Date: 04/10/2000

Beginning_Time: unknown

Ending_Date: changes annually

Currentness_Reference: publication date

Status:

Progress: In work

Maintenance_and_Update_Frequency: As needed

Spatial_Domain:

Bounding_Coordinates:

West_Bounding_Coordinate: -76.977947

East_Bounding_Coordinate: -75.860381

North_Bounding_Coordinate: 39.189665

South_Bounding_Coordinate: 37.958141

Keywords:

Theme:

Theme_Keyword_Thesaurus: none

Theme_Keyword: oyster repletion

Theme_Keyword: dredged shell

Theme_Keyword: seed

Theme_Keyword: oyster propagation

Place:

Place_Keyword_Thesaurus: none

Place_Keyword: Maryland

Stratum:

Stratum_Keyword: Chesapeake Bay

Temporal:

Access_Constraints: none

Use_Constraints: The Department of Natural Resources makes no warranty, expressed or implied, as to the use or appropriateness of Spatial Data, and there are no warranties of merchantability or fitness for a particular purpose or use. The information contained in Spatial Data is from publicly available sources, but no representation is made as to the accuracy or completeness of Spatial Data. The Department of Natural Resources shall not be subject to liability for human error, error due to software conversion, defect, or failure of machines, or any material used in the connection with the machines, including tapes, disks, CD-ROM's or DVD-ROM's and energy. The Department of Natural Resources shall not be liable for any lost profits, consequential damages, or claims against the Department of Natural Resources by third

parties. The liability of the Department of Natural Resources for damage regardless of the form of the action shall not exceed any distribution fees that may have been paid in obtaining Spatial Data.

Point_of_Contact:

Contact_Information:

Contact_Person_Primary:

Contact_Person: Kelly Greenhawk

Contact_Organization: Maryland Department of Natural Resources

Contact_Position: DP Programmer / Analyst

Contact_Address:

Address_Type: mailing address

Address: 904 South Morris Street

City: Oxford

State_or_Province: Maryland

Postal_Code: 21654

Country: USA

Contact_Voice_Telephone: 410-226-0078

Contact_TDD/TTY_Telephone: none

Contact_Facsimile_Telephone: 410-226-0120

Contact_Electronic_Mail_Address: kgreenhawk@dnr.state.md.us

Hours_of_Service: Monday through Friday 0830 - 1700 (EST)

Data_Set_Credit: Data set was produced by MDNR, Fisheries Service staff at the Oxford Laboratory, Oxford, Maryland.

Security_Information:

Security_Classification_System: none

Security_Classification: Unclassified

Security_Handling_Description: none

Native_Data_Set_Environment: Microsoft Windows XP Version 5.1 (Build 2600) Service Pack 2; ESRI ArcCatalog 9.1.0.722

Data_Quality_Information:

Attribute_Accuracy:

Attribute_Accuracy_Report: Dataset was derived from coordinates collected by MDNR's Shellfish Program personnel in the field as material was deployed. Beginning in 1999, coordinates were taken from DGPS (Northstar DGPS model 951XD) ; prior to 1999, coordinates were taken from Loran.

Positional_Accuracy:

Horizontal_Positional_Accuracy:

Horizontal_Positional_Accuracy_Report: No report available.

Vertical_Positional_Accuracy:

Vertical_Positional_Accuracy_Report: No report available.

Lineage:

Source_Information:

Type_of_Source_Media: planting form completed in the form

Process_Step:

Process_Description: This spatial layer was created using MapInfo software (version 5x). MapInfo's Universal Translator was then used to convert the file to an ArcView shapefile.

Process_Step:

Process_Description: Metadata imported.

Source_Used_Citation_Abbreviation: C:\DOCUME~1\KGREEN~1\LOCALS~1\Temp\xml6D.tmp

Process_Step:

Process_Description: Metadata imported.

Source_Used_Citation_Abbreviation: C:\DOCUME~1\KGREEN~1\LOCALS~1\Temp\xml86.tmp

Process_Step:

Process_Description: Metadata imported.

Source_Used_Citation_Abbreviation: Z:\KGreenhawk\Metadata from SCOLdata\DNRRepletion.shp.xml

Spatial_Data_Organization_Information:

Direct_Spatial_Reference_Method: Vector

Point_and_Vector_Object_Information:

SDTS_Terms_Description:

SDTS_Point_and_Vector_Object_Type: G-polygon
 Point_and_Vector_Object_Count: 181
 Spatial_Reference_Information:
 Horizontal_Coordinate_System_Definition:
 Planar:
 Grid_Coordinate_System:
 Grid_Coordinate_System_Name: State Plane Coordinate System 1983
 State_Plane_Coordinate_System:
 SPCS_Zone_Identifier: 1900
 Lambert_Conformal_Conic:
 Standard_Parallel: 38.300000
 Standard_Parallel: 39.450000
 Longitude_of_Central_Meridian: -77.000000
 Latitude_of_Projection_Origin: 37.666667
 False_Easting: 400000.000000
 False_Northing: 0.000000
 Planar_Coordinate_Information:
 Planar_Coordinate_Encoding_Method: coordinate pair
 Coordinate_Representation:
 Abscissa_Resolution: 0.000256
 Ordinate_Resolution: 0.000256
 Planar_Distance_Units: meters
 Geodetic_Model:
 Horizontal_Datum_Name: North American Datum of 1983
 Ellipsoid_Name: Geodetic Reference System 80
 Semi-major_Axis: 6378137.000000
 Denominator_of_Flattening_Ratio: 298.257222
 Entity_and_Attribute_Information:
 Detailed_Description:
 Entity_Type:
 Entity_Type_Label: DNRRepletion
 Attribute:
 Attribute_Label: FID
 Attribute_Definition: Internal feature number.
 Attribute_Definition_Source: ESRI
 Attribute_Domain_Values:
 Unrepresentable_Domain: Sequential unique whole numbers that are automatically generated.
 Attribute:
 Attribute_Label: Shape
 Attribute_Definition: Feature geometry.
 Attribute_Definition_Source: ESRI
 Attribute_Domain_Values:
 Unrepresentable_Domain: Coordinates defining the features.
 Attribute:
 Attribute_Label: DATE_
 Attribute_Definition: Date of repletion activity; if seed planting, may be the date buoys were deployed
 Attribute_Definition_Source: Maryland Department of Natural Resources
 Attribute:
 Attribute_Label: OPERATORS
 Attribute_Definition: From fieldsheet, names of staff operating vessel
 Attribute_Definition_Source: Maryland Department of Natural Resources
 Attribute:
 Attribute_Label: NAME
 Attribute_Definition: From fieldsheet, code assigned to bar by Shellfish Program staff, usually taken from Maryland Oyster Bars publication (file mdoysbrs)
 Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:
Attribute_Label: NOB
Attribute_Definition: From fieldsheet, NOB number within which activity falls
Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:
Attribute_Label: ACCURACY
Attribute_Definition: From fieldsheet, estimated accuracy of GPS
Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:
Attribute_Label: DESC_
Attribute_Definition: From fieldsheet, description of repletion activity; usually bar name, current year and code for material deployed
Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:
Attribute_Label: ACRES
Attribute_Definition: From fieldsheet, estimated number of acres covered by deployed material
Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:
Attribute_Label: DEPTH
Attribute_Definition: From fieldsheet, depth range in feet at planting location
Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:
Attribute_Label: BOTTOM
Attribute_Definition: From fieldsheet, bottom type(s) of are where material was deployed; codes are comma delimited and are as follows: SH = shell; M or MD mud; S = sand; CL = clay
Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:
Attribute_Label: COMMENTS
Attribute_Definition: From fieldsheet, comments recorded by field staff (eg. funding agency) or comments from data producer (eg. incomplete coordinates on fieldsheet)
Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:
Attribute_Label: NUMCOORDS
Attribute_Definition: Number of coordinates provided on fieldsheet
Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:
Attribute_Label: LAT1
Attribute_Definition: From fieldsheet, latitude of first corner; expressed in degrees and decimal minutes to three places
Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:
Attribute_Label: LONG1
Attribute_Definition: From fieldsheet, longitude of first corner; expressed in degrees and decimal minutes to three places
Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:
Attribute_Label: LAT2
Attribute_Definition: From fieldsheet, latitude of second corner; expressed in degrees and decimal minutes to three places
Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:
Attribute_Label: LONG2
Attribute_Definition: From fieldsheet, longitude of second corner; expressed in degrees and decimal minutes to three places
Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:
Attribute_Label: LAT3

Attribute_Definition: From fieldsheet, latitude of third corner; expressed in degrees and decimal minutes to three places
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LONG3
Attribute_Definition: From fieldsheet, longitude of third corner; expressed in degrees and decimal minutes to three places
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LAT4
Attribute_Definition: From fieldsheet, latitude of fourth corner; expressed in degrees and decimal minutes to three places
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LONG4
Attribute_Definition: From fieldsheet, longitude of fourth corner; expressed in degrees and decimal minutes to three places
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LAT5
Attribute_Definition: From fieldsheet, latitude of center coordinate; expressed in degrees and decimal minutes to three places
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LONG5
Attribute_Definition: From fieldsheet, longitude of center coordinate; expressed in degrees and decimal minutes to three places
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: YEAR
Attribute_Definition: Four digit year, derived from field DATE_
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: RECTYPE
Attribute_Definition: Record type; "SHELL" for dredged shell or fresh shell planting; "SEED" for seed plant; OTHER for other types of material
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: ACCESS
Attribute_Definition: Accession number; internal tracking number used by data producer; first two characters represent 2 digit year; Characters 3 through 5 are alphabetic and are a code for group responsible for planting activity ("DNR" denotes Md Department of Natural Resources in this dataset); Characters 6 through 11 are the oyster bar code for the oyster bar within which the activity falls (from MDNR - Maryland Oyster Bar publication); Characters 12 through 14 denote type of material deployed (DSH = dredged shell; FSH = fresh shell; SSE = seed from state seed area; HSE = hatchery seed; characters 15-18 represent month and day of planting; the last character is intended to avoid duplicate accession / tracking numbers. Value will be the letter "A" if the activity on this bar was the only one of its kind that day, but will be assigned a "B", "C", etc. for additional activities. Duplication will only occur if the same group plants the same material on different locations on the same bar during the same day.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: TRIB
Attribute_Definition: Tributary name within which activity falls, derived from barcode
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: BARNAME
Attribute_Definition: Barname within which activity falls or intersects

Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: BARCODE
Attribute_Definition: Six character barcode, derived from MdOysBrS file
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: MATERIAL
Attribute_Definition: Material code for type of material deployed; DSH = dredged shell; FSH = fresh shell; SSE = seed from state seed area; HSE = hatchery seed
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: MATSOURCE
Attribute_Definition: From summary report, source of material deployed; Prior to 2002 code is a 2-4 character alphabetic code. Definitions for 2000 are as follows: BEA = Bald Eagle Addition #2; B = Bugby SSA; WCT = Wild Cherry Tree SSA; GM = Great Marsh; OC = Oyster Creek SSA; BC-N = Back Cove SSA; HSB = Horse Shoe Bend; GR = Gravelly Run; Definitions for 2001 are as follows: B = Bugby SSA; BN = Bugby North SSA; WCT = Wild Cherry Tree SSA; BC = Back Cove SSA (63 acres); Beginning in 2002, Shellfish staff used 6 character code from MdOysBrS file
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: BUSHEL
Attribute_Definition: From summary report, number of bushels of material deployed; A negative 8 in this field denotes data that is missing at publication time, but will be updated in the near future.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: X1
Attribute_Definition: Calculated field, derived from Long1 (decimal degree equivalent) and used to generate polygon in MapInfo software
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: Y1
Attribute_Definition: Calculated field, derived from Lat1 (decimal degree equivalent) and used to generate polygon in MapInfo software
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: X2
Attribute_Definition: Calculated field, derived from Long2 (decimal degree equivalent) and used to generate polygon in MapInfo software
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: Y2
Attribute_Definition: Calculated field, derived from Lat2 (decimal degree equivalent) and used to generate polygon in MapInfo software
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: X3
Attribute_Definition: Calculated field, derived from Long3 (decimal degree equivalent) and used to generate polygon in MapInfo software
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: Y3
Attribute_Definition: Calculated field, derived from Lat3 (decimal degree equivalent) and used to generate polygon in MapInfo software
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: X4

Attribute_Definition: Calculated field, derived from Long4 (decimal degree equivalent) and used to generate polygon in MapInfo software

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: Y4

Attribute_Definition: Calculated field, derived from Lat4 (decimal degree equivalent) and used to generate polygon in MapInfo software

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: X5

Attribute_Definition: Calculated field, derived from Long5 (decimal degree equivalent) and used to generate polygon in MapInfo software

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: Y5

Attribute_Definition: Calculated field, derived from Lat5 (decimal degree equivalent) and used to generate polygon in MapInfo software

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: PAGE

Attribute_Definition: From summary report, page number where plant can be referenced

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: LOCDETAILS

Attribute_Definition: From summary report, details regarding location of plant on oyster bar

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: CATEGORY

Attribute_Definition: From summary report, activity purpose / category; Categories are as follows: "SEED TRANSPLANTED FOR NOB IMPROVEMENT BY DNR", "DSH FOR SEED OYSTER PRODUCTION BY DNR", "DSH FOR NOB IMPROVEMENT BY DNR", "DSH FOR CO SEED OYSTER PRODUCTION AND/OR GROWOUT, FSH FOR NOB IMPROVEMENT BY DNR; NOT LISTED (planting not found in summary report)

Attribute_Definition_Source: Maryland Department of Natural Resources

Overview_Description:

Entity_and_Attribute_Detail_Citation: *****

Distribution_Information:

Resource_Description: Downloadable Data

Standard_Order_Process:

Digital_Form:

Digital_Transfer_Information:

Transfer_Size: 0.175

Distribution_Information:

Distributor:

Contact_Information:

Contact_Person_Primary:

Contact_Person: Kelly Greenhawk

Contact_Organization: Maryland Department of Natural Resources

Contact_Position: DP Programmer / Analyst

Contact_Address:

Address_Type: mailing and physical address

Address: Sarbanes Cooperative Oxford Laboratory

Address: 904 South Morris Street

City: Oxford

State_or_Province: Maryland

Postal_Code: 21654

Country: USA

Contact_Voice_Telephone: 410-226-0078
Contact_TDD/TTY_Telephone: *****
Contact_Facsimile_Telephone: 410-226-0120
Contact_Electronic_Mail_Address: kgreenhawk@dnr.state.md.us
Hours_of_Service: Monday through Friday 0800 - 1700

Distribution_Liability: This data represents the results of data generated for specific Maryland Department of Natural Resources management activity. As such, they are only valid for their intended use, content, time, and accuracy specifications. The user is responsible for any application of the data for other than it's intended purpose.

Standard_Order_Process:

Digital_Form:

Digital_Transfer_Information:

Format_Name: ARCE shapefile

Format_Information_Content: Data is normally distributed as a statewide dataset - Md State Plane NAD 83 meters

File-Decompression_Technique: no compression applied

Transfer_Size: 0.175

Digital_Transfer_Option:

Online_Option:

Online_Computer_and_Operating_System: <http://dnrweb.dnr.state.md.us/gis/data/index.html>

Metadata_Reference_Information:

Metadata_Date: 20050629

Metadata_Contact:

Contact_Information:

Contact_Person_Primary:

Contact_Person: Kelly Greenhawk

Contact_Organization: Maryland Department of Natural Resources

Contact_Position: DP Programmer / Analyst

Contact_Address:

Address_Type: mailing and physical address

Address: Sarbanes Cooperative Oxford Laboratory

Address: 904 South Morris Street

City: Oxford

State_or_Province: Maryland

Postal_Code: 21654

Country: USA

Contact_Voice_Telephone: 410-226-0078

Contact_TDD/TTY_Telephone: none

Contact_Facsimile_Telephone: 410-226-0120

Contact_Electronic_Mail_Address: kgreenhawk@dnr.state.md.us

Hours_of_Service: Monday through Friday 0830 - 1700 (EST)

Metadata_Standard_Name: FGDC Content Standards for Digital Geospatial Metadata

Metadata_Standard_Version: FGDC-STD-001-1998

Metadata_Time_Convention: local time

Metadata_Extensions:

Online_Linkage: <http://www.esri.com/metadata/esriprof80.html>

Profile_Name: ESRI Metadata Profile

Metadata_Extensions:

Online_Linkage: <http://www.esri.com/metadata/esriprof80.html>

Profile_Name: ESRI Metadata Profile

"MdSancRes.shp"

Identification_Information:

Citation:

Citation_Information:

Originator: Kelly Greenhawk, MDNR, Fisheries Service

Publication_Date: Unpublished Material

Title: **MdSancRes**

Geospatial_Data_Presentation_Form: vector digital data

Online_Linkage: \\OX0057178\C\$\Documents and Settings\kgreenhawk\My Documents\ArcData\Bndfile\Oysbound\mdsanres.shp

Description:

Abstract: Polygon representation of oyster recovery activities undertaken in Maryland's oyster sanctuaries and reserves.

Purpose: File was created for the purpose of tracking oyster recovery efforts in Maryland's oyster sanctuaries and reserves.

Supplemental_Information:

File was created by Kelly Greenhawk, MDNR, Fisheries Service, Cooperative Oxford Laboratory but is maintained and distributed by Eric Campbell, MDNR, Fisheries Service, Shellfish Program. File is updated on a continual basis, as planting forms are submitted by the various Maryland partners. Due to the occasional receipt of incomplete forms, some fields in this file may be missing data, or may contain incomplete information. Records are flagged as suspect in this instance, in either the field COMMENTS or in the field OBJCOMM.

Time_Period_of_Content:

Time_Period_Information:

Range_of_Dates/Times:

Beginning_Date: 1997

Ending_Date: unknown

Currentness_Reference: ground condition

Status:

Progress: In work

Maintenance_and_Update_Frequency: Continually

Spatial_Domain:

Bounding_Coordinates:

West_Bounding_Coordinate: -76.979840

East_Bounding_Coordinate: 2.567953

North_Bounding_Coordinate: 39.216303

South_Bounding_Coordinate: -0.000001

Keywords:

Theme:

Theme_Keyword_Thesaurus: none

Theme_Keyword: oysters

Theme_Keyword: sanctuary

Theme_Keyword: reserve

Theme_Keyword: restoration

Theme_Keyword: recovery

Place:

Place_Keyword_Thesaurus: none

Place_Keyword: Maryland

Place_Keyword: Chesapeake Bay

Stratum:

Stratum_Keyword_Thesaurus: none

Stratum_Keyword: estuary floor

Use_Constraints: The Department of Natural Resources makes no warranty, expressed or implied, as to the use or appropriateness of Spatial Data, and there are no warranties of merchantability or fitness for a particular purpose or use. The information contained in Spatial Data is from publicly available sources, but no representation is made as to the accuracy or completeness of Spatial Data. The Department of Natural Resources shall not be subject to liability for human error, error due to software conversion, defect, or failure of machines, or any material used in the connection with the machines, including tapes, disks, CD-ROM's or DVD-ROM's and energy. The Department of Natural Resources shall not be liable for any lost profits, consequential damages, or claims against the Department of Natural Resources by third parties. The liability of the Department of Natural Resources for damage regardless of the form of the action shall not exceed any distribution fees that may have been paid in obtaining Spatial Data.

Point_of_Contact:

Contact_Information:

Contact_Person_Primary:

Contact_Person: Eric Campbell

Contact_Organization: MDNR, Fisheries Service, Shellfish Program

Contact_Position: Biologist

Contact_Address:

Address_Type: mailing and physical address

Address: Maryland Department of Natural Resources

Address: Tawes State Office Building

Address: 580 Taylor Avenue

City: Annapolis

State_or_Province: MD

Postal_Code: 21401

Country: USA

Contact_Voice_Telephone: 410-260-8344

Contact_Facsimile_Telephone: 410-260-8279

Contact_Electronic_Mail_Address: ecampbell@dnr.state.md.us

Data_Set_Credit: MDNR, Fisheries Service

Native_Data_Set_Environment: Microsoft Windows XP Version 5.1 (Build 2600) Service Pack 2; ESRI ArcCatalog 9.1.0.722

Data_Quality_Information:

Completeness_Report: Records from forms that were submitted to MDNR with only one set of coordinates (as opposed to 4 sets of corner coordinates) are represented with a circular object, as opposed to a polygon object. These objects were created by using the buffer feature of ArcView 8x in conjunction with the centroid coordinate values.

Lineage:

Source_Information:

Type_of_Source_Media: paper forms, designed by MDNR's Shellfish Program

Source_Time_Period_of_Content:

Time_Period_Information:

Range_of_Dates/Times:

Beginning_Date: 1997

Ending_Date: 2002

Source_Currentness_Reference: ground condition

Process_Step:

Source_Used_Citation_Abbreviation: C:\DOCUME~1\KGREEN~1\LOCALS~1\Temp\xml15.tmp

Process_Step:

Process_Description: Metadata imported.

Source_Used_Citation_Abbreviation: J:\DATA\OysterData\MdSancRes.shp.xml

Process_Step:

Process_Description: Dataset copied.

Source_Used_Citation_Abbreviation:

Process_Step:

Process_Description: Dataset copied.

Source_Used_Citation_Abbreviation: C:\Documents and Settings\kgreenhawk\My Documents\ArcData\Help Campbell with Mdsanres updates (Geo coordsys)\MdSRsp83m

Process_Step:

Process_Description: Dataset moved.

Source_Used_Citation_Abbreviation: C:\Documents and Settings\kgreenhawk\My Documents\ArcData\Metadata templates\MdSR83mBKUP

Process_Step:

Process_Description: Metadata imported.

Source_Used_Citation_Abbreviation: C:\Documents and Settings\kgreenhawk\My Documents\ArcData\Help Campbell with Mdsanres updates (Geo coordsys)\MdSR83mBKUP.shp.xml

Spatial_Data_Organization_Information:

Direct_Spatial_Reference_Method: Vector

Point_and_Vector_Object_Information:
 SDTS_Terms_Description:
 SDTS_Point_and_Vector_Object_Type: G-polygon
 Point_and_Vector_Object_Count: 158

Spatial_Reference_Information:
 Horizontal_Coordinate_System_Definition:
 Planar:
 Grid_Coordinate_System:
 Grid_Coordinate_System_Name: State Plane Coordinate System 1983
 State_Plane_Coordinate_System:
 SPCS_Zone_Identifier: 1900
 Lambert_Conformal_Conic:
 Standard_Parallel: 38.300000
 Standard_Parallel: 39.450000
 Longitude_of_Central_Meridian: -77.000000
 Latitude_of_Projection_Origin: 37.666667
 False_Easting: 400000.000000
 False_Northing: 0.000000

Planar_Coordinate_Information:
 Planar_Coordinate_Encoding_Method: coordinate pair
 Coordinate_Representation:
 Abscissa_Resolution: 0.016384
 Ordinate_Resolution: 0.016384
 Planar_Distance_Units: meters

Geodetic_Model:
 Horizontal_Datum_Name: North American Datum of 1983
 Ellipsoid_Name: Geodetic Reference System 80
 Semi-major_Axis: 6378137.000000
 Denominator_of_Flattening_Ratio: 298.257222

Entity_and_Attribute_Information:
 Detailed_Description:
 Entity_Type:
 Entity_Type_Label: mdsancre

Attribute:
 Attribute_Label: FID
 Attribute_Definition: Internal feature number.
 Attribute_Definition_Source: ESRI
 Attribute_Domain_Values:
 Unrepresentable_Domain: Sequential unique whole numbers that are automatically generated.

Attribute:
 Attribute_Label: Shape
 Attribute_Definition: Feature geometry.
 Attribute_Definition_Source: ESRI
 Attribute_Domain_Values:
 Unrepresentable_Domain: Coordinates defining the features.

Attribute:
 Attribute_Label: RECTYPE
 Attribute_Definition: Type of record - SEED, SHELL or MULTI; object represents the boundaries of a seed planting activity, or deployment of shell or other material used as substrate for oysters
 Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:
 Attribute_Label: GROUP
 Attribute_Definition: Three character code for the name of the group which served as project leader for the activity;
 ANS = Academy of Natural Sciences; CBF = Chesapeake Bay Foundation; ORP = Oyster Recovery Partnership; DNR = Md. Department of Natural Resources; SRF = South River Federation; PRF = Potomac River Fisheries Commission; LCF = Living Classrooms Foundation; COE = Army Corps of Engineers

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute_Domain_Values:

Enumerated_Domain:

Enumerated_Domain_Value: ORP

Enumerated_Domain_Value_Definition: Oyster Recovery Partnership

Enumerated_Domain:

Enumerated_Domain_Value: CBF

Enumerated_Domain_Value_Definition: Chesapeake Bay Foundation

Enumerated_Domain:

Enumerated_Domain_Value: DNR

Enumerated_Domain_Value_Definition: Maryland Department of Natural Resources

Enumerated_Domain:

Enumerated_Domain_Value: COE

Enumerated_Domain_Value_Definition: Army Corp of Engineers

Enumerated_Domain:

Enumerated_Domain_Value: ANS

Enumerated_Domain_Value_Definition: Academy of Natural Science

Enumerated_Domain:

Enumerated_Domain_Value: LCF

Enumerated_Domain_Value_Definition: Living Classroom Foundation

Enumerated_Domain:

Enumerated_Domain_Value: PRF

Enumerated_Domain_Value_Definition: Potomac River Fisheries Commission

Enumerated_Domain:

Enumerated_Domain_Value: SRF

Enumerated_Domain_Value_Definition: South River Federation

Enumerated_Domain:

Enumerated_Domain_Value: MRA

Enumerated_Domain_Value_Definition: Magothy River Association

Attribute:

Attribute_Label: ACCESS

Attribute_Definition: Internal DNR tracking / accession number to identify activity; For more information on coding scheme, see Entity and Attribute Overview under Overview Description

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: OTHGROUPS

Attribute_Definition: Listing of other groups involved in activity

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: FUNDSOURCE

Attribute_Definition: Funding source for activity

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: LOCATION

Attribute_Definition: Location of activity, usually tributary name.

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: BARNAME

Attribute_Definition: Barname of oyster bar where activity took place, from form; barname will not always match barcode from MdOysbrs file since this field is reserved for information on the form

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: BOTTYPE

Attribute_Definition: Bottom type at location; Codes are as follows: ND = no data provided; S = sand; SH = shell; M = mud; GR = gravel; ST = stone; BSH = broken shell; CL = clay

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: DEPTH

Attribute_Definition: Depth or depth range in feet (minimum depth, followed by "-", followed by the maximum depth)

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: STARTDATE

Attribute_Definition: Date or beginning date of deployment if a range of dates is provided on form; when material was deployed

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: ENDDATE

Attribute_Definition: If a date range was provided, ending date of project. Field will be null if project was completed in one day or only start date was provided.

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: SANCT

Attribute_Definition: Logical field indicating whether activity took place in a sanctuary ("Y" or "N")

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: RESERVE

Attribute_Definition: Logical field indicating whether activity took place in a reserve ("Y" or "N")

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: ORA

Attribute_Definition: Logical field indicating whether activity took place in a designated Oyster Recovery Area

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: CLOSNAME

Attribute_Definition: Name of sanctuary or reserve / name of closure followed by a dash, then an "S" if the closure is a sanctuary or an "R" if the closure is a reserve; "NONE" is used for projects that do not fall within a designated sanctuary or reserve, once entered into the GIS.

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: LOCNOTES

Attribute_Definition: Notes on location of activity

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: MATERIAL1

Attribute_Definition: Material deployed; DSH = dredged shell; FSH = fresh shell; HSE = hatchery seed; USE = seed from unknown source; MUL = multiple materials; BBO = buy back oysters; CUL = cultchless oysters; 1YC = one year old cultch; SOS = spat on shell; STO = stone; UNK = material unknown; RUB = rubble; SLG = slag; SHL = shell of unknown type; RFB = reef ball(s); SPT = spat; OG1 = oyster gardeners' 1 year old oysters; FNS = fines; OTH = other material; If code OTH is used, see field OTHER for description of material. See Value Definition Source for more information.

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute_Domain_Values:

Enumerated_Domain:

Enumerated_Domain_Value_Definition_Source:

The abundance of material codes used in this file stems from the fact that the various partners use a variety of terms to describe the material they have deployed. Although a more streamlined coding system, with fewer material codes may be desirable in the future, the dataset creator has chosen to retain a more lengthy, specific coding system for now.

It is hoped that with the establishment of the Technical Committee called for in Maryland's Comprehensive Oyster Management Plan, data management guidelines can be written that will include a directive to the data producers to utilize a more streamlined coding system.

Attribute:

Attribute_Label: MATERIAL2

Attribute_Definition: Material deployed if more than one material was deployed; DSH = dredged shell; FSH = fresh shell; HSE = hatchery seed; USE = seed from unknown source; MUL = multiple materials; BBO = buy back oysters; CUL = cultchless oysters; 1YC = one year old cultch; SOS = spat on shell; STO = stone; UNK = material unknown; RUB = rubble; SLG = slag; SHL = shell of unknown type; RFB = reef ball(s); SPT = spat; OG1 = oyster gardeners' 1 year old oysters; FNS = fines; OTH = other material

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: MATERIAL3

Attribute_Definition: Material deployed if more than two different materials were deployed; DSH = dredged shell; FSH = fresh shell; HSE = hatchery seed; USE = seed from unknown source; MUL = multiple materials; BBO = buy back oysters; CUL = cultchless oysters; 1YC = one year old cultch; SOS = spat on shell; STO = stone; UNK = material unknown; RUB = rubble; SLG = slag; SHL = shell of unknown type; RFB = reef ball(s); SPT = spat; OG1 = oyster gardeners' 1 year old oysters; FNS = fines; OTH = other material

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: MATERIAL4

Attribute_Definition: Material deployed, if more than 3 types of material were deployed; DSH = dredged shell; FSH = fresh shell; HSE = hatchery seed; USE = seed from unknown source; MUL = multiple materials; BBO = buy back oysters; CUL = cultchless oysters; 1YC = one year old cultch; SOS = spat on shell; STO = stone; UNK = material unknown; RUB = rubble; SLG = slag; SHL = shell of unknown type; RFB = reef ball(s); SPT = spat; OG1 = oyster gardeners' 1 year old oysters; FNS = fines; OTH = other material

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: OTHER

Attribute_Definition: Additional description of material in MATERIAL1

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: AVGSIZE

Attribute_Definition: Average size of oysters, if seed was deployed, in centimeters. ND = no data provided

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: MATSOURCE

Attribute_Definition: Material source, i.e. hatchery name or vendor

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: BSTUSED

Attribute_Definition: Broodstock used, if applicable

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: TESTED

Attribute_Definition: Was the material, if seed material, tested for disease ? ("Y" or "N")

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: DISEASE

Attribute_Definition: Was disease present ? If the value of TESTED is "Y", field can be "Y", "N" or blank. A blank in this field denotes seed / oysters were tested but results of test were not indicated on form.

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: LAB_NAME

Attribute_Definition: Name of lab performing disease diagnosis

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: DERMOPREV

Attribute_Definition: Percent of sample testing positive for Dermo

Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: MSXPREV
Attribute_Definition: Percent of sample testing positive for MSX
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: CONTACT
Attribute_Definition: Contact name for laboratory
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: CONTACTPH
Attribute_Definition: Contact phone number for laboratory
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: OYS
Attribute_Definition: Number of oysters deployed, if seed activity. Value will be a negative nine if number of oysters was not provided on form.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: BU
Attribute_Definition: Number of bushels of material deployed; Value will be a negative nine if number of bushels was not provided on form.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: CUYDS
Attribute_Definition: Number of cubic yards of material deployed; Value will be a negative nine if number of cubic yards was not provided on form.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: CONFIG
Attribute_Definition: Configuration of planting, if noted on form; flat, mound, etc.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: ATTACH
Attribute_Definition: Logical field reflecting whether an attachment describing configuration details was submitted with the form.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: COMMENTS
Attribute_Definition: Additional information pertaining to activity.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: INCXY
Attribute_Definition: Logical field. Is coordinate information provided on form incomplete ? ("Y" or "N")
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: NUMCOORDS
Attribute_Definition: Number of corner coordinates provided on form
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: PRJORIGIN
Attribute_Definition: Project origin; Field will contain "ND" for no data for activities where project origin was not reported on form.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: PURP1

Attribute_Definition: Project purpose (text field 1 of 2 due to ESRI text field limitations). Field will contain "ND" for no data for activities where project purpose was not reported on form.

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: PURP2

Attribute_Definition: Project purpose (text field 2 of 2). Field will contain "ND" for no data for activities where project origin was not reported on form.

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: EXPRESULTS

Attribute_Definition: Expected results. Field will contain "ND" for no data for activities where expected results were not reported on form.

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: DDY1

Attribute_Definition: Decimal equivalent of corresponding LAT field

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: DDX1

Attribute_Definition: Decimal equivalent of corresponding LONG field

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: DDY2

Attribute_Definition: Decimal equivalent of corresponding LAT field

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: DDX2

Attribute_Definition: Decimal equivalent of corresponding LONG field

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: DDY3

Attribute_Definition: Decimal equivalent of corresponding LAT field

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: DDX3

Attribute_Definition: Decimal equivalent of corresponding LONG field

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: DDY4

Attribute_Definition: Decimal equivalent of corresponding LAT field

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: DDX4

Attribute_Definition: Decimal equivalent of corresponding LONG field

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: DDY5

Attribute_Definition: Decimal equivalent of corresponding LAT field

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: DDX5

Attribute_Definition: Decimal equivalent of corresponding LONG field

Attribute_Definition_Source: Maryland Department of Natural Resources

Attribute:

Attribute_Label: DDY6

Attribute_Definition: Decimal equivalent of corresponding LAT field

Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: DDX6
Attribute_Definition: Decimal equivalent of corresponding LONG field
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: OBJECTCOMM
Attribute_Definition: Comments pertaining to spatial object location or shape, from database manager.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: ACRES
Attribute_Definition: Area of object in acres, calculated in GIS.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: SQMETERS
Attribute_Definition: Area of object in square meters, calculated in GIS
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LAT1
Attribute_Definition: Latitude value of first corner of planted area, recorded from GPS. Values are expressed in degrees and decimal minutes.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LONG1
Attribute_Definition: Longitude value of first corner of planted area, recorded from GPS. Values are expressed in degrees and decimal minutes.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LAT2
Attribute_Definition: Latitude value of second corner of planted area, recorded from GPS. Values are expressed in degrees and decimal minutes.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LONG2
Attribute_Definition: Longitude value of second corner of planted area, recorded from GPS. Values are expressed in degrees and decimal minutes.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LAT3
Attribute_Definition: Latitude value of third corner of planted area, recorded from GPS. Values are expressed in degrees and decimal minutes.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LONG3
Attribute_Definition: Longitude value of third corner of planted area, recorded from GPS. Values are expressed in degrees and decimal minutes.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LAT4
Attribute_Definition: Latitude value of fourth corner of planted area, recorded from GPS. Values are expressed in degrees and decimal minutes.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LONG4
Attribute_Definition: Longitude value of fourth corner of planted area, recorded from GPS. Values are expressed in degrees and decimal minutes.

Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LAT5
Attribute_Definition: Latitude value of fifth corner of planted area, recorded from GPS. Values are expressed in degrees and decimal minutes.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LONG5
Attribute_Definition: Longitude value of fifth corner of planted area, recorded from GPS. Values are expressed in degrees and decimal minutes.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LAT6
Attribute_Definition: Latitude value of sixth corner of planted area, recorded from GPS. Values are expressed in degrees and decimal minutes.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: LONG6
Attribute_Definition: Longitude value of sixth corner of planted area, recorded from GPS. Values are expressed in degrees and decimal minutes.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: PRJPREP
Attribute_Definition: Description of preparations undertaken to ready site for deployment.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: CENTROIDX
Attribute_Definition: X value of centroid coordinate, expressed in meters. Field is calculated in GIS and used to center polygon objects for records where only one coordinate is provided.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: CENTROIDY
Attribute_Definition: Y value of centroid coordinate, expressed in meters. Field is calculated in GIS and used to center polygon objects for records where only one coordinate is provided.
Attribute_Definition_Source: Maryland Department of Natural Resources
Attribute:
Attribute_Label: TONS
Attribute_Definition: Amount of material deployed, expressed in tons. Value will be a negative nine if number of tons was not provided on form.
Attribute_Definition_Source: Maryland Department of Natural Resources
Overview_Description:
Entity_and_Attribute_Overview:
characters 1-2 represent year activity took place;
characters 3 through 5 represent project leader (see GROUP for more info);
characters 6 through 11 represent barcode from MdOysBrs file/publication;
characters 12-14 represent type of material deployed (see MATERIAL fields for more info)
characters 15-18 represent month and day of activity (MMDD);
character 19 is used to avoid duplicate accession / tracking codes, when the same group deploys the same material more than once on the same bar on the same date. The letter "A" is used for the first planting, the letter "B" is used for the second planting, the letter "C" is used for the third planting, etc..

Distribution_Information:
Distributor:
Contact_Information:
Contact_Person_Primary:
Contact_Person: Eric Campbell

Contact_Organization: MDNR, Fisheries Service, Shellfish Program
Contact_Position: Biologist
Contact_Voice_Telephone: 410-260-8344
Resource_Description: Downloadable Data
Standard_Order_Process:
Digital_Form:
Digital_Transfer_Information:
Transfer_Size: 0.087
Metadata_Reference_Information:
Metadata_Date: 20050629
Metadata_Contact:
Contact_Information:
Contact_Person_Primary:
Contact_Person: Eric Campbell
Contact_Organization: MDNR, Fisheries Service, Shellfish Program
Contact_Position: Biologist
Contact_Address:
Address_Type: REQUIRED: The mailing and/or physical address for the organization or individual.
City: REQUIRED: The city of the address.
State_or_Province: REQUIRED: The state or province of the address.
Postal_Code: REQUIRED: The ZIP or other postal code of the address.
Contact_Voice_Telephone: 410-260-8344
Contact_Facsimile_Telephone: 410-260-8279
Contact_Electronic_Mail_Address: ecampbell@dnr.state.md.us
Metadata_Standard_Name: FGDC Content Standards for Digital Geospatial Metadata
Metadata_Standard_Version: FGDC-STD-001-1998
Metadata_Time_Convention: local time
Metadata_Extensions:
Online_Linkage: <http://www.esri.com/metadata/esriprof80.html>
Profile_Name: ESRI Metadata Profile

"Lselines.dxf" and "Lease91.dbf"

Required Disclaimer

Any documents, maps, data, or publications that use the Natural Oyster Bar/Lease lines/coordinates provided must also include the following disclaimer:

"The Natural Oyster Bar/lease lines shown are for oyster management purposes only. For the official boundary(ies) consult the current official Natural Oyster Bar Chart."

The oyster bars/leases in this product are part of the conversion of the present mylar based Natural Oyster Bar Charts to digital format. They are based on bars that were delineated by the Bay Bottom Survey from 1975-1985, and amendments, Potomac River Oyster Survey of 1928, and amendments, and the Potomac River Bottom Survey of 1994, and amendments. They may differ from the bars shown on the current official mylar based Natural Oyster Bar Charts. Leases are based on the conversion of the coordinates from the original lease documents. They are based on surveys dating from 1912 to the present. Coordinates based on early surveys were converted by NADCON to NAD 83-91 values. Surveys performed after 1995 were done by GPS in the same coordinates.

It is the responsibility of the licensee to verify that this data is current. This office does not automatically notify licensees of changes/updates to this data.

Some leases have corner points that fall on the shoreline. In these instances this file may exhibit lines that cross the shoreline, or that connect from point to point over the water. In these cases the true extent of the lease should be verified from the lease document.

Any requests for all or part of this data must be referred to this office. The data is copyrighted by the State of Maryland, Department of Natural Resources.

The data is available only in DXF format, with coordinates in NAD 1983 (1991 adjustment) Maryland Zone 1900 State Plane coordinate Meters.

"Potential.shp"

Metadata was not available for the source file "Potential". See the document "Virginia Oyster Reef Restoration Map Atlas", dated August 2002 for more information on this dataset.

"Privlease.shp"

Identification_Information:

Citation:

Citation_Information:

Originator:

Comprehensive Coastal Inventory, Virginia Institute of Marine Science and Virginia Marine Resources Commission

Publication_Date: 2002

Title: Privateleases - coverage

Geospatial_Data_Presentation_Form: vector digital data

Publication_Information:

Publication_Place: Gloucester Point, Virginia

Publisher: Virginia Insititute of Marine Science

Description:

Abstract:

This polygon coverage delineates general survey boundaries of private oyster ground leases in Virginia. The coverage was generated from data provided by the Virginia Marine Resources Commission; the state agency responsible for regulating private oyster ground leases in Virginia. Data was originally received in AutoCad. Considerable manipulation was required to make the conversion. While private lease boundaries are legally defined, this coverage does not meet legal survey standards.

Purpose:

The data conversion and subsequent development of the GIS coverage was necessary for an oyster reef restoration targeting model being developed at the Virginia Institute of Marine Science.

Time_Period_of_Content:

Time_Period_Information:

Single_Date/Time:

Calendar_Date: 2002

Currentness_Reference: Data layer accurate to 2002.

Status:

Progress: Complete

Maintenance_and_Update_Frequency:

This coverage is not updated to reflect current status of private oyster ground leases in Virginia. The Virginia Marine Resources Commission does update their AutoCad system to reflect current standings, however, updates to the GIS record do not occur.

Spatial_Domain:

Bounding_Coordinates:

West_Bounding_Coordinate: -77.06352286

East_Bounding_Coordinate: -75.25314337

North_Bounding_Coordinate: 38.34948166

South_Bounding_Coordinate: 36.77505899

Keywords:

Theme:

Theme_Keyword_Thesaurus: none

Theme_Keyword: Private Oyster Ground Leases

Place:

Place_Keyword_Thesaurus: none

Place_Keyword: Virginia's Waters

Access_Constraints: none

Use_Constraints:

Acknowledgement the Virginia Marine Resources Commission should be included in products derived from this data. While private lease boundaries are legally defined, this coverage does not meet legal survey standards.

Point_of_Contact:

Contact_Information:

Contact_Person_Primary:

Contact_Person: Marcia Berman

Contact_Organization: Virginia Institute of Marine Science (VIMS)

Contact_Position: Director Comprehensive Coastal Inventory Program

Contact_Address:

Address_Type: mailing address

Address: P.O. Box 1346

City: Gloucester Point

State_or_Province: Virginia

Postal_Code: 23062

Country: USA

Contact_Voice_Telephone: (804) 684-7188

Contact_Facsimile_Telephone: (804) 684-7179

Contact_Electronic_Mail_Address: marcia@vims.edu

Data_Set_Credit:

The Virginia Marine Resources Commission is credited with origination of the database. The GIS coverage was generated by the Comprehensive Coastal Inventory Program at the Virginia Institute of Marine Science. The coverage was worked on by Sharon Killeen, Dan Schatt and Helen Woods.

Native_Data_Set_Environment:

SunOS, 5.7, sun4u UNIX ARC/INFO version 7.2.1

Cross_Reference:

Citation_Information:

Originator: The Virginia Marine Resources Commission

Publication_Date: 2002

Title: Virginia Private Oyster Leases

Geospatial_Data_Presentation_Form: map

Publication_Information:

Publication_Place: Newport News, Va

Publisher: Virginia Marine Resources Commission

Data_Quality_Information:

Logical_Consistency_Report:

Polygon topology is present with all polygons closed and coded for private.

Completeness_Report:

The most current data available was used to create the coverage.

Lineage:

Source_Information:

Source_Citation:

Citation_Information:

Originator: Virginia Marine Resources Commission

Publication_Date: 2002

Title: Virginia Private Oyster Leases

Geospatial_Data_Presentation_Form: map

Publication_Information:

Publication_Place: Newport News, VA

Publisher: Virginia Marine Resources Commission

Type_of_Source_Media: AutoCAD file

Source_Time_Period_of_Content:

Time_Period_Information:

Single_Date/Time:

Calendar_Date: 2002

Source_Currentness_Reference: Data layer accurate to 2002

Source_Citation_Abbreviation: VMRC

Source_Contribution: location of private oyster leases

Process_Step:

Process_Description:

An autocad file (VMRC) was turned into a shapefile using ArcInfo the conversion tool and projected using ArcInfo projection tools.

Process_Date: 2002

Process_Step:

Process_Description:

The coverage was then cleaned of all dangles and all polygons were closed using ArcInfo Arcedit.

Process_Date: 2002

Spatial_Data_Organization_Information:

Direct_Spatial_Reference_Method: Vector

Point_and_Vector_Object_Information:

SDTS_Terms_Description:

SDTS_Point_and_Vector_Object_Type: Point

Point_and_Vector_Object_Count: 5935

SDTS_Terms_Description:

SDTS_Point_and_Vector_Object_Type: String

Point_and_Vector_Object_Count: 12508

SDTS_Terms_Description:

SDTS_Point_and_Vector_Object_Type: GT-polygon composed of chains

Point_and_Vector_Object_Count: 5939

Spatial_Reference_Information:
 Horizontal_Coordinate_System_Definition:
 Planar:
 Grid_Coordinate_System:
 Grid_Coordinate_System_Name: Universal Transverse Mercator
 Universal_Transverse_Mercator:
 UTM_Zone_Number: 18
 Transverse_Mercator:
 Scale_Factor_at_Central_Meridian: 0.999600
 Longitude_of_Central_Meridian: -75.0000
 Latitude_of_Projection_Origin: 0.0000
 False_Easting: 500000
 False_Northing: 0.0000
 Planar_Coordinate_Information:
 Planar_Coordinate_Encoding_Method: coordinate pair
 Coordinate_Representation:
 Abscissa_Resolution: 0.84691375494
 Ordinate_Resolution: 0.84691375494
 Planar_Distance_Units: Meters
 Geodetic_Model:
 Horizontal_Datum_Name: North American Datum of 1983
 Ellipsoid_Name: GRS1980
 Semi-major_Axis: 6378206.4
 Denominator_of_Flattening_Ratio: 294.98

Entity_and_Attribute_Information:
 Overview_Description:
 Entity_and_Attribute_Overview:
 Each polygon in this coverage is a separate private lease polygon. Items listed in the .aat are from the autocad conversion.

PRILEASE83.PAT:

| COLUMN | ITEM NAME | WIDTH | OUTPUT | TYPE | N.DEC | ALTERNATE NAME |
|--------|---------------|-------|--------|------|-------|----------------|
| 1 | AREA | 4 | 12 | F | 3 | |
| 5 | PERIMETER | 4 | 12 | F | 3 | |
| 9 | PRILEASE83# | 4 | 5 | B | - | |
| 13 | PRILEASE83-ID | 4 | 5 | B | - | |
| 17 | LEASE | 8 | 8 | C | - | |

PRILEASE83.AAT:

| COLUMN | ITEM NAME | WIDTH | OUTPUT | TYPE | N.DEC | ALTERNATE NAME |
|--------|---------------|-------|--------|------|-------|----------------|
| 1 | FNODE# | 4 | 5 | B | - | |
| 5 | TNODE# | 4 | 5 | B | - | |
| 9 | LPOLY# | 4 | 5 | B | - | |
| 13 | RPOLY# | 4 | 5 | B | - | |
| 17 | LENGTH | 4 | 12 | F | 3 | |
| 21 | PRILEASE83# | 4 | 5 | B | - | |
| 25 | PRILEASE83-ID | 4 | 5 | B | - | |
| 29 | ENTITY | 14 | 14 | C | - | |
| 43 | HANDLE | 16 | 16 | C | - | |
| 59 | LAYER | 32 | 32 | C | - | |

| | | | | |
|---------------|-----|-----|---|---|
| 91 ELEVATION | 8 | 19 | F | 5 |
| 99 THICKNESS | 8 | 19 | F | 5 |
| 107 COLOR | 4 | 6 | B | - |
| 111 LINETYPE | 32 | 32 | C | - |
| 143 LINEWIDTH | 8 | 19 | F | 5 |
| 151 STYLE | 32 | 32 | C | - |
| 183 TEXT | 254 | 254 | C | - |

Entity_and_Attribute_Detail_Citation: none

Distribution_Information:

Distributor:

Contact_Information:

Contact_Organization_Primary:

Contact_Organization: Virginia Institute of Marine Science (VIMS)

Contact_Position: Director Comprehensive Coastal Inventory Program

Contact_Address:

Address_Type: mailing address

Address: P.O. Box 1346

City: Gloucester Point

State_or_Province: Virginia

Postal_Code: 23062

Country: USA

Contact_Voice_Telephone: (804) 684-7188

Contact_Instructions: Contact via email

Contact_Electronic_Mail_Address: marcia@vims.edu

Distribution_Liability:

The Comprehensive Coastal Inventory Program (CCI) at VIMS performs a service by distributing data generated by either CCI or public agencies which offer data without restriction or charge. CCI assumes no responsibility for data accuracy or precision, metadata completeness or correctness for digital information. CCI assumes no liability for misuse of any data which may arise as a result of any alteration, conversion, or combination with other data sources. As well, the timeliness and scale of these products must be considered when evaluating appropriate use.

Metadata_Reference_Information:

Metadata_Date: 20041103

Metadata_Contact:

Contact_Information:

Contact_Organization_Primary:

Contact_Organization: Virginia Institute of Marine Science (VIMS)

Contact_Person: Tamia Rudnicki

Contact_Position: GIS Programmer/Analyst

Contact_Address:

Address_Type: mailing address

Address: P.O. Box 1346

City: Gloucester Point

State_or_Province: Virginia

Postal_Code: 23062

Country: USA

Contact_Voice_Telephone: (804) 684-7181
Contact_Facsimile_Telephone: (804) 684-7179
Contact_Electronic_Mail_Address: tamia@vims.edu
Metadata_Standard_Name: FGDC Content Standards for Digital Geospatial
Metadata
Metadata_Standard_Version: FGDC-STD-001-1998
Metadata_Access_Constraints: none
Metadata_Use_Constraints: none

Generated by mp version 2.5.4 on Mon Nov 22 12:14:01 2004

Appendix 2.

Frequency data used to establish polygon reduction rates for Maryland oyster habitat, derived from data collected by MDNR. The shaded areas of the table contain the specific frequencies used.

Table 2. Relative abundance of classed bottom interrogations from 15 pooled acoustic benthic habitat surveys. Distributions came from bottom interrogations 1) within Yates Oyster Bars, and 2) within Maryland Bay Bottom Survey (MBBS) shell bottom polygons within Yates Bars. Because the Terrapin Sands survey was not within Yates bar boundaries it was not included in this analysis.

| Bottom class | Inside Yates bars | | MBBS shell bottom | |
|------------------------------------|-------------------|------------------------|-------------------|------------------------|
| | Frequency | Relative frequency (%) | Frequency | Relative frequency (%) |
| Clean shell | 181 | 0.62 | 81 | 1.16 |
| Mud | 9,490 | 32.52 | 1,431 | 20.58 |
| Sand | 15,572 | 53.36 | 3,338 | 48.00 |
| Heavily sedimented shell with mud | 2,432 | 8.33 | 1,505 | 21.64 |
| Heavily sedimented shell with sand | 864 | 2.96 | 359 | 5.16 |
| Lightly sedimented shell with sand | 95 | 0.33 | 84 | 1.21 |
| Gravel/cobble /boulder | 310 | 1.06 | 95 | 1.37 |
| Unidentified | 239 | 0.82 | 61 | 0.88 |
| Total | 29,183 | 100 | 6,954 | 100 |

Appendix 3.

Images showing the various bottom classifications identified by the acoustic surveys referenced in Appendix 2.





5. Heavily Sedimented Shell With Sand



6. Lightly Sedimented Shell With Sand



7. Clean Shell



8. Gravel, Cobble, Boulder

Appendix 4.

VBA code used to scale habitat polygons.

Appendix 4. VBA code used to scale habitat polygons. The line that reads ".Scale pPoint, 0.349857, 0.349857" contains the scale factor needed to reduce the Yates bar polygons to 12.24% of their original acreage. The scale factor is the square root of the percent reduction in area desired. For the MBBS polygons, a scale factor of .540092 was used.

```
Sub Scale_polygon()  
  
    Dim pMxDoc As IMxDocument  
    Dim pMap As IMap  
    Dim pTransform2D As ITransform2D  
    Dim pEnumFeature As IEnumFeature  
    Dim pFeature As IFeature  
    Dim pArea As IArea  
    Dim pPoint As IPoint  
    Dim pPolygon As IPolygon  
  
    Set pMxDoc = ThisDocument  
    Set pEnumFeature = pMxDoc.FocusMap.FeatureSelection  
    pEnumFeature.Reset  
    Set pFeature = pEnumFeature.Next  
  
    If pFeature.Shape.GeometryType = esriGeometryPolygon Then  
        Set pArea = pFeature.Shape  
    End If  
  
    Do While Not pFeature Is Nothing  
  
        Set pArea = pFeature.Shape  
        Set pPoint = pArea.Centroid  
  
        Set pTransform2D = pFeature.ShapeCopy  
        With pTransform2D  
            .Scale pPoint, 0.349857, 0.349857  
  
        End With  
        Set pFeature.Shape = pTransform2D  
        pFeature.Store  
        Set pFeature = pEnumFeature.Next  
    Loop  
  
    pMxDoc.ActiveView.Refresh  
  
End Sub
```

**Appendix B. Oyster population estimates for the
Maryland Portion of Chesapeake Bay 1994 – 2006 (Greenhawk and OConnell 2007)**

Oyster population estimates for the Maryland Portion of Chesapeake Bay 1994 – 2006

Prepared by: Maryland Department of Natural Resources, Fisheries Service

Kelly Greenhawk and Tom O'Connell

June 2007

INTRODUCTION

The Chesapeake Bay 2000 Agreement established a goal of increasing the Chesapeake Bay oyster population 10-fold by 2010 from its 1994 level of abundance. Upon the adoption of this goal, the National Oceanic and Atmospheric Administration Chesapeake Bay Stock Assessment Committee and U.S. Environmental Protection Agency Chesapeake Bay Program jointly funded an interstate research project to estimate the 1994 baseline population of oysters and to develop a methodology for producing annual oyster population estimates that could be used to measure progress toward achieving the 10-fold goal. This collaborative project between the Maryland Department of Natural Resources (MD DNR) and Virginia Institute of Marine Sciences resulted in annual oyster population estimates of small and market size oysters (spat estimates were not included) for the period 1994-2002. Estimates of the oyster population in Virginia have been produced annually since 2002. Oyster population estimates in Maryland have not been updated since 2002 until now primarily because of a lack of documentation of the methodology used to compute the 1994-2002 estimates. This report provides documentation for the methodology used to estimate Maryland's oyster population from 1994 to 2002, presents the assumptions associated with the estimates, and provides estimates of abundance through 2006.

METHODS

Oyster abundance was based on estimates of suitable oyster habitat and estimates of oyster density in each of eight basins in the Maryland portion of Chesapeake Bay. The steps used to calculate the population estimates are presented as follows.

1.0 Step 1 - Delineation of Maryland Basins

The Maryland portion of the Chesapeake Bay was divided into eight basins. This was done in an attempt to account for the significant observed variation in oyster abundance among basins. The following eight basins were established: 1) Chester River; 2) Eastern Bay; 3) Choptank River; 4) Little Choptank River; 5) Tangier Sound; 6) Potomac River; 7) Patuxent River; and 8) Chesapeake Bay mainstem (Figure 1).

Step 2 - Calculation of Available Oyster Habitat in Maryland Basins

Maryland's oyster population estimates are based on multiplying the annual measured oyster density by the amount of available oyster habitat within each basin. The most recent comprehensive survey of available oyster habitat is the Maryland Bay Bottom Survey (MBBS), conducted between 1976 and 1983. (Note: A digital dataset of the MBBS is available from MD DNR.) It is widely accepted in the scientific community that the amount and quality of existing oyster habitat in Maryland has declined significantly since the MBBS survey was conducted. Because the oyster population estimates are based on the amount and quality of available habitat, changes in habitat availability and quality since the MBSS was completed needed to be taken into account in our study.

A survey was conducted by the MD DNR Cooperative Oxford Laboratory between 1999 and 2000 at 15 oyster bars located throughout the Maryland portion of the Chesapeake Bay (Smith et al. 2005). The objective of the 1999-2000 MD DNR survey was to assess the relative abundance of oyster habitat within the cultch bottom classifications of both the 1978-1983 MBBS and a survey conducted by C.C. Yates between 1906-1911. The 15 bars surveyed were believed to be a fair representation of bars typically found in the Chesapeake Bay based upon a similarity of bottom types found at each bar. The MBBS classified oyster habitat into the following seven bottom classes: cultch, mud with cultch, sand with cultch, mud, sand, hard bottom and leased bottom (Smith et al. 2001). Areas of bottom with generous amounts of oyster shell were classified as cultch, while areas of scattered oyster shell were classified as either sand with cultch or mud with cultch. The cultch, sand with cultch and mud with cultch classifications were later categorized as either high quality habitat, low quality habitat, or lost habitat, for the purposes of this oyster population estimation effort (Table 1).

Results from the MD DNR survey confirmed anecdotal reports of significant loss of available oyster habitat within Maryland over the past 20 years. Only 2.37% of high quality oyster bottom habitat reported by the MBBS remained. Of the remaining high quality bottom habitat defined by the MBBS, 26.80% had degraded to low quality bottom classification consisting of heavily sedimented shell with sand or mud, and 68.58% was completely lost to sand or mud (Table 1). Overall, there was a 70.83% decline of available oyster habitat within the high quality bottom classifications defined by the MBBS. Given the significant level of habitat degradation on the high quality bottom since the MBBS was conducted, these population estimate calculations assumed a complete loss of low quality habitat identified in the MBBS survey.

MBBS-based adjustments to high quality habitat could not be applied to certain sections of Eastern Bay or the Choptank River (Broad Creek and Harris Creek) due to missing MBBS data. Therefore, oyster bar boundaries from the 1906-1911 Yates survey were used in these two areas (Table 1). The 1999-2000 MD DNR survey found that 85.88% of the charted oyster bars from the Yates survey have been lost to mud or sand. Only 0.95% of the charted Yates oyster bars fit the criteria for high quality and 11.29% meet low quality bottom habitat criteria. Habitat degradation of the charted Yates bars was more significant than the results based upon an assessment of the MBSS. This was expected given the earlier time period of the Yates Survey. These habitat adjustments were applied like the MBSS adjustments.

Significant amounts of fossil dredge shell and, to a lesser extent, shell retrieved from oyster packing houses have been planted in Maryland waters of Chesapeake Bay over the past four decades. The 1999-2000 MD DNR survey included an assessment of length of time after planting that fossil shell and shell from packing houses were suitable oyster habitat. Results indicated that shell plantings became moderately sedimented after an average of 5.5 years and heavily sedimented after an average of 18.6 years. For the purpose of estimating and assessing habitat for the biomass calculations, it was determined that only shell plantings 5 years old or less would be included in habitat estimates (Smith et al. 2005). Any GIS polygons designated as low quality habitat after adjustments to the MBBS and Yates surveys adjustments that overlapped shell plantings 5 years old or less were re-classified as high quality habitat. This did not result in an increase in the total amount of habitat, but simply changed the preliminary bottom classification. Total habitat only increased when the shell plantings did not overlap adjusted MBBS and Yates habitat polygons. These population estimates assumed that 1999-2003 shell plantings were constant throughout the 1994-2006 time period.

Leased oyster bottom within Maryland is another potential source of available habitat. Maryland leases are not legally permitted on chartered oyster bottom, and therefore, should not overlap any habitat derived from the MBBS and Yates surveys. Because leases cannot legally occur on natural oyster bottom, lease holders commonly plant shell substrate to enhance spat settlement and/or support seed plantings. Maryland lease harvest data were reviewed to determine the level of lease activity. The average lease harvest was 1,424 bushels between 2000 and 2004 and 3,767 bushels between 1990 and 1999 (MD DNR 2005). Based on this level of activity, it was assumed that there were little or no remaining shell plantings on lease grounds over the past 5 years. Given the functional longevity of shell plantings reported by MD DNR's Cooperative Oxford Laboratory, it was further assumed that any plantings prior to 2000 were no longer providing any suitable oyster habitat

Table 2 presents the sequence of calculated values in the estimation of available oyster habitat for the Choptank River basin. Table 3 presents a summary of estimates of high and low quality habitat for all eight basins delineated in this study.

Assumptions for Step 2

1. The degradation of oyster bottom habitat reported in the MD DNR 1999-2000 survey was representative of all bars within Maryland's portion of the Chesapeake Bay.
2. There was a complete loss of available oyster habitat within the "mud with cultch" and "sand" bottom classifications reported by the MBBS survey.
3. The amount of available high and low quality habitat remained constant throughout the 1994-2006 time series.
4. Shell plantings did not provide suitable oyster habitat after 5 years.
5. Shell plantings between 1999 and 2003 were representative of shell plantings between 1994 and 2006.
6. Maryland leases did not provide any suitable oyster bottom habitat between 1994 and 2006.

2.0 Step 3 – Estimation of Oyster Density in High and Low Quality Habitat

Oyster density estimates for high quality habitat

For high quality habitat, annual estimates of oyster density were calculated using data collected in the Maryland DNR Fall Oyster Dredge Survey (Fall Survey) and estimates of the average tow area and efficiency of the oyster dredge used to collect data in the survey. Maryland's Fall Survey collects oyster size and abundance data at 43 sentinel stations located throughout Maryland's Chesapeake Bay. Bushel count data collected in 2004 and basin averages are presented in Table 4. A detailed description of the survey can be found in the MD DNR report for the 2003 and 2004 Fall Surveys (MD DNR, 2005).

The Fall Survey provides number of oysters per bushel of dredged material. In order to convert number of oysters per bushel of dredge material to number of oysters per m², area swept to collect 1 bushel and the average efficiency of the dredge are needed. Historically, data on the Fall Survey dredge tow area was not recorded. Despite this limitation, the Fall Survey data are believed to be the best available data for estimating Maryland's oyster population. As a result, a study was conducted in 2001 and 2002 in order to study tow area of the dredge (unpublished MDNR study, 2002). The average tow area for collecting a one bushel sample of material was estimated to be 45 m². The original data used to develop this estimate are unavailable, so variance cannot be assigned.

A study of dredge efficiency initiated by MD DNR in 2001 estimated an average dredge efficiency of 10% (unpublished MDNR study, 2002).

Oyster density on high quality habitat (DH) for each basin was estimated as follows:

$$DH = ((BC_{\text{basin}} / (DE * TA)) * M) \quad (\text{Eqn 1})$$

where:

DH = oyster density (oysters/acre)

BC_{basin} = annual basin average of small and market oysters per bushel of cultch material (oysters/bushel), as determined from the Fall Survey.

DE = dredge efficiency (0.10)

TA = tow area/bu (45 m²/bu)

M = conversion factor (4046.856 m²/acre)

An example of estimating oyster density on high quality habitat for the Chester River in 2004 is provided below. Note that the mean bushel count (BC) is the only variable input parameter in these calculations.

$$\text{Oysters/acre} = ((59.5 \text{ oysters/bu}) / (0.10 * 45 \text{ m}^2/\text{bu})) * 4046.856 \text{ m}^2/\text{acre} = 53,509 \text{ oysters/acre (Eqn. 2)}$$

where:

BC = mean bushel count

Chester River Site 1: Buoy Rock = 34 oysters/bushel

Chester River Site 2: Old Field = 85 oysters/bushel

Mean bushel count = 59.5 oysters/bushel

Oyster density estimates for low quality habitat.

Density estimates used for low quality habitat are not well documented. For the period 1994-2001, the mean oyster density was 2.02 oysters/m² (personal communication, J. Vanisko).

The 2.02 oysters/m² density used is believed to be based upon data from a 1994 patent tong survey of areas categorized as "mud with cultch" and "sand with cultch" by the MBBS. The researchers could not recall the basis for the 2002 oyster density estimate of 0.36 oyster/m². We assumed that the reduction was made to account for the increase in disease mortality in recent years.

Examples of estimating oyster density on low quality habitat for the Chester River before/after 2002 are provided below.

$$\text{Before 2002: Oysters/acre} = 2.02 \text{ oysters/m}^2 * 4046.856 \text{ m}^2/\text{acre} = 8,175 \text{ oysters/acre (Eqn. 3)}$$

$$\text{2002 and later: Oysters/acre} = 0.36 \text{ oysters/m}^2 * 4046.856 \text{ m}^2/\text{acre} = 1,457 \text{ oysters/acre (Eqn. 4)}$$

Assumptions for Step 3

1. Oyster abundance statistics developed from data collected by the Fall Survey were representative of the associated basins.
2. The Fall Survey provided a representative sample of the classified high quality habitat within each basin, including natural oyster bars, managed oyster harvest reserves and oyster sanctuaries.
3. The defined dredge tow area for obtaining one bushel of sample material in the Fall Survey was representative and consistent throughout the 1994-2006 time series.
4. The defined dredge efficiency of the Fall Survey was representative and consistent throughout the 1994-2006 time series.
5. Oyster densities for low quality habitat derived from a 1994 patent tong survey were representative and consistent from 1994-2001.
6. Oyster density defined for low quality habitat in 2002 was representative and consistent from 2002-2006.

Step 4 – Calculation of the Estimated Oyster Population in MD Chesapeake Bay

The estimated density of oysters on high and low quality habitat for each basin was multiplied by the estimated amount of high and low quality habitat for each basin. Table 5 provides an example of the values in this calculation for the Chester River Basin in 2004.

The annual estimate of Maryland's Chesapeake Bay oyster population was calculated as the sum of the annual populations for Maryland's eight basins (Figure 2). Summary statistics from the 2004 calculation are presented in Table 6.

Step 5 – Calculation of Estimated Oyster Biomass in MD Chesapeake Bay

The estimated annual biomass of small and market oysters for each basin was calculated in two steps. In order to distribute the population among size groups, the total estimated number of small and market oysters for each basin was multiplied by the relative abundance within each 5-mm size class ≥ 35 mm. The relative abundance values were taken from the Fall Survey. This distribution was converted to biomass (grams per dry tissue weight) by the formula given below:

$$\log(10) \text{ weight} = -3.7595 + 2.062584 * \log(10) \text{ size class (Jordan et al. 2002)} \quad (\text{Eqn 5})$$

where:

size class = length in mm of the midpoint of a given size class

Total biomass was calculated as the sum of biomass in all size classes for all basins.

RESULTS

The 1994 baseline population of small and market sized oysters in Maryland Chesapeake Bay was estimated to be 582 million oysters. The biomass estimate was 1.1 million grams dry tissue weight. Maryland's oyster population remained at a relatively stable but historically low level of abundance between 1994 and 2000. In 2001, the population experienced another significant decline in response to increased disease mortality due to a multiple-year drought. The population has remained at this low level of abundance since 2001 (Figure 2, Table 7). Maryland's 2006 oyster population estimate is 59% (based on number of oysters) and 63% (based on oyster biomass) below the 1994 baseline period which was established for measuring progress towards achieving a 10-fold oyster population increase.

DISCUSSION

The methodology presented in this report was developed through an interstate research project jointly funded in 2000 by the National Oceanic and Atmospheric Administration Chesapeake Bay Stock Assessment Committee and U.S. Environmental Protection Agency Chesapeake Bay Program to estimate the 1994 baseline population of oysters and to develop a methodology for producing annual oyster population estimates that could be used to measure progress toward achieving the Chesapeake Bay 2000 Agreement goal of increasing the Chesapeake Bay oyster population 10-fold from its 1994 level of abundance by 2010. This methodology is based upon several critical assumptions that are identified in this report.

REFERENCES

- Jordan, Stephen J., Kelly N. Greenhawk, Carol B. McCollough, Jessica Vanisko, and Mark L. Homer. 2002. Oyster biomass, abundance and harvest in northern Chesapeake Bay: trends and forecasts. *Journal of Shellfish Research*, vol 21, no 2 p 733-741.
- MD DNR. 2005. Maryland Oyster Population Status Report 2003 and 2004 Fall Surveys. Maryland Department of Natural Resources, Fisheries Service. Annapolis, Maryland.
- Smith, G.F., D.G. Bruce, E.B. Roach, A. Hansen, R.I.E. Newell, and A.M. McManus. 2005. Assessment of Recent Habitat Conditions of Eastern Oyster *Crassostrea virginica* Bars in Mesohaline Chesapeake Bay. *North American Journal of Fisheries Management* 25:1569-1590.
- Smith, G.F., K.N. Greenhawk, D.G. Bruce, E.B. Roach, and S.J. Jordan. 2001 A Digital Presentation of the Maryland Oyster Habitat and Associated Bottom Types in the Chesapeake Bay (1974-1983). *Journal of Shellfish Research* 20:197-206.

Table 1. Relative abundance of high quality bottom classifications remaining from the values reported by the MBBS and Yates surveys, as reported by the 1999-2000 MD DNR Cooperative Oxford Laboratory acoustic benthic habitat survey.

| Habitat Classification | Bottom Classification | MBBS | Yates |
|------------------------|------------------------------------|-------------------|--------------|
| | | Percent Remaining | |
| High Quality Habitat | Clean shell | 1.16 | 0.62 |
| | Lightly sedimented shell with sand | 1.21 | 0.33 |
| | Total | 2.37 | 0.95 |
| Low Quality Habitat | Heavily sedimented shell with mud | 21.64 | 8.33 |
| | Heavily sedimented shell with sand | 5.16 | 2.96 |
| | Total | 26.80 | 11.29 |
| Total Habitat | Total | 29.17 | 12.24 |
| Habitat Lost | Mud | 20.58 | 32.52 |
| | Sand | 48.00 | 53.36 |
| | Total | 68.58 | 85.88 |
| Other | Gravel/cobble/boulder | 1.37 | 1.06 |
| | Unidentified | 0.88 | 0.82 |

Table 2. Values from the 2004 oyster habitat estimate for the Choptank River basin.

| Choptank River Basin | Bottom Class | Original Estimate | Adjustment Factor | Current Estimate of High Quality Habitat | Current Estimate of Low Quality Habitat |
|----------------------|---------------|-------------------|-------------------|--|---|
| MBBS | High Quality | 14,397.42 acres* | 2.37% | 340.18 acres | |
| | | | 26.80% | | 3,859.55 acres |
| | Low Quality | | 0% | | |
| Yates | Not specified | 1,028.43 acres* | 0.95% | 10.20 acres | |
| | | | 11.29% | | 115.68 acres |
| Shell Plantings | High Quality | 164.32 acres | 0 % | 164.32 acres | |
| Leased Bottom | Not specified | | 0% | | |

* Value may vary by a maximum of 1.12 acres due to differences in decimal precision in software employed.

Table 3. Acreages for each habitat type in each basin. Total acreage shown in bold text were used to calculate the annual Maryland oyster population from 1994 – 2006.

| <u>Basin</u> | <u>Habitat Type*</u> | <u>High Quality Habitat (ac)</u> | <u>Low Quality Habitat (ac)</u> | <u>Sum</u> |
|-----------------|----------------------|----------------------------------|---------------------------------|------------|
| CHESTER | B | 126.27 | 1,432.65 | 1,558.92 |
| | P | 481.23 | 0.00 | 481.23 |
| Total | | 607.50 | 1,432.65 | 2,040.15 |
| CHOPTANK | B | 340.18 | 3,859.55 | 4,199.73 |
| | P | 164.32 | 0.00 | 164.32 |
| | Y | 10.20 | 115.68 | 125.88 |
| Total | | 514.70 | 3,975.23 | 4,489.93 |
| EASTERN BAY | B | 76.28 | 865.48 | 941.76 |
| | P | 192.11 | 0.00 | 192.11 |
| | Y | 58.09 | 659.11 | 717.20 |
| Total | | 326.48 | 1,524.59 | 1,851.07 |
| LITTLE CHOPTANK | B | 79.36 | 900.41 | 979.77 |
| | P | 1.00 | 0.00 | 1.00 |
| Total | | 80.36 | 900.41 | 980.77 |
| MD MAINSTEM | B | 1,111.05 | 12,605.65 | 13,716.70 |
| | P | 684.33 | 0.00 | 684.33 |
| Total | | 1,795.38 | 12,605.65 | 14,401.03 |
| MD POTOMAC | B | 300.31 | 3,407.26 | 3,707.57 |
| | P | 12.86 | 0.00 | 12.86 |
| Total | | 313.17 | 3,407.26 | 3,720.43 |
| PATUXENT | B | 129.24 | 1,466.32 | 1,595.56 |
| | P | 56.25 | 0.00 | 56.25 |
| Total | | 185.49 | 1,466.32 | 1,651.81 |

| | | | | |
|--------------|---|---------------|-----------------|-----------|
| TANGIER | B | 563.15 | 6,389.26 | 6,952.41 |
| | P | 56.89 | 0.00 | 56.89 |
| Total | | 620.04 | 6,389.26 | 7,009.30 |
| Grand Total | | | | 36,144.49 |

* Habitat codes are as follows:

B = Maryland Bay Bottom Survey

P = shell plantings

Y = Yates survey

Table 4. Small- and market-size oysters per bushel of cultch material in samples collected by the 2004 MD DNR Fall Oyster Survey.

| Basin | Water Body | Oyster Bar | Site Code | Oysters/Bu | Basin Average | |
|-----------------|-------------------|-------------------|------------------|-------------------|----------------------|-------|
| Chester | Chester River | Buoy Rock | CHBR | 34 | 59.5 | |
| Chester | Chester River | Old Field | CHOF | 85 | | |
| Choptank | Broad Creek | Deep Neck | BCDN | 48 | 24.88 | |
| Choptank | Choptank River | Cook Point | CRCP | 1 | | |
| Choptank | Choptank River | Lighthouse | CRLI | 8 | | |
| Choptank | Choptank River | Oystershell Point | CROS | 20 | | |
| Choptank | Choptank River | Royston | CRRO | 43 | | |
| Choptank | Choptank River | Sandy Hill | CRSH | 20 | | |
| Choptank | Choptank River | Tilghman Wharf | CRTW | 39 | | |
| Choptank | Tred Avon River | Double Mills | TADM | 20 | | |
| Eastern Bay | Eastern Bay | Bugby | EBBU | 64 | | 80.83 |
| Eastern Bay | Eastern Bay | Hollicutts Noose | EBHN | 96 | | |
| Eastern Bay | Eastern Bay | Parsons Island | EBPI | 57 | | |
| Eastern Bay | Miles River | Bruffs Island | MRBI | 74 | | |
| Eastern Bay | Miles River | Long Point | MRLP | 42 | | |
| Eastern Bay | Miles River | Turtle Back | MRTU | 152 | | |
| Little Choptank | Little Choptank | Cason | LCCA | 27 | 22.50 | |
| Little Choptank | Little Choptank | Ragged Point | LCRP | 18 | | |
| Mainstem | Bay North | Swan Point | BNSP | 37 | 57.57 | |
| Mainstem | Middle Bay | Stone Rock | MESR | 54 | | |
| Mainstem | Upper Bay | Hacketts | UBHA | 76 | | |
| Mainstem | Western Shore | Butlers | WSBU | 123 | | |
| Mainstem | Western Shore | Flagpond | WSFP | 34 | | |
| Mainstem | Western Shore | Hog Island | WSHI | 74 | | |

| | | | | | | |
|----------|------------------|-------------------|------|-----|-------|-------|
| Mainstem | Western Shore | Holland Point | WSHP | 5 | | |
| Patuxent | Patuxent River | Broome Island | PXBI | 17 | 17.0 | |
| Potomac | Potomac River | Cornfield Harbor | PRCH | 54 | 56.57 | |
| Potomac | Potomac River | Lower Cedar Point | PRLC | 18 | | |
| Potomac | Potomac River | Ragged Point | PRRP | 0 | | |
| Potomac | St. Mary's River | Chickencock | SMCC | 67 | | |
| Potomac | St. Mary's River | Pagan | SMPA | 214 | | |
| Potomac | Wicomico River | Lancaster | WWLA | 27 | | |
| Potomac | Wicomico River | Mills West | WWMW | 16 | | |
| Tangier | Fishing Bay | Goose Creek | FBGC | 7 | | 75.20 |
| Tangier | Holland Straits | Holland Straits | HOHO | 151 | | |
| Tangier | Honga River | Normans | HRNO | 12 | | |
| Tangier | Manokin River | Georges | MAGE | 176 | | |
| Tangier | Nanticoke River | Wilson Shoal | NRWS | 33 | | |
| Tangier | Pocomoke Sound | Marumsco | PSMA | 106 | | |
| Tangier | Tangier Sound | Back Cove | TSBC | 120 | | |
| Tangier | Tangier Sound | Old Womans Leg | TSOW | 7 | | |
| Tangier | Tangier Sound | Piney Island | TSPI | 85 | | |
| Tangier | Tangier Sound | Sharkfin Shoal | TSSS | 55 | | |

Table 5. Habitat and density data for the 2004 Chester River oyster population estimate.

| Habitat Quality | Estimated Oyster Density (oysters/m ²) | Estimated Oyster Density (oysters/acre) | Estimated Available Habitat (acres) | Estimated Oyster Population (# of oysters) |
|-----------------|--|---|-------------------------------------|--|
| High | 59.50 | 53,508.65 | 607.50 | 32,506,371 |
| Low | 0.36 | 1,456.87 | 1432.65 | 2,087,182 |

Table 6. Input data and estimates of the 2004 oyster population by basin and for all basins combined.

| | | Chester | Eastern Bay | Choptank | Little Choptank | Tangier Sound | Potomac | Patuxent | Mainstem | Total |
|--|-------------------------|---------|-------------|----------|-----------------|---------------|---------|----------|----------|-----------|
| Habitat (ac) | | | | | | | | | | |
| High quality | | | | | | | | | | |
| | MBBS | 126.27 | 76.28 | 340.18 | 79.36 | 563.15 | 300.31 | 129.24 | 1111.05 | 2,725.84 |
| | Yates | 0 | 58.09 | 10.20 | 0 | 0 | 0 | 0 | 0 | 68.29 |
| | Shell plantings | 481.23 | 192.11 | 164.32 | 1.00 | 56.89 | 12.86 | 56.25 | 684.33 | 1,648.99 |
| | Leased bottom | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Total | 607.5 | 326.48 | 514.70 | 80.36 | 620.04 | 313.17 | 185.49 | 1795.38 | 4443.12 |
| Low quality | | | | | | | | | | |
| | MBBS | 1432.65 | 865.48 | 3859.55 | 900.41 | 6389.26 | 3407.26 | 1466.32 | 12605.65 | 30,926.58 |
| | Yates | 0 | 659.11 | 115.68 | 0 | 0 | 0 | 0 | 0 | 774.79 |
| | Shell plantings | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| | Leased bottom | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| | Total | 1432.65 | 1524.59 | 3975.23 | 900.41 | 6389.26 | 3407.26 | 1466.32 | 12605.65 | 31701.37 |
| Density (oysters/m²) | | | | | | | | | | |
| | on high quality habitat | 13.22 | 17.96 | 5.53 | 5.00 | 16.71 | 12.57 | 3.78 | 12.79 | |

| | | | | | | | | | | |
|-------------------------------|-------------------------|---------------|---------------|---------------|--------------|---------------|---------------|--------------|----------------|----------------|
| | on low quality habitat | .36 | .36 | .36 | .36 | .36 | .36 | .36 | .36 | |
| Population | | | | | | | | | | |
| | on high quality habitat | 32,506,370.82 | 23,732,981.86 | 11,513,901.11 | 1,626,026.74 | 41,931,730.46 | 15,932,448.95 | 2,835,793.87 | 92,954,116.29 | 223,033,370.10 |
| | on low quality habitat | 2,087,182.17 | 2,221,126.63 | 5,791,386.02 | 1,311,778.66 | 9,308,309.46 | 4,963,928.61 | 2,136,234.92 | 18,364,770.12 | 46,184,716.58 |
| Total Basin Population | | 34,593,552.99 | 25,954,108.49 | 17,305,287.12 | 2,937,805.40 | 51,240,039.92 | 20,896,377.55 | 4,972,028.79 | 111,318,886.41 | 269,218,086.68 |

Table 7. Maryland Oyster Population Estimates by Basin (1994-2006)

| | Chester | Eastern Bay | Choptank | Little Choptank | Tangier Sound | Potomac | Patuxent | Mainstem | Total |
|-------------|----------------|--------------------|-----------------|------------------------|----------------------|----------------|-----------------|-----------------|----------------|
| 1994 | 60,061,223.12 | 24,892,220.04 | 97,124,389.50 | 16,683,089.13 | 139,327,515.80 | 64,586,301.14 | 13,321,142.73 | 173,281,327.49 | 589,277,208.95 |
| 1995 | 58,422,246.44 | 21,662,577.15 | 89,082,016.36 | 20,657,821.16 | 102,023,888.56 | 72,673,225.99 | 17,491,427.84 | 180,316,316.44 | 562,329,519.95 |
| 1996 | 64,158,664.82 | 30,911,099.97 | 93,479,285.13 | 18,309,115.87 | 86,578,291.04 | 55,050,971.85 | 18,492,296.27 | 206,611,029.24 | 573,590,754.19 |
| 1997 | 101,581,965.68 | 29,002,674.63 | 77,163,103.66 | 14,225,982.05 | 83,176,913.97 | 63,017,196.32 | 18,992,730.48 | 163,939,784.79 | 551,100,351.58 |
| 1998 | 108,684,197.96 | 116,790,240.52 | 118,127,133.73 | 17,911,642.67 | 78,548,810.74 | 83,254,625.16 | 17,658,239.24 | 181,238,937.94 | 722,213,827.97 |
| 1999 | 109,503,686.30 | 70,596,560.36 | 106,555,373.82 | 16,827,624.84 | 95,165,374.14 | 51,751,828.38 | 18,492,296.27 | 192,771,706.71 | 661,664,450.83 |
| 2000 | 107,318,384.06 | 53,371,798.27 | 88,677,004.77 | 17,803,240.88 | 87,191,654.12 | 45,394,942.19 | 24,831,129.63 | 166,015,683.17 | 590,603,837.08 |
| 2001 | 51,802,808.13 | 23,018,069.50 | 38,770,901.75 | 5,647,849.97 | 36,184,764.36 | 15,987,895.81 | 3,971,160.37 | 81,794,998.36 | 257,178,448.23 |
| 2002 | 23,667,041.79 | 15,433,302.10 | 14,701,641.14 | 1,528,582.23 | 43,210,559.62 | 9,711,476.53 | 4,471,594.58 | 70,031,574.21 | 182,755,772.20 |
| 2003 | 39,510,483.03 | 18,809,746.94 | 13,775,900.35 | 2,287,394.70 | 86,759,338.20 | 38,478,398.24 | 10,977,239.35 | 63,573,223.70 | 274,171,724.51 |
| 2004 | 34,593,552.99 | 25,954,108.49 | 17,305,287.12 | 2,937,805.40 | 51,240,039.92 | 20,896,377.55 | 4,972,028.79 | 111,318,886.41 | 269,218,086.68 |
| 2005 | 33,774,064.65 | 25,905,174.51 | 30,265,658.22 | 3,226,876.82 | 29,437,770.49 | 14,096,923.33 | 7,307,388.45 | 141,996,051.34 | 286,009,907.81 |
| 2006 | 36,778,855.23 | 19,690,558.64 | 30,902,105.01 | 4,419,296.43 | 23,415,660.27 | 15,746,495.07 | 9,475,936.71 | 96,556,942.38 | 236,985,849.74 |

Table 8. Maryland Oyster Biomass Estimates by Basin (1994-2006)

| | Chester | Eastern Bay | Choptank | Little Choptank | TangierSound | Potomac | Patuxent | Mainstem | Total |
|-------------|----------------|--------------------|-----------------|------------------------|---------------------|----------------|-----------------|-----------------|---------------|
| 1994 | 100,604,202 | 42,492,811 | 89,003,017 | 58,934,152 | 114,605,118 | 33,526,888 | 27,440,701 | 609,560,420 | 1,076,167,309 |
| 1995 | 83,018,760 | 36,564,486 | 107,353,882 | 21,951,048 | 115,619,792 | 74,609,509 | 30,259,873 | 272,420,755 | 741,798,106 |
| 1996 | 84,401,894 | 33,925,831 | 118,572,292 | 21,735,808 | 112,748,480 | 68,748,028 | 24,039,045 | 259,758,883 | 723,930,262 |
| 1997 | 116,205,172 | 40,108,786 | 113,908,877 | 19,006,144 | 123,478,596 | 86,500,348 | 38,790,667 | 271,806,654 | 809,805,244 |
| 1998 | 126,640,415 | 108,261,204 | 127,855,090 | 23,603,996 | 120,699,104 | 97,199,120 | 40,131,468 | 239,120,894 | 883,511,292 |
| 1999 | 168,252,067 | 84,869,759 | 138,946,576 | 21,301,739 | 122,908,065 | 60,204,250 | 48,857,214 | 283,108,217 | 928,447,887 |
| 2000 | 151,286,084 | 77,214,467 | 132,172,207 | 20,039,015 | 80,800,250 | 50,947,144 | 33,327,946 | 272,765,205 | 818,552,319 |
| 2001 | 83,792,975 | 36,420,227 | 57,078,960 | 6,776,996 | 41,429,392 | 20,935,500 | 5,434,841 | 110,525,021 | 362,393,912 |
| 2002 | 46,141,627 | 21,617,947 | 26,480,472 | 2,000,058 | 40,872,830 | 15,292,220 | 8,809,732 | 109,005,021 | 270,220,086 |
| 2003 | 68,103,750 | 26,515,210 | 20,581,190 | 1,493,491 | 76,380,446 | 22,077,915 | 8,655,666 | 74,869,463 | 298,677,131 |
| 2004 | 51,407,440 | 34,581,494 | 26,750,684 | 2,964,143 | 64,446,800 | 19,018,801 | 6,208,066 | 159,467,442 | 364,844,871 |
| 2005 | 62,418,078 | 36,978,617 | 41,481,615 | 4,224,597 | 47,944,247 | 23,701,085 | 13,173,224 | 207,064,520 | 436,985,983 |
| 2006 | 81,982,665 | 33,809,416 | 48,614,516 | 5,899,524 | 41,059,409 | 29,945,828 | 13,712,229 | 144,731,010 | 399,754,597 |




Figure 1. Map showing the 8 Maryland basins used in oyster population estimates.




Figure 2. Maryland Chesapeake Bay Oyster Population Estimates (1994-2006)

Appendix C. Development of Starting Population (Barker 2007)

**Spatially Distributed Estimate
of the 2004 Maryland Chesapeake Bay
Population of *Crassostrea virginica***

**Linda S. Barker
Maryland DNR Fisheries Service
March 14, 2007**

Introduction

The Maryland Department of Natural Resources (MdDNR), the Army Corps of Engineers (Norfolk district) and the Virginia Marine Resources Commission (VMRC) are the lead agencies in preparing an Environmental Impact Statement (EIS) to evaluate several native and non-native oyster restoration alternatives for Chesapeake Bay. The overall purpose of the EIS is to evaluate alternatives that may restore the ecological and economic benefits of the oyster resource in Chesapeake Bay.

The University of Maryland and Versar, Inc. have been tasked with modeling to support the ecological risk assessment component of the EIS. The model will be used to estimate the probability of achieving the stated goal of re-establishing an oyster population comparable to that observed during the 1920 - 1970 time period. The model is intended to be both spatially and temporally explicit, since several parameters vary over space or time. MdDNR has agreed to supply a spatially distributed estimate of the 2004 population of oysters in Maryland's portion of Chesapeake Bay as the starting point for the model.

This document describes the methods used to determine the spatial distribution and site-specific size composition of the estimated 2004 Maryland Chesapeake Bay oyster population. The starting point for this exercise is the estimate of the 2004 population of "small"- and "market"-size oysters for each of eight basins. The final product was a file of total oyster population (including spat), distributed by 5-mm groups on the bars in the Maryland portion of Chesapeake Bay in 2004 (a matrix of 2115 bars and 28 5-mm height groups).

Methods

The spatially- and size-distributed 2004 Maryland Chesapeake Bay oyster population was calculated in three steps:

Estimation of the 2004 Maryland Chesapeake Bay population of "small-" and "market-"size oysters as reported by MdDNR for the Chesapeake Bay Oyster Population Estimation (Greenhawk and O'Connell 2007).

Expansion of the 2004 Maryland Chesapeake Bay oyster population to include spat, apportioned among NOAA regions.

Distribution of each NOAA region's total oyster population among its bars by 5-mm height groups.

Check of the final answer file consisted of mathematical examination of the values and patterns in the final matrix and subjective consideration of the spatial pattern of population density.

Estimation of the number of "small-" and "market-"size oysters on each oyster bar in Maryland's portion of Chesapeake Bay in 2004 .

The estimated 2004 Maryland Chesapeake Bay "small-" and "market-" size oyster population was supplied by MdDNR Fisheries Service staff. Methods used to develop this estimate are found in Greenhawk and O'Connell (2007).

The basin-specific oyster population and available habitat estimates presented in Greenhawk and O'Connell (2007) were used to calculate oyster density (oysters/acre) as follows. The oyster population in each basin was divided by the amount of available oyster habitat in that basin. The mean oyster density in each basin was then multiplied by each bar area to produce a file of the "small-" and "market-" size oyster population distributed among bars.

Expansion of the 2004 Maryland Chesapeake Bay oyster population to include spat, apportioned among NOAA regions.

Development of representative height frequencies for each NOAA region

The annual MdDNR Fall Oyster Survey samples approximately 400 oyster bars in Maryland's portion of Chesapeake Bay from October through December of each year (Tarnowski, 2005). A subset of these, the "Modified Fall Survey" (MFS) sites, have been deemed representative of the Bay population. Additional measurements are taken at these sites, including oyster counts by 5-mm size group and counts of spat.

One half of the NOAA regions had at least one appropriate MFS site. For those NOAA regions without a MFS site, the closest MFS site in a similar environmental setting was assigned to provide the representative height frequency. For NOAA regions with more than one MFS site, the pooled absolute counts for each height class were used to develop the representative height-frequency distribution.

It should be noted that the NOAA region coding used in this analysis is the "new" system, provided by the NOAA Northeast Fisheries Science Center. This system was used because it encompasses both Maryland and Virginia, enabling the modelers to use a consistent coding system in later work.

Estimation of total population (including spat) for each bar

Since the population values supplied by Greenhawk and O'Connell (2007) included only "small- and market"-size oysters, the next step in the analysis was to "scale up" these values to include spat. A population multiplier was developed for each NOAA region to calculate the total population (including spat). The 5-mm height frequency data for each NOAA region specifically identified spat. The 5-mm height-class data were collapsed into two groups (small <75 mm and market >75 mm) to produce a height frequency of spat, small- and market-size (SSM) oysters. Population multipliers were developed at the NOAA-region scale and were then applied to produce total population for each bar within the NOAA regions as follows:

$$\text{Population Multiplier} = \frac{\text{Spat Count} + \text{Small Count} + \text{Market Count}}{\text{Small Count} + \text{Market Count}} \quad \text{Eqn 1.}$$

$$\text{Total Population} = \text{Population Multiplier} * \text{Small \& Market Population} \quad \text{Eqn 2.}$$

Distribution of each NOAA region's total population among its bars by 5-mm height groups.

The absolute frequency of each bar's total population was a straightforward multiple of the absolute population and the appropriate relative height frequency. For every NOAA region, the associated MFS site provided counts of SSM oysters for each 5-mm height group. Each 5-mm height-group proportion was calculated by dividing the count of each height group by the total count. The final product was a file of total oyster population (including spat), distributed by 5-mm height groups on the oyster bars in Maryland's portion of Chesapeake Bay in 2004 (a matrix of 2115 bars and 28 5-mm height groups).

Checks.

Two checks were performed on the final distributed bar population. A cursory visual check was made to verify that the height-frequency distribution pattern was consistent throughout each NOAA region. Additionally, the "original" SSM population value used to develop the distributed population was compared to the sum of the 5-mm counts for each bar. The percent difference between these values was calculated. All percent differences were less than 2%, which was consistent with rounding error.

RESULTS AND DISCUSSION

Allocation of the basin populations among bars.

One source of error occurred during the distribution of the basin-scale "small"- and "market"-size population among bars. Since the total area of each basin was known, each bar could be assigned a portion of the basin population equal to its proportion of total basin area. However, this method neglected

to consider habitat quality. Because the relative proportion of high and low quality habitat was known only at the basin scale, there were no means to allocate the quality ratings among bars within a basin.

Representative height frequencies

Between one and seven MFS sites were used to develop a representative height frequency for each NOAA region. For almost half of the 38 NOAA regions, there was no MFS site within the NOAA region boundaries (Figure 1, Table 1) so the closest MFS site in a similar environmental setting was assigned to provide the representative height frequency. In other NOAA regions where more than one MFS site was within the boundaries (Table 1), the representative height frequency was developed from the sum of counts in all height classes (Table 2). Although these are the best available data, the extent to which these height frequencies are representative of oysters within each bar in a given region is not known.

Graphical representation of the MFS height frequencies used with the most heavily populated NOAA regions are presented in Figures 2-4. These histograms not only demonstrate the variability among the MFS height frequencies, but also the variability of sample sizes.

2004 oyster population.

The 2004 Maryland Chesapeake Bay total population of small and market oysters was 269,218,086 (Greenhawk and O'Connell, 2007). This number increases to 291,751,774 oysters when spat are included. In this report, this population was distributed among 38 NOAA regions with 1 to 273 bars per region, for a final matrix of 28 5-mm height groups x 2115 bars. Table 3 provides the basic statistics on the bar populations and the aggregated NOAA populations.

Although this report provides the spatial distribution and the size composition of the 2004 oyster population in the Maryland portion of Chesapeake Bay, including spat, there are two limitations of note:

- 1.) The spatial variability portrayed was limited to the scale of each NOAA region.
- 2.) Total population estimates and height-frequency distributions were developed without variance estimates. Without the ability to assign measures of variance to these values, no associated measures of precision could be developed for our estimates.

ACKNOWLEDGEMENTS

This analysis could not have been performed without the technical expertise and support of Lisa Warner and Kelly Greenhawk of MdDNR. Thanks also to Mitchell Tarnowski and Luke Whitman of MdDNR for review and comments.

References

Dew, J. R., Berkson, J. M., Hallerman, E. M., and Allen, S. K. Jr. 2003. A model for assessing the likelihood of self-sustaining populations resulting from commercial production of triploid Suminoe oysters (*Crassostrea ariakensis*) in Chesapeake Bay. Fishery Bulletin 101:758-768, referenced in (Draft

July 9, 2004) Integrated Approach For The Demographic Model Used In The Ecological Risk Assessment by University of Maryland, College Park, MD and Versar, Inc., Columbia, MD)

Dew, J.R.. 2002. A population dynamic model assessing options for managing eastern oysters (*Crassostrea virginica*) and triploid Suminoe oysters (*Crassostrea ariakensis*) in Chesapeake Bay. M.S. Thesis, pg 127. Virginia Polytechnic Institute and State University, Blacksburg, VA., referenced in (Draft July 9, 2004) Integrated Approach For The Demographic Model Used In The Ecological Risk Assessment by University of Maryland, College Park, MD and Versar, Inc., Columbia, MD)

Greenhawk, K. and O'Connell, T. 2007. Oyster population estimates for the Maryland Portion of Chesapeake Bay 1994 – 2006. Maryland DNR.

Tarnowski, M.(ed.) 2005. Maryland Oyster Population Status Report for the 2003 and 2004 fall surveys. Maryland DNR.

Table 1. Assignment of Modified Fall Survey (MFS) sites within NOAA regions

| NOAA | ASSOCIATED MFS SITE(S) | TITLE* |
|------|--|--------|
| 212 | NONE - USE MAGE | |
| 217 | UBHA, WSHP, MESR, WSFP | 4A |
| 218 | BNSP | |
| 220 | WSHI, WSBU, HOHO | 2E |
| 224 | BCDN, TADM, CRTW, CRRO, CRCP, CRLI, CRSH | 3B |
| 225 | CROS | |
| 226 | EBHN, EBPI, EBBU, MRBI | 3A |
| 228 | FBGC | |
| 230 | NONE - USE WSHP | |
| 231 | HRNO | |
| 232 | NONE - USE TSSS | |
| 235 | LCRP, LCCA | 2A |
| 236 | NONE - USE UBHA | |
| 237 | MAGE | |
| 239 | MRLP, MRTU | 2B |
| 243 | CHBR, CHOF | 1A |
| 245 | NRWS | |
| 246 | NONE - USE BNSP | |
| 248 | PXBI | |
| 249 | NONE - USE PXBI | |
| 251 | PSMA | |
| 252 | PRCH | |
| 253 | PRLC | |
| 254 | NONE - USE PRLC | |
| 257 | NONE - USE WWLA | |
| 258 | WWMW, WWLA | 2F |
| 268 | NONE - USE UBHA | |
| 270 | NONE - USE SMCC | |
| 271 | NONE - USE UBHA | |
| 273 | SMPA, SMCC | 2C |
| 276 | TSSS, TSPI, TSOW, TSBC | 4C |
| 279 | NONE - USE NRWS | |
| 308 | NONE - USE WSBU | |
| 346 | NONE - USE PSMA | |
| 358 | NONE - USE TSOW | |
| 380 | USE PRCH | |
| 381 | USE PRLC | |
| 382 | NONE - USE PRLC | |

* Where more than one MFS was assigned to a NOAA region, a title was assigned to the group.

Table 2. NOAA regions, associated MFS sites, spat, small- and market-size counts from the 2004 MdDNR Fall Oyster Survey, and population multiplier used to estimate total population (including spat).

| NOAA code | MFS code | Spat Count | Small Count | Market Count | Population Multiplier |
|------------------|-----------------|-------------------|--------------------|---------------------|------------------------------|
| 212 | MAGE | 4 | 149 | 27 | 1.02 |
| 217 | 4A | 0 | 106 | 63 | 1.00 |
| 218 | BNSP | 0 | 3 | 34 | 1.00 |
| 220 | 2E | 1 | 117 | 231 | 1.00 |
| 224 | 3B | 206 | 92 | 87 | 2.15 |
| 225 | CROS | 0 | 1 | 19 | 1.00 |
| 226 | 3A | 0 | 149 | 142 | 1.00 |
| 228 | FBGC | 0 | 1 | 6 | 1.00 |
| 230 | WSHP | 0 | 2 | 3 | 1.00 |
| 231 | HRNO | 0 | 6 | 6 | 1.00 |
| 232 | TSSS | 2 | 36 | 19 | 1.04 |
| 235 | 2A | 7 | 41 | 4 | 1.16 |
| 236 | UBHA | 0 | 38 | 38 | 1.00 |
| 237 | MAGE | 4 | 149 | 27 | 1.02 |
| 239 | 2B | 1 | 126 | 68 | 1.01 |
| 243 | 1A | 1 | 64 | 55 | 1.01 |
| 245 | NRWS | 0 | 17 | 16 | 1.00 |
| 246 | BNSP | 0 | 3 | 34 | 1.00 |
| 248 | PXBI | 0 | 12 | 5 | 1.00 |
| 249 | PXBI | 0 | 12 | 5 | 1.00 |
| 251 | PSMA | 14 | 55 | 51 | 1.13 |
| 252 | PRCH | 0 | 32 | 22 | 1.00 |
| 253 | PRLC | 0 | 6 | 12 | 1.00 |
| 254 | PRLC | 0 | 6 | 12 | 1.00 |
| 257 | WWLA | 0 | 6 | 21 | 1.00 |
| 258 | 2F | 0 | 14 | 29 | 1.00 |
| 268 | UBHA | 0 | 38 | 38 | 1.00 |
| 270 | SMCC | 2 | 61 | 6 | 1.03 |
| 271 | UBHA | 0 | 38 | 38 | 1.00 |
| 273 | 2C | 8 | 274 | 7 | 1.03 |
| 276 | 4C | 109 | 188 | 79 | 1.41 |
| 279 | NRWS | 0 | 17 | 16 | 1.00 |
| 308 | WSBU | 0 | 58 | 65 | 1.00 |
| 346 | PSMA | 14 | 55 | 51 | 1.13 |
| 358 | TSOW | 29 | 3 | 4 | 5.14 |
| 380 | PRCH | 0 | 32 | 22 | 1.00 |
| 381 | PRCH | 0 | 32 | 22 | 1.00 |
| 382 | PRLC | 0 | 6 | 12 | 1.00 |

Table 3. Basic statistics on 2004 Maryland Chesapeake Bay oyster population, by NOAA region and by bar.

| | NOAA | bar |
|---------------|-------------|------------|
| mean | 7,677,678 | 137,944 |
| median | 2,839,501 | 25,759 |
| min | 18,678 | 461 |
| max | 41,026,637 | 8,838,600 |
| SE | 1,792,304 | 8,688 |

Figure 1. NOAA regions and locations of assigned MFS height frequency sites.




Figure 2. Height frequency for NOAA 217, population 4.10×10^7 , representing 14% of the total Maryland Chesapeake Bay 2004 oyster population. Developed from length group totals of MFS sites UBHA, WSHP, MESR, WSFP (total $n = 169$).




Figure 3. Height frequency for NOAA 218 and 246, total population 3.65×10^7 , representing 13% of the total Maryland Chesapeake Bay 2004 oyster population. Developed from MFS site BNSP ($n = 37$).




Figure 4. Height frequency for NOAA 235, total population 3.55×10^7 , representing 12% of the total Maryland Chesapeake Bay 2004 oyster population. Developed from length group totals of MFS sites LCRP and LCCA (n = 52).





Figure 5. Maryland Chesapeake Bay 2004 Estimated Oyster Population Density by NOAA Region.



Appendix D. Estimation of annual mortality rates for eastern oysters (*Crassostrea virginica*) in Chesapeake Bay based on box counts and application of those rates to project population growth of *C. virginica* and *C. ariakensis* (Vølstad, J. H., J. Dew, and M. Tarnowski, *Journal of Shellfish Management*, In Press)

**Estimation of annual mortality rates for eastern oysters (*Crassostrea virginica*)
in Chesapeake Bay based on box counts and application of those rates to project population growth of *C. virginica* and *C. ariakensis***

by

Jon H. Vølstad¹, Jodi Dew¹, and Mitchell Tarnowski²

¹Versar, Inc. 9200 Rumsey Road,

Columbia, MD 21045, USA.

² Maryland Dept. of Natural Resources, Tawes State Office Building, B-2,

Annapolis, MD 21401, USA.

Keywords: natural mortality, *Crassostrea virginica*, *Crassostrea ariakensis*, Chesapeake Bay, population model, ecological risk assessment

Abstract

In an effort to restore the ecological role of oysters in Chesapeake Bay and the economic benefits of a commercial fishery, the states of Maryland and Virginia are considering introducing the non-native Asian oyster (*Crassostrea ariakensis*) into the Bay. As part of an ecological risk assessment (ERA) to evaluate the proposed action and alternatives, demographic modeling is being employed to project the change in populations of both the Asian and the native eastern oyster (*C. virginica*) in the Bay across space and time. Annual mortality rates are vital input to the demographic model. We present two approaches for parameterizing mortality rates for *C. virginica* by salinity ranges and disease-intensity categories and discuss how these rates could be applied to project population growth for the Asian oyster. We estimated mortality rates from empirical data collected during annual dredge surveys of oyster beds in Maryland. We compared counts of recent boxes (dead oysters without fouling or sedimentation on the inner valve surfaces, including “gapers” of one or two weeks old with tissue remaining in the shell), old boxes (dead oysters without tissue remnants but with fouling, sedimentation or both on the inner valve surfaces), and live oysters in market-size and small classes. Our mortality estimates based on counts of recent boxes consistently differentiated between years with high disease intensity and those with low disease intensity, between wet and dry years, and between salinity zones. In contrast, traditional estimates of yearly mortality based on total box counts often were out of phase with measured levels of disease intensity and weather (dry or wet). To model populations of *C. ariakensis*, we propose to adjust the mortality rates for *C. virginica* based on research results that provide estimates of differences between the two species’ resistance to MSX and dermo and to other mortality factors, such as predation.

Introduction

The eastern oyster (*Crassostrea virginica*) historically supported a valued fishery and formed an important component of the Chesapeake Bay’s ecosystem. In recent years, the abundance of eastern oyster in the Chesapeake Bay has declined to less than 1% of estimated virgin stock due to intense fishing pressure during the 19th and 20th centuries, habitat destruction, degraded water quality, and disease (NRC 2004; Gottlieb and Schweighofer 1996).

The states of Maryland and Virginia recognized the need to reverse the decline in oyster stocks to restore the ecological role of oysters in the Bay and the economic benefits of a commercial fishery. To achieve those objectives, managers in Maryland and Virginia have proposed options for increasing the biomass of oyster stocks (NRC 2004), including the deliberate release of diploid Asian oysters (*C. ariakensis*) into Chesapeake Bay on a large scale to establish a self-recruiting population.

A comprehensive study of the potential effects of introducing the non-native Asian oyster into Chesapeake Bay is being conducted to support a programmatic Environmental Impact Statement (EIS) that will evaluate alternative restoration strategies. The EIS will address the proposed action to introduce diploid *C. ariakensis* and seven restoration alternatives, which include stocking and aquaculture of the Asian oyster and of the native eastern oyster. The comprehensive study, known as an ecological risk assessment (ERA), is being conducted as one element of the overall EIS. The ERA will identify the ecological risks and benefits posed by each of the eight restoration alternatives.

Evaluating the ecological effects of the proposed restoration alternatives requires constructing a demographic model that can project the change in populations of Asian and eastern oyster in the Bay that might result from implementing each alternative. The model must account for the primary population dynamics: growth, stock recruitment, and mortality. The reliability of model projections depends on the validity of the various input parameters, including mortality. Accurate and precise estimates of annual natural mortality rates (M) for Asian and eastern oyster are critical to the performance of the demographic model and the reliability of conclusions to be drawn from the ERA and the EIS.

The accuracy and precision of estimates of mortality are determined by the quantity and quality of the data and the validity of methods used to calculate the estimates. The Maryland Department of Natural Resources (DNR) collects empirical data through annual surveys of oyster beds in Maryland. Samples of oysters are collected by dredging at representatively selected oyster bars during fall. Oysters from each sample are sorted by size and classified into one of two mortality categories: live oyster or “box.” The box category refers to dead,

articulated shells. DNR classifies boxes further as “recent” (open shells with tissue remaining inside them, known as “gapers,” and empty shells without fouling or sedimentation on the inner valve surfaces) or “old” (empty shells with fouling, sedimentation or both on the inner valve surfaces). DNR estimates annual mortality as the proportion of boxes to live oysters (Tarnowski 2003). Natural mortality of oysters in Delaware Bay and Virginia is also estimated from annual collections of live oysters and boxes (Southworth et al. 2005; Ford et al. 2006). Ford et al. (2006) found that mortality rates obtained from counts of recent boxes and rates based on total box counts both provided reliable indices of total mortality for the year prior to the survey.

Here we describe a method for estimating annual mortality based solely on counts of recent boxes of small (shell height from 40 to 76 mm) and market-size (shell height \geq 76 mm) oysters. We compare estimated mortality rates for Chesapeake Bay based on counts of recent boxes with estimates based on total box counts (Jordan et al. 2002; Jordan and Coakley 2004). We estimate oyster mortality rates for specific ranges of salinity and disease intensity to enhance their applicability for spatially explicit modeling of oyster population dynamics, and we discuss how the rates for *C. virginica* could be adapted to model population growth for *C. ariakensis*.

Material and methods

We analyzed data from Maryland DNR’s annual fall survey of oysters for the period from 1980 to 2005. The survey is conducted mostly during October (early November in some years) using a standard oyster dredge to collect samples of live oysters and boxes from 200 to 400 representative oyster bars (Tarnowski 2003). A fixed set of 43 “disease bars” has been sampled every year since 1980; length frequency data and tissue samples from each size class (for disease studies) have been collected annually from these bars since 1990. A composite sample of one Maryland bushel ($\sim 0.046 \text{ m}^3$) is collected at each of the disease bars by pooling two, $\frac{1}{2}$ -bushel subsamples from replicate tows, and at each bar in seed-production areas by pooling five, $\frac{1}{5}$ -bushel subsamples from replicate tows. At all other bars, DNR collects a $\frac{1}{2}$ -bushel sample from a single tow. DNR has reported counts of recent and old boxes-per-bushel separately for small and market-size oysters since 1991 and for spat (age 0) since 1992. The counts of live oysters and boxes of age 1+ in each sample is classified into two size categories (shell

height; market: ≥ 76 mm; small: < 75 mm). Spat is identified morphologically by their asymmetric valves. Spat have one valve that is thinner and narrower than the other (the lower valve if they are oriented parallel to the substrate; M. Tarnowski, MD DNR personal communication). All counts are standardized to a volume of one bushel. We estimated mortality rates using only the counts of small and market-size boxes. Box counts for spat are considered to be unreliable because boxes can break apart easily during collection.

We made the following assumptions to estimate mortality from counts of recent boxes: (1) time since death (TSD) is between one and two weeks; (2) the instantaneous (1-week) mortality rate is constant from June to October; and (3) the cumulative natural mortality from October through May is negligible. Data from an intense mortality study conducted by Maryland DNR in the Choptank River during 2002 provide some support for assumptions 1 and 2. The Choptank study was conducted when freshwater run-off into the Bay was low to estimate the extent to which mortality might increase because of increased *Haplosporidium nelsoni* (MSX disease) infection. MSX is likely to cause additional mortality during years in which salinity is high due to drought. Maryland DNR counted live oysters and boxes from one-bushel samples at each of six oyster bars located in waters that normally are moderately salty every month from June through August, and during the annual fall oyster survey in October 2002. DNR's empirical data from the study in the Choptank River during 2002 (results summarized later in this paper) support assumptions (1) and (2). Assumption (3) is supported by numerous studies showing that natural mortality from *Perkinsus marinus* (dermo disease) and *H. nelsoni* (MSX) occurs primarily from early summer to October (Andrews 1996; Burreson and Ragone Calvo 1996; Ford and Tripp 1996; Ford and Haskin 1982) and that predation on oysters is greatest during summer months (Carriker 1955; Manzi 1970; Gunter 1979; Garton and Stickle 1980; Pearse and Wharton 1938; Landers and Rhodes 1970). When using total box counts to estimate mortality rates, we followed Jordan et al. (2002) and assumed that the TSD of a box is up to one year. According to that assumption, all boxes observed in the fall survey died within one year prior to the survey, and shells from oysters that died during the year remained articulated until the fall.

To refine the parameters for use in our demographic model, we estimated mortality rates for specific levels of salinity and disease intensity. Dermo infections in Chesapeake Bay are heaviest and cause most mortality at

medium and high salinities (NRC 2004). We calculated average annual mortality rates by post-stratifying the counts of live oysters and boxes from all oyster bars across years by salinity class and disease tier. The mean salinity from May through September for each bar by year was estimated from the nearest Maryland Chesapeake Bay Monitoring Program station, or interpolated by kriging. We classified the yearly observations of live oysters and boxes from each bar into three salinity (S) classes (low: $S < 11$ ppt; medium: $11 \leq S < 15$ ppt; and high: $S \geq 15$ ppt). The salinity thresholds were defined in consultation with Chesapeake Bay oyster biologists. Maryland DNR uses an index of disease intensity ranging from one to seven based on pathogen concentration in hemolymph or solid tissue (see Giesecker 2001) to classify dermo disease-intensity into three tiers (Tarnowski 2003). Tier 1 (dermo intensity > 2.85) and Tier 2 ($2 < \text{dermo intensity} \leq 2.85$) represent years with high and medium disease intensity, respectively; Tier 3 (dermo intensity ≤ 2) represents years with relatively low disease intensity (Figure 1). Disease tiers 1 and 2 are generally associated with dry (lower 25 percentile of yearly USGS flow estimates from 1937-2003) (USGS 2004) and average (normal range of flow, 25-75 percentiles) years; Tier 3, indicating low disease-intensity, generally is associated with wet years (> 75 percentile of yearly flow estimates) (Table 1).

We calculated mean annual mortality rates for each size class by salinity zone j and disease tier k based on counts of live oysters and recent boxes from each bar over time ($i = 1, \dots, n_{jk}$), using the following equations:

$$s_{jk} = \frac{\sum_i live_{ijk}}{\sum_i (live_{ijk} + newbox_{ijk})} \quad (0.1)$$

$$m_{jk} = -\log(s_{jk}) \quad (0.2)$$

$$totm_{jk} = 1 - \exp^{(-m_{jk} \times T)} \quad (0.3)$$

where: s_{jk} = 1- or 2 -week survival rate,

$live_{jk}$ = number of live oysters,

$newbox_{jk}$ = number of recent boxes,

m_{jk} = instantaneous (1- or 2 -week) mortality,

T = expansion factor to total number of weeks (20) when natural mortality occurs ($T = 20$ for 1-week mortality, and $T = 10$ for 2-week mortality)

$totm_{jk}$ = mean annual mortality rate

across all bars by salinity zone j and disease tier k during the time series. The numerical representation of instantaneous and annual proportional mortality rates above follows Ricker (1975).

For comparison, we also calculated the annual mortality rates by size class (except spat) from counts of recent and old boxes using the estimator

$$totm_{jk} = \frac{\sum_i allboxes_{ijk}}{\sum_i (live_{ijk} + allboxes_{ijk})}, \quad (0.4)$$

where $allboxes_{ijk}$ is the total number of recent and old boxes. The ratio estimators (equations 1.1 and 1.4) provide a weighted mean mortality across individual bars, with weights proportional to the number of live and dead oysters at each bar. Mortality estimates for individual bars are highly skewed (Appendix A). We applied the same assumption as Jordan et al. (2002), Jordan and Coakley (2004), and Ford et al. (2006) that shells from dead oysters (boxes) remain articulated for no more than one year, on average. Our estimates are based on the total box counts from Maryland DNR's dredge surveys of 200 to 400 bars per year from 1991 to 2005. Jordan and Coakley (2004)

based their yearly estimates on data for the 43 disease bars sampled from 1985 to 2000. The mean annual mortality rate, or actual mortality rate (Ricker 1975), based on either method is an estimate of the annual expectation of death for an individual oyster.

We calculated the mortality rates from equations (0.3) and (0.4) and the associated variances by bootstrapping (Efron and Tibshirani 1993). We ran 2000 bootstrap re-samples, each consisting of n_{jk} randomly selected observations (with replacement) where n_{jk} is the number of samples collected across all bars over the time series of fall surveys (or yearly) in the respective salinity zone and disease tier. The standard error and the 95% confidence interval (CI) for the annual mean mortality rates were estimated directly from the distribution of the bootstrap estimates. The mean annual mortality rates based on recent box counts were calculated under the assumption that the TSD was 1-week, and 2-weeks. We then pooled the two bootstrap distributions to obtain estimates of mean mortality rates and variances that represent uncertainty in the TDS (i.e., allowing TSD to vary between 1 and 2 weeks). Our estimates of mortality are expressed as means with associated 95% confidence limits in brackets, unless otherwise noted.

Results

From 1980 to 1985, a period when disease intensity of dermo and MSX were limited, the estimated average annual mortality rate of market-size oysters across salinity zones based on total box counts was 0.153 (0.147-0.158). The natural mortality increased significantly after 1985, with an average annual rate of 0.293 (0.289-0.298) from 1986 to 2006. Mortality estimates were generally higher for dry years (e.g., 2002) with elevated disease levels than for wet years (e.g., 1994) with below-average disease levels (Figures 2 and 3). The discrepancy between our yearly mortality estimates from total box counts and those reported by Jordan and Coakley (2004; Figure 2) is due primarily to a bias in their estimates due to an offset of one year (see Discussion).

The annual mortality rate estimated across all years and salinity regimes based on recent boxes (TSD 1-2 weeks) averaged 0.37 (0.17 - 0.56) and did not differ significantly from our estimated average mortality rate based on total boxes, 0.30 (0.29 - 0.31). The average mortalities by salinity zones based on recent (TSD of 1-2 weeks) and total boxes did not differ significantly (Figure 4), and the 95% confidence intervals for our estimates overlapped with mortality rates provided by Jordan and Coakley (2004). Table 2 shows estimates of the average annual mortality rates by salinity-class and disease-intensity based on recent boxes according to different assumptions for TSD.

Our mortality estimates for market-size oysters by year based on recent boxes (TSD of 1-2 weeks) differed significantly from estimates based on total boxes for 12 of 15 years (Figure 3). Estimates based on the number of recent boxes generally improved the separation of mortality rates for dry and wet years (Figure 3). Mortality estimates based on recent boxes were significantly lower than those derived using total boxes for 5 of 6 wet years and significantly higher for 4 of 5 dry years. In 2003, a wet year, the mortality rate for market-size oysters based on total boxes, 0.40 (0.38 - 0.42), was significantly higher than the mortality rate based on recent boxes, 0.17 (0.06 - 0.28).

Mortality rates based on recent boxes (TSD of 1-2 weeks) for market-size and small oysters by salinity zone increased consistently with increasing salinity and differed greatly from mean mortality rates across salinity

zones in dry years such as 1991 and 2002 (Figure 5). In 2002, for example, which was the second dry year in a row, mortality rates for market-size oysters were 0.88 (0.73 – 1.00), 0.69 (0.44 - 0.97), and 0.43 (0.19 - 0.66) for high, medium, and low salinity classes, respectively, as compared to 0.75 (0.52 - 0.98) for a mean 2002 annual mortality rate across salinity zones. Using a mean mortality rate regardless of salinity would generally underestimate mortality for oysters in high salinity areas and overestimate mortality for oysters in low salinity areas.

To ensure that results from the predictive demographic model being used in the ERA correctly account for the effects of salinity, the mean annual mortality rate based on recent box counts was estimated individually for each salinity class and disease tier and was compared to mortality rates based on total box counts. In contrast to the mean mortality rates based on total box counts, the mean mortality rates based on recent box counts consistently differentiated between salinity classes and disease tiers (Figure 6). In high salinity, mean mortality rates based on recent box counts (TSD of 1-2 weeks) for market-size oysters were 0.79 (0.57 – 1.00), 0.51 (0.27 - 0.75), and 0.23 (0.09 - 0.37) for disease tiers 1, 2, and 3 respectively, whereas respective mean mortality rates based on total box counts for market-size oysters were 0.45 (0.44 - 0.45), 0.53 (0.50 - 0.53), and 0.26 (0.24 - 0.28).

Figure 7 shows monthly mortality rates for market-size and small oysters in June, July, August, and October from the study conducted by Maryland DNR in a medium salinity section of the Choptank River during 2002. For comparison, we included the cumulative mortality rate from June to October and an annual mortality rate using our method based on recent box counts (TSD of 1-2 weeks) in the Maryland DNR fall oyster survey. Monthly mortality rates for market-sized oysters were 0.31 (0.11 - 0.51), 0.36 (0.15 - 0.58), 0.33 (0.14 - 0.52), and 0.22 (0.08 - 0.36) for June to August and October, respectively. The cumulative mortality rate from June to October was 0.78 (0.44 - 0.91); September's mortality rate was imputed using the mean of the August and October estimates. The estimated average annual mortality rate based on the recent box counts from the fall oyster survey for medium salinity and Tier 1 disease intensity was 0.59 (0.34 - 0.83).

Discussion

Estimates of natural mortality rates for oysters (age 1+) based on counts of recent boxes appear to be more accurate (i.e., closer to the true average mortality) than estimates based on total box counts for Chesapeake Bay. Mortality estimates based on recent boxes increase consistently with increasing disease intensity and salinity, and they are higher during years of drought than during wet years, when reduced salinity commonly decreases disease mortality from MSX and dermo (Matthiessen et al. 1990; Gottlieb and Schweighofer 1996). The mortality rates estimated from total box counts, in contrast, did not always reflect variation in mortality due to changes in disease levels linked with freshwater runoff into Chesapeake Bay. These mortality estimates were similar for 2003 (wet) and 2001 (dry), for example. The difference between the mortality rates for 2003 based on recent boxes and those based on total boxes might be attributable to a longer time-to-disarticulation than the one year assumed in this study. Estimates based on total boxes may incorporate mortality over more than one year because boxes may take longer than one year to disarticulate (Christmas et al. 1997). Given the high annual mortality rates from 1999 to 2002 (Figure 4), which probably resulted from three successive dry years and one year near the 25th percentile of runoff, a significant proportion of the old boxes counted in 2003 could represent oysters that died in 2002 and 2001, thus biasing the estimated mortality rate for 2003. The lower than average estimate of mortality for 2003 based on recent boxes is more credible for a wet year with below-average disease intensity.

One reason for the difference between our estimates of annual mortality based on total box counts and those reported by Jordan and Coakley (2004) is sampling variability. Our estimates are based on counts of live oysters and boxes collected from 200 to 400 bars per year, while Jordan and Coakley (2004) restricted their analysis to data from 43 disease bars. This difference, however, probably is small. Of more concern is a systematic error (bias) in the estimates provided by Jordan and Coakley (2004). They used fall box counts to estimate natural mortality for the following year (i.e., October 1985 counts were used to estimate mortality for 1986); however, most natural deaths in a give year probably occur between May and October, just before the fall survey. For example, by following a cohort of market-size oysters through one year from October 1985 to October 1986, we see that individuals are subject to mortality due to fishing from October 1985 to April 1986 and to natural mortality

from October 1985 to October 1986. Hence, natural mortality during 1985 (i.e., November 1984 through October 1985) would have occurred primarily between May 1985 and October 1985, and therefore should be estimated from October 1985 box counts, not from October 1984 counts. The bias caused by the one year offset in Jordan and Coakley's (2004) estimates of M is readily apparent in Figure 3. Estimating natural mortality for the current year based on box counts from the previous year's fall survey could cause a bias in stock assessments of an unknown magnitude and direction. The natural mortality due to disease is strongly linked to climatic conditions; consequently, the offset could result in very large biases for adjacent wet and dry years, with unknown implications for long-term projections.

Estimates of natural mortality are based on important assumptions, such as a qualitative assumption about the time an oyster shell sits before being subject to sedimentation or fouling (recent-box method) and an assumption about the typical time between death and disarticulation of the shell (total-box method). Our estimates of annual mortality using counts of recent boxes are based on the assumption that most natural deaths within a year occur during a 20-week period from May to October. We also assumed that instantaneous (1-week) mortality rates were uniform over the 20-week time period. This assumption was necessary because live oysters and recent boxes only were counted during the fall each year. The mortality study in the Choptank showed variability in monthly mortality rates based on counts of recent boxes (1-2 weeks TSD). Ford and Tripp (1996; Figure 7) also demonstrated that mortality was not uniform over this period and that samples of recent boxes collected in November for some years may significantly underestimate disease mortality in prior months. An additional source of bias in our estimates of annual mortality from counts of recent boxes is that mortality can occur outside this 20-week period, for example due to late winter epizootic outbreaks of MSX, and due to winter and spring freshets. Ford and Tripp (1996) showed that increased mortality caused by MSX can occur during the late winter/early spring; 15% of the small and market-size oysters that survived to the fall survey were killed when MSX outbreaks occurred two or more years in a row. Sources of bias in mortality estimates based on total box counts include violations of the assumption of a one-year disarticulation rate and the assumption that all individuals stay in the same size class for one year. Individuals in a size class (e.g., small oysters) may stay in the same size class between

two fall surveys, or grow larger (i.e., achieve market size) within one year. The possibility that oysters will grow to the next larger size class within a year introduces a bias of unknown magnitude in estimates of annual mortality rates. If recent boxes really are only one or two weeks old, then live oysters within the size class of the recent boxes will not have sufficient growth time to reach the next size class before the boxes of their size cohorts are collected and counted. We recognize that our mortality estimates by salinity zones are subject to bias if predation varies geographically, and if the TSD for recent and old boxes varies by salinity and temperature. TSD for recent and old boxes may decrease progressively with increasing salinity (see Christmas et al. 1997; Ford et al. 2006).

We recommend conducting experiments in Chesapeake Bay to estimate the average TSD of recent and total boxes by salinity zone. Results from experimental studies similar to those conducted by Ford et al. (2006) in Delaware Bay could help to refine the TSD and, hence, the annual mortality estimates for Chesapeake Bay oysters. Mortality rates can then be calibrated through forecasting, starting with the first year in a time series of survey data, for areas where accurate abundance and demographic data are available. Such calibration was conducted using abundance-at-age data from 1994 to 2004 for 23 oyster bars in the James River, Virginia, (data provided by R. Mann, Virginia Institute of Marine Science) to help determine the average TSD of recent boxes. Results indicated that mortality parameters based on the assumption that recent boxes have a TSD of one to two weeks produced population projections that matched the survey observations reasonably.

We used mean estimates of annual mortality by salinity and disease tier and allowed TSD to vary from one to two weeks (Table 2) in the oyster demographic model. After the model randomly selects years between 2004 and 2014 and the climatic condition of each year (dry, average, or wet rainfall year) using block bootstrapping from historic USGS data, the disease tier for each year is assigned randomly; probabilities are based on the historic data (Table 1). Next, the model applies randomly selected mortality rates for the appropriate salinity class and disease tier by drawing randomly from a normal distribution with estimated means and associated variances. The empirical mortality estimates for *C. virginica* presented here can also be useful for specifying approximate mortality rates for *C. ariakensis* in the Chesapeake Bay for oysters that are one-year old or older. A mortality rate for a disease and salinity category may be selected based on an evaluation of this species' susceptibility to

mortality caused by dermo and MSX and to predation. Estimated mortality rates for *C. virginica* for low disease levels (Tier 3; Table 2) could be justified for *C. ariakensis* since *C. ariakensis* appears to experience low disease-related mortality. The Virginia Seafood Council grew triploid *C. ariakensis* from 2003 to spring 2005, and results from their trials in October 2004 showed no MSX prevalence in the oysters, and low levels of dermo (<http://www.vims.edu/vsc/>). Mortality rates from these trials supported our assumption that most mortality occurs during the summer months, and mortality rates for *C. ariakensis* were lower than rates for *C. virginica* (S. Allen, pers. com.). Laboratory studies (Newell et al. in review) comparing the relative susceptibility of juvenile diploids (shell height <25 mm) of both oyster species to invertebrate predators of eastern oyster juveniles suggest that *C. ariakensis* generally will have weaker shells than *C. virginica*. Both species developed stronger shells in response to cues from predators, but *C. virginica* showed inducible changes in shell composition that make it comparatively stronger. If this comparative difference in shell strength persists over time following introduction to Chesapeake Bay, juvenile *C. ariakensis* probably would experience greater mortality due to predation than *C. virginica*, resulting in reduced recruitment to the spawning stock. This differential predation mortality may not hold for oysters larger than 40 mm. We note that the study conducted by Newell et al. (in review) covered a fairly short time period. If *C. ariakensis* is introduced to Chesapeake Bay, it may adapt genetically to develop shell strength similar to the native oyster's. Freeman and Byers (2006) showed that New England mussels can adapt genetically to grow thicker shells in response to invasive predators in fewer than 15 years.

Acknowledgements

We would like to thank the Maryland Department of Natural Resources for providing funding for this project. Ed Weber of Versar, Inc. provided statistical support by running bootstrap analyses. We appreciate the editorial effort by Beth Franks and Carol DeLisle of Versar.

References

Andrews, J. D. 1996. History of *Perkinsus marinus*, a pathogen of oysters in Chesapeake Bay 1950-1984. *J. Shellfish Res.* 15:13-16.

Burreson, E. M. & L. M. Ragone Calvo . 1996. Epizootiology of *Perkinsus marinus* disease of oysters in Chesapeake Bay, with emphasis on data since 1985. *J. Shellfish Res.* 15:17-34.

Carriker, M. R. 1955. Seasonal vertical movement of oyster drills (*Urosalpinx cinerea*). *Proc. Natl. Shellfish Assoc.* 45:190-198.

Christmas, J. F., M. R. McGinty, D. A. Randle, G. F. Smith & S. J. Jordan. 1997. Oyster shell disarticulation in three Chesapeake Bay tributaries. *J. Shellfish Res.* 16:115-123.

Efron, B. & R. J. Tibshirani. 1993. An Introduction to the Bootstrap. Chapman & Hall/CRC Press. 436 pp.

Ford, S. E., M. J. Cummings & E.N. Powell. 2006. Estimating mortality in natural assemblages of oysters. *Estuaries and Coasts* 29:361-374.

Ford, S. E. & M. R. Tripp. 1996. Disease and Defense Mechanisms. In: V. S. Kennedy, R. I. E. Newell & A. F. Eble, editors. The Eastern Oyster *Crassostrea virginica*. Maryland Sea Grant College, University of Maryland, College Park, MD. pp. 581-642.

- Ford, S. E. & H. H. Haskin. 1982. Infection and mortality patterns in strains of oysters *Crassostrea virginica* selected for resistance to the parasite *Haplosporidium nelsoni* (MSX). *J. Parasitol.* 73:368-376.
- Freeman, A. S. & J. E. Byers. 2006. Divergent induced responses to an invasive predator in marine mussel populations. *Science* 313:831-833.
- Garton, D. & W. B. Stickle. 1980. Effects of salinity and temperature on predation rate of *Thais haemastoma* on *Crassostrea virginica* spat. *Biol. Bull.* 158:49-57.
- Giesecker, C.M. 2001. Year 2000 maryland Oyster Disease Status Report. Maryland DNR, Fisheries Service, Sarbanes Cooperative Oxford laboratory. FS-SCOL-01-1. Oxford, Maryland.
- Gottlieb, S. J. & M. E. Schweighofer. 1996. Oysters and the Chesapeake Bay ecosystem: a case for exotic species introduction to improve environmental quality. *Estuaries* 19:639-650.
- Gunter, G. 1979. Studies of the southern oyster borer, *Thais haemastoma*. *Gulf Research Reports* 6:249-260.
- Jordan, S. J. & J. M. Coakley. 2004. Long-term projections of eastern oyster populations under various management scenarios. *J. Shellfish Res.* 23:63-72.

Jordan, S. J., K. N. Greenhawk, C. B. McCollough, J. Vanisko & M.L. Homer. 2002. Oyster biomass, abundance and harvest in northern Chesapeake Bay: trends and forecasts. *J. Shellfish Res.* 21:733-741.

Landers, W. S. & E. W. Rhodes. 1970. Some factors influencing predation by the flatworm, *Stylochus ellipticus* on oysters. *Ches. Sci.* 11:55-60.

Manzi, J. J. 1970. Combined effects of salinity and temperature on feeding, reproductive, and survival rates of *Eupleura caudata* and *Urosalpinx cinerea*. *Biol. Bull.* 138:35-46.

Matthiessen, G. C., S. Y. Feng & L. Leiboviyiz. 1990. Patterns of MSX (*Haplosporidium nelsoni*) infections and subsequent mortality in resistant and susceptible strains of the eastern oyster *Crassostrea virginica* (Gmelin, 1791) in New England. *J. Shellfish Res.* 9:359-366.

NRC (National Research Council). 2004. Non-Native Oysters in the Chesapeake Bay. The National Academies Press, Washington D.C. 325 pp.

Newell, R. I. E., V. S. Kennedy & K. S. Shaw. In Review. Comparative vulnerability to predators, and induced defense responses, of native and non-native oysters in Chesapeake Bay. *Ecol. App.* (In Review).

Pearse, A. S. & G. W. Wharton. 1938. The oyster "leach" *Stylochus inimicus* Palami, associated with oysters on the coasts of Florida. *Ecol. Monograph* 8:605-655.

Ricker, W. E. 1975. Computations and interpretation of biological statistics of fish populations. *Fish. Res. Bd. Canada Bull.* 191. 382 pp.

Southworth, M., J. A. Harding & R. Mann. 2005. The Status of Virginia's Public Oyster Resource 2004. Molluscan Ecology Program, Virginia Institute of Marine Science, Gloucester Point, Virginia.

Tarnowski, M. 2003. Maryland Oyster Population Status Report: 2002 Fall Survey. Maryland Department of Natural Resources, Shellfish Program and Sarbanes Cooperative Oxford Laboratory. 32 pp.

USGS. 2004. Annual mean inflow of stream water into the Chesapeake Bay.
<http://md.water.usgs.gov/monthly/bay.html>.

Figure Captions

Figure 1. Annual mean dermo disease intensity across 43 MD oyster bars and mean discharge (liters/sec) into the Chesapeake Bay from 1991 to 2005 (USGS 2004). High, medium, and low dermo intensity correspond to disease Tiers 1-3; high, medium and low flow correspond to dry (lower 25th percentile), average (25-75th percentiles), and wet years (upper 75th percentile) based on USGS flow estimates from 1937-2005.

Figure 2. Yearly natural mortality rates for market-size oysters based on total boxes across all Maryland bars surveyed, versus estimates of mortality reported in Jordan and Coakley (2004). Error bars for our estimates represent 95% confidence intervals.

Figure 3. Yearly mortality estimates for market-size oysters (> 76 mm) based on recent and total boxes, under the assumptions of average time-since-death (TSD) of 1-2 -weeks and 1-year, respectively. The climatic condition (dry, average, wet) for each year is based on USGS flow estimates. Error bars represent 95% confidence intervals.

Figure 4. Mean annual natural mortality rates by salinity class for market-size oysters over all years (1991-2005). Error bars represent 95% confidence intervals. We assumed an average TSD of 1-year for total boxes, and 1-2-weeks for recent boxes. Mortality estimates per salinity class reported in Jordan and Coakley (2004) are also shown.

Figure 5. Annual natural mortality rates for small and market-size oysters based on counts of recent boxes (TSD=1-2 -weeks) across all bars in each salinity class. Error bars represent 95% confidence intervals. Market-size oysters are defined as being greater than or equal to 76 mm in shell length. Small oysters are defined as being between 40 mm and 75 mm in shell length.

Figure 6. Mean annual natural mortality rates by disease level (tier 1 = high dermo intensity; tier 2 = medium dermo intensity; and tier 3 = low dermo intensity) and salinity class for small and market-size oysters over all years (1991-2005). Error bars represent 95% confidence intervals. We assumed an average TSD of 1-year for total boxes, and 1-2 -weeks for recent boxes.

Figure 7. Monthly and cumulative natural mortality at 6 oyster bars in Choptank River (medium salinity) compared to (1) the estimated 2002 mortality rate for medium salinity and (2) the mortality rate across all medium salinity bars in years with high disease intensity. Error bars represent 95% confidence intervals. All estimates are based on counts of recent boxes (TSD = 1-2 -weeks).

Table 1. Proportion of years (1991-2005) in each disease tier for three climatic condition classes (dry, average, or wet years) as defined by run-off measured by USGS. Tiers 1-3 represent high, medium, and low disease intensity, respectively.

| Condition | Disease Tier | | |
|-----------|--------------|------|------|
| | 1 | 2 | 3 |
| Dry | 0.80 | 0.20 | 0.00 |
| Average | 0.00 | 0.75 | 0.25 |
| Wet | 0.00 | 0.17 | 0.83 |

Table 2. Estimates of proportional annual natural mortality by size class, salinity zone, and disease intensity (Tier), based on counts of live oysters and recent boxes from 1991- 2005 dredge surveys. The relative standard error, $RSE=S.E./Mean$; LCL, and UCL are lower and upper 95% confidence limits.

| TSD Recent | Market-Size Oysters | | | | | | Small Oysters | | | |
|---------------|---------------------|----------|------|------|------|------|---------------|------|------|------|
| | Boxes (Weeks) | Salinity | Tier | Mean | RSE | LCL | UCL | Mean | RSE | LCL |
| 1 | High | 1 | 0.90 | 0.01 | 0.88 | 0.91 | 0.81 | 0.02 | 0.78 | 0.84 |
| 1 | High | 2 | 0.63 | 0.05 | 0.57 | 0.68 | 0.59 | 0.03 | 0.56 | 0.62 |
| 1 | High | 3 | 0.30 | 0.09 | 0.26 | 0.34 | 0.44 | 0.05 | 0.40 | 0.47 |
| 1 | Med | 1 | 0.71 | 0.02 | 0.68 | 0.74 | 0.68 | 0.03 | 0.65 | 0.71 |
| 1 | Med | 2 | 0.54 | 0.04 | 0.51 | 0.57 | 0.49 | 0.05 | 0.45 | 0.53 |
| 1 | Med | 3 | 0.17 | 0.07 | 0.15 | 0.19 | 0.21 | 0.05 | 0.20 | 0.23 |
| 1 | Low | 1 | 0.44 | 0.10 | 0.37 | 0.52 | 0.37 | 0.10 | 0.31 | 0.43 |
| 1 | Low | 2 | 0.28 | 0.10 | 0.23 | 0.32 | 0.20 | 0.10 | 0.17 | 0.23 |
| 1 | Low | 3 | 0.13 | 0.26 | 0.07 | 0.19 | 0.11 | 0.08 | 0.09 | 0.12 |
| 2 | High | 1 | 0.68 | 0.02 | 0.65 | 0.70 | 0.56 | 0.03 | 0.53 | 0.59 |
| 2 | High | 2 | 0.39 | 0.07 | 0.35 | 0.44 | 0.36 | 0.04 | 0.34 | 0.39 |
| 2 | High | 3 | 0.16 | 0.10 | 0.14 | 0.19 | 0.25 | 0.06 | 0.22 | 0.27 |
| 2 | Med | 1 | 0.46 | 0.03 | 0.44 | 0.49 | 0.43 | 0.04 | 0.41 | 0.46 |
| 2 | Med | 2 | 0.32 | 0.04 | 0.30 | 0.34 | 0.28 | 0.06 | 0.26 | 0.31 |
| 2 | Med | 3 | 0.09 | 0.08 | 0.08 | 0.10 | 0.11 | 0.05 | 0.10 | 0.12 |
| 2 | Low | 1 | 0.25 | 0.12 | 0.20 | 0.30 | 0.21 | 0.11 | 0.17 | 0.24 |
| 2 | Low | 2 | 0.15 | 0.10 | 0.13 | 0.18 | 0.10 | 0.11 | 0.09 | 0.12 |
| 2 | Low | 3 | 0.06 | 0.10 | 0.05 | 0.08 | 0.06 | 0.08 | 0.05 | 0.06 |

| | | | | | | | | | | |
|-----|------|---|------|------|------|------|------|------|------|------|
| 1-2 | High | 1 | 0.79 | 0.14 | 0.57 | 1.00 | 0.69 | 0.18 | 0.45 | 0.93 |
| 1-2 | High | 2 | 0.51 | 0.24 | 0.27 | 0.75 | 0.47 | 0.25 | 0.25 | 0.70 |
| 1-2 | High | 3 | 0.23 | 0.31 | 0.09 | 0.37 | 0.34 | 0.28 | 0.16 | 0.53 |
| 1-2 | Med | 1 | 0.59 | 0.21 | 0.34 | 0.83 | 0.56 | 0.22 | 0.31 | 0.80 |
| 1-2 | Med | 2 | 0.43 | 0.26 | 0.21 | 0.64 | 0.39 | 0.27 | 0.18 | 0.59 |
| 1-2 | Med | 3 | 0.13 | 0.32 | 0.05 | 0.21 | 0.16 | 0.31 | 0.06 | 0.26 |
| 1-2 | Low | 1 | 0.34 | 0.29 | 0.15 | 0.54 | 0.29 | 0.30 | 0.12 | 0.46 |
| 1-2 | Low | 2 | 0.22 | 0.31 | 0.08 | 0.35 | 0.15 | 0.33 | 0.05 | 0.25 |
| 1-2 | Low | 3 | 0.10 | 0.33 | 0.03 | 0.16 | 0.08 | 0.33 | 0.03 | 0.13 |




Figure 1




Figure 2




Figure 3




Figure 4







Figure 5



| | |
|---|---|
| ↑ | High Salinity; Tier 1 (High Dermo Prevalence) |
| ↔ | Medium Salinity; Tier 2 (Medium Dermo Prevalence) |
| ↓ | Low Salinity; Tier 3 (Low Dermo Prevalence) |

Figure 6

Market Oysters



Small Oysters




Figure 7

Appendix E. Description of Alternative Restoration Strategies (P. Jones)

Alternative 1--No Action

Under this alternative, a non-native oyster is not introduced into Chesapeake Bay. Oyster management activities are restricted to the native oyster. Oyster population restoration and habitat rehabilitation efforts continue at current levels.

Population restoration

Oyster population restoration activities carried out under Alternative 1 are summarized in Table 1. Maryland hatcheries will produce 225 million seed each year, of which 25 million will be allocated to the Potomac River. Virginia will produce 50 million hatchery seed annually.

Hatchery seed will be planted on oyster sanctuaries, harvest reserves and open harvest areas in the three management regions (Figure 1). The number of hatchery seed planted will range from one million to five million per acre. Differences in the number planted per acre are due to salinity dependent differences in the natural mortality of stocked hatchery seed and different management objectives for sanctuaries, harvest reserves and open harvest areas. The number of seed that will be stocked per acre each year in each region is as follows: Maryland sanctuaries – 2 million per acre, Maryland harvest reserves - 1 million per acre, Potomac River open harvest areas - 1 million per acre and Virginia sanctuaries - 5 million per acre.

In Maryland, 40 million hatchery seed will be planted on 20 acres of oyster bars located in sanctuaries and 160 million hatchery seed will be planted on 160 acres of oyster habitat in harvest reserves each year. In the Potomac River, 25 million hatchery seed will be planted on 25 acres of oyster bars that are open to harvest. Virginia will plant 50 million hatchery seed on 10 acres of oyster bars located in sanctuaries each year. Several hatchery seed plantings will occur on these bars over the 10 year period. As a result, for some bars, the total number of acres planted with hatchery seed will exceed the total size of the bar.

Table 1. Total number of hatchery seed planted, allocation of seed within each management area and number of acres to be planted annually under Alternative 1 – No Action.

| Region | Number of hatchery seed planted annually (in millions) | Allocation of hatchery seed (in millions) | | | Number of acres planted with hatchery seed annually | | |
|----------|--|---|----------|---------------|---|----------|---------------|
| | | sanctuaries | reserves | harvest areas | sanctuaries | reserves | harvest areas |
| Maryland | 200 | 40 | 160 | | 20 | 160 | |
| Potomac | 25 | | | 25 | | | 25 |
| Virginia | 50 | 50 | | | 10 | | |
| Bay wide | 275 | 90 | 160 | 25 | 30 | 160 | 25 |

Under Alternative 1, ten sanctuaries in Maryland and three in Virginia will receive hatchery seed annually (Table 2). In Maryland, approximately 30 million hatchery seed will be planted on sanctuaries in low salinity waters (5 – 12 ppt) and 10 million on sanctuaries in intermediate to high salinity waters (above 12 pt). A total of 200 acres will be planted in Maryland and 100 acres will be planted in Virginia over the 10 year period. Detailed information the location of each site and the number of seed planted on each bar each year is presented in Appendix X.

Table 2. Summary of hatchery seed planting information for Maryland and Virginia sanctuaries under Alternative 1 - No Action.

| Region | Sanctuary | | | Total acres planted with hatchery seed at each location over the 10 year period |
|----------|-------------------|-------------------|--------------|---|
| | Location | Name | size (acres) | |
| Maryland | Upper Bay | Gales Lump | 248 | 20 |
| | Severn River | Severn River | 148 | 20 |
| | Upper Bay | Thomas Pt. South | 46 | 20 |
| | Chester River | Strong Bay | 76 | 20 |
| | Eastern Bay | Mill Hill | 76 | 20 |
| | Choptank River | Cambridge | 142 | 20 |
| | Patuxent River | Trent Hall | 41 | 20 |
| | Potomac River | Dukeheart Channel | 138 | 20 |
| | Upper Bay | Outer Magothy | 167 | 20 |
| | Chester River | Hells Delight | 175 | 20 |
| | | Total | 1,257 | 200 |
| Virginia | Great Wicomico R. | Shell Bar | 14 | 33 |
| | Great Wicomico R. | Haynie Bar | 6 | 17 |
| | Piankatank River | Palaces | 53 | 50 |
| | | Total | 73 | 100 |

Summary information on hatchery seed plantings on Maryland’s harvest reserves is presented in Table 3. In Maryland, ten harvest reserves will receive 160 million hatchery seed annually. A total of 1600 acres will be planted over 10 years. Each year, approximately 128 million seed will be planted on harvest reserves in low salinity waters (5-12 ppt) and 32 million in intermediate to high salinity sites (above 12ppt). Detailed information on the location of each site and the number of seed planted on each bar each year is presented in Appendix 1.

Table 3. Summary of hatchery seed planting information for Maryland harvest reserves under Alternative 1 - No Action.

| Location | Reserve | | Total Acres Planted With Hatchery Seed at Each Location Over the 10 Year Period |
|----------------|--------------------|--------------|---|
| | Name | Size (acres) | |
| Chester River | Blunt | 63 | 100 |
| Choptank River | Bolingbroke Sand | 22 | 40 |
| Chester River | Emory Hollow | 120 | 200 |
| Patuxent River | Broadneck | 271 | 300 |
| Choptank River | Howell Point | 71 | 140 |
| Choptank River | Black Buoy | 88 | 150 |
| Upper Bay | Swan Point | 184 | 220 |
| Upper Bay | Belvedere | 234 | 300 |
| Chester River | Devil's Playground | 38 | 50 |
| Chester River | Boat House | 58 | 100 |
| | Total | 1,149 | 1,600 |

Information on hatchery seed plantings on Potomac River open harvest areas is presented in Table 4. Hatchery seed will be planted on seven oyster bars each year. A total of 250 acres will be planted over the 10 year period. Detailed information on the location of each site and the number of seed planted on each bar each year is presented in Appendix 1.

Table 4. Summary of hatchery seed planting information for Potomac River open harvest areas under Alternative 1 – No Action.

| Open harvest area | | Total acres planted with hatchery seed at each location over the 10 year period |
|-------------------|--------------|---|
| Name | Size (acres) | |
| Ragged Point | 165 | 40 |
| Colonial Beach | 99 | 40 |
| Sheepshead | 107 | 40 |
| Cobb Bar | 68 | 30 |
| Swan Point | 113 | 40 |
| Gum Bar | 44 | 30 |
| Lower Cedar Point | 42 | 30 |
| Total | 638 | 250 |

Habitat rehabilitation

Oyster bar habitat rehabilitation activities to be carried out under Alternative 1 are summarized in Table 5. The locations in which these activities will be undertaken are presented in Figure 1. Maryland will rehabilitate 200 acres of oyster bar habitat each year of the 10 year period. Of this total, approximately 40 acres will be in sanctuaries and 160 acres in open harvest areas. In the Potomac River, habitat rehabilitation activities will be undertaken on 55 acres of oyster bottom in open harvest areas annually. Virginia will add shell to 223-1,484 acres of oyster bottom each year.

Table 5. Total acres of oyster bars rehabilitated within each management area and acres restored in sanctuaries, reserves and open harvest areas under Alternative 1 – No Action.

| Region | Total Number of Acres of Oyster Habitat Rehabilitated Annually | Number of Acres Rehabilitated Annually | | |
|----------|--|--|----------|---------------|
| | | Sanctuaries | Reserves | Harvest Areas |
| Maryland | 200 | 40 | | 160 |
| Potomac | 55 | | | 55 |
| Virginia | 223-1,484 | 223-1,484 | | |

Habitat rehabilitation will occur on three sanctuaries (Table 6) and 14 open harvest areas in Maryland (Table 7) and 12 open harvest areas in the Potomac River. Virginia will plant shell on 19-55 sanctuary sites each year. Details of shell planting activities for Virginia are presented in Appendix _.

Over the 10 year study period, 400 acres of sanctuaries and 1600 acres of open harvest areas in Maryland and 550 acres of open harvest areas in the Potomac River will be rehabilitated. In Virginia, a total of 9,501 acres of oyster bars in sanctuaries will receive shell over the 10 year study period.

Table 6. Summary of habitat rehabilitation information for Maryland sanctuaries under Alternative 1 – No Action.

| Location | Sanctuaries | | Total acres rehabilitated at each location over the 10 year period |
|------------------|-----------------|--------------|--|
| | Name | Size (acres) | |
| Choptank River | Black Walnut | 211 | 210 |
| St. Mary's Shore | Point Look-In | 111 | 110 |
| Lower Bay | NW Middleground | 103 | 80 |
| | Total | 425 | 400 |

Table 7. Summary of habitat rehabilitation information for Maryland and Potomac River open harvest areas under Alternative 1 – No Action.

| Open Harvest Areas | | | | | |
|--------------------|------------------|----------------------|--------------|--|-------|
| Region | Location | Name | Size (acres) | Total acres rehabilitated at each location over the 10 year period | |
| Maryland | Tangier Sound | Piney Island West(a) | 154 | 150 | |
| | Tangier Sound | Piney Island West(b) | 204 | 200 | |
| | Manokin River | Drum Point(a) | 22 | 20 | |
| | Manokin River | Drum Point(b) | 19 | 10 | |
| | Tangier Sound | Harris Addition | 113 | 100 | |
| | Little Annessex | Great Rock | 323 | 320 | |
| | Pocomoke Sound | Gumby | 117 | 100 | |
| | Pocomoke Sound | Marumsco | 128 | 120 | |
| | Eastern Bay | Rich Neck | 82 | 80 | |
| | Eastern Bay | Lowes Point | 76 | 70 | |
| | St. Mary's River | Cherry | 36 | 30 | |
| | Lower Bay | Butler | 65 | 50 | |
| | Choptank | Wild Cherry Tree | 283 | 250 | |
| | Main Bay | Lulu | 152 | 100 | |
| | | Total | | 1,774 | 1,600 |
| | Potomac | | Jones Shore | 4 | 4 |
| | | | Poseys Bluff | 110 | 110 |
| | | Great Neck | 98 | 94 | |
| | | Kitts Point | 3 | 2 | |
| | | Hog Island | 22 | 20 | |
| | | Heron Island | 38 | 30 | |
| | | St. Georges | 17 | 10 | |
| | | Ragged Point | 165 | 160 | |
| | | Thickett Point | 11 | 10 | |
| | | Bonums | 12 | 10 | |
| | | Tall Timbers | 42 | 40 | |
| | King Copsico | 62 | 60 | | |
| | Total | | 584 | 550 | |

Alternative 2--Expand Native Oyster Restoration Program

Under Alternative 2, a non-native oyster is not introduced in the Bay. Instead, native oyster recovery efforts undergo significant expansion from current levels. Changes from the status quo include large-scale increases in oyster population restoration and habitat rehabilitation activities. Changes from the status quo also include an assessment of cultch limitations and development of potential long-term solutions, and evaluation of the feasibility of the development, production, and deployment of large quantities of disease resistant strain(s) of the native oyster for brood stock enhancement. For alternative 2, the evaluation of Maryland sanctuaries is subset into Alternative 2a and Alternative 2b. The two subsets differ in the number of sanctuaries that receive seed and the location of the sanctuaries.

Population restoration

Under Alternative 2, hatchery seed plantings increase from 200 million to 2 billion annually in Maryland, 25 million to 125 million a year in the Potomac River and 50 to 200 million in Virginia (Table 8). The number of acres of sanctuaries planted with hatchery seed annually increases from 75 to 750 in Maryland and from 10 to 40 in Virginia. Harvest reserve plantings increase from 50 to 500 acres per year and open harvest area hatchery seed

plantings increase from 25 to 125 acres annually over the 10 year period. The number of seed stocked per acre each year is the same as in Alternative 1: Maryland sanctuaries – 2 million per acre, Maryland harvest reserves and Potomac River open harvest areas - 1 million per acre, and Virginia sanctuaries - 5 million per acre.

Table 8. Total number of hatchery seed planted (millions), allocation of seed within each management area and number of acres to be planted annually under Alternative2 – Expand Native Oyster Restoration Program

| Year | Number of Hatchery Seed planted annually (in billions) | | | Allocation of Hatchery Seed (in billions) | | | | Number of Acres Planted with hatchery seed annually | | | |
|-------|--|-------|-------|---|-------|-------------|-------------------|---|-----|-------------|-------------------|
| | MD | Pot | VA | Sanctuaries | | Reserves MD | Harvest Areas Pot | Sanctuaries | | Reserves MD | Harvest Areas Pot |
| | | | | MD | VA | | | MD | VA | | |
| 1 | 0.200 | 0.025 | 0.050 | 0.150 | 0.050 | 0.050 | 0.025 | 75 | 10 | 50 | 25 |
| 2 | 0.350 | 0.045 | 0.100 | 0.263 | 0.100 | 0.087 | 0.045 | 131 | 20 | 87 | 45 |
| 3 | 0.500 | 0.070 | 0.200 | 0.375 | 0.200 | 0.125 | 0.070 | 188 | 40 | 125 | 70 |
| 4 | 0.750 | 0.090 | 0.200 | 0.563 | 0.200 | 0.187 | 0.090 | 281 | 40 | 187 | 90 |
| 5 | 1.000 | 0.120 | 0.200 | 0.750 | 0.200 | 0.250 | 0.120 | 375 | 40 | 250 | 120 |
| 6 | 1.500 | 0.125 | 0.200 | 1.125 | 0.200 | 0.375 | 0.125 | 563 | 40 | 375 | 125 |
| 7 | 2.000 | 0.125 | 0.200 | 1.500 | 0.200 | 0.500 | 0.125 | 750 | 40 | 500 | 125 |
| 8 | 2.000 | 0.125 | 0.200 | 1.500 | 0.200 | 0.500 | 0.125 | 750 | 40 | 500 | 125 |
| 9 | 2.000 | 0.125 | 0.200 | 1.500 | 0.200 | 0.500 | 0.125 | 750 | 40 | 500 | 125 |
| 10 | 2.000 | 0.125 | 0.200 | 1.500 | 0.200 | 0.500 | 0.125 | 750 | 40 | 500 | 125 |
| Total | 12.300 | 0.975 | 1.750 | 9.226 | 1.750 | 3.074 | 0.975 | 4,613 | 350 | 3,074 | 975 |

As was the case in Alternative 1, hatchery seed in Alternative 2 will be planted on oyster bars located in sanctuaries, harvest reserves and open harvest areas in the three management regions (Figure 2 and 3). Several hatchery seed plantings will occur on these bars over the 10 year period. As a result, for some bars, the total number of acres planted with hatchery seed will exceed the total size of the bar.

In alternatives 2a and 2b there are differences in the number of sanctuaries that receive seed and the location of the sanctuaries (Table 9). In Alternative 2a, a total of 32 sanctuaries, which are all located in low salinity waters (5 -12 ppt), will be planted with hatchery seed (Figure 2). In Alternative 2b, a total of 39 sanctuary areas receive seed. Of these, 26 are in low salinity, and 13 are in moderate to high salinity waters (Figure 3).

Table 9. Summary of hatchery seed planting information for Maryland sanctuaries under Alternatives 2a and 2b-Expanded Native Oyster Restoration Program.

| Sanctuary | | | Total acres planted with hatchery seed at each location over the 10 year period | |
|-----------------|--------------------|--------------|---|----------------|
| Location | Name | Size (acres) | Alternative 2a | Alternative 2b |
| Upper Bay | Gales Lump | 248 | 295 | 270 |
| Upper Bay | Severn River | 208 | 253 | 190 |
| Upper Bay | Thomas Point South | 46 | 70 | 50 |
| Middle Bay | Hering Bay | 30 | 53 | 50 |
| Chester River | Strong Bay | 76 | 115 | 110 |
| Chester River | Ringgold | 16 | 30 | 30 |
| Chester River | ORA Zone A | 116 | 165 | 105 |
| Eastern Bay | Mill Hill | 76 | 135 | 130 |
| Eastern Bay | Cabin Creek | 22 | 40 | 40 |
| Miles River | Miles River | 49 | 70 | 70 |
| Choptank River | Howell Point | 5 | 10 | 10 |
| Choptank River | Oxford Lab | 4 | 10 | 10 |
| Choptank River | Cambridge | 95 | 134 | 234 |
| Choptank River | States Bank | 20 | 40 | 40 |
| Patuxent River | Elbow/Teague | 32 | 60 | 60 |
| Patuxent River | Kitts Marsh | 13 | 20 | 20 |
| Patuxent River | Trent Hall | 63 | 122 | 80 |
| Wicomico River | Russell | 38 | 71 | 71 |
| Potomac River | Dukeheart Channel | 138 | 222 | 212 |
| Upper Bay | Bodkin Point South | 634 | 705 | 680 |
| Upper Bay | Outer Magothy | 377 | 490 | 355 |
| Upper Bay | Tea Table | 274 | 306 | 251 |
| Chester River | Hells Delight | 175 | 235 | 175 |
| Choptank River | Royston | 175 | 231 | 168 |
| Choptank River | Castle Haven | 19 | 40 | 40 |
| Potomac River | Bluff Point Lumps | 5 | 10 | 10 |
| Upper Bay | Broad Creek | 108 | 107 | |
| Upper Bay | Love Point | 94 | 91 | |
| Upper Bay | Swan Point | 75 | 75 | |
| Chester River | Long Point | 59 | 58 | |
| Upper Bay | Graveyard | 43 | 40 | |
| Upper Bay | Craighill | 310 | 310 | |
| | Subtotal | 3,643 | | 3,461 |
| Middle Bay | Plum Point | 91 | | 109 |
| Patuxent River | Mears | 20 | | 32 |
| Patuxent River | Neal Addition | 77 | | 120 |
| Choptank river | Cook Point | 72 | | 120 |
| Lower Bay | Dorchester PD | 14 | | 20 |
| Tangier Sound | Somerset PD | 369 | | 270 |
| Lower Bay | St Mary's PD | 114 | | 107 |
| Lower Bay | SW Middleground | 149 | | 140 |
| Pocomoke Sound | Kitts Creek | 33 | | 50 |
| St Mary's River | Piny Point | 13 | | 20 |
| Potomac River | Jones Shore | 6 | | 10 |
| Tangier Sound | Sharkfin Shoal | 90 | | 109 |
| Potomac River | Cooper Creek | 27 | | 50 |
| | Subtotal | 1,075 | | 1,157 |
| | Total | 4,718 | 4,613 | 4,618 |

Under Alternative 2, three sanctuaries in Virginia will receive hatchery seed annually (Table 10). A total of 350 acres of hatchery seed will be planted over the 10 year period.

Table 10. Summary of hatchery seed planting information for Virginia sanctuaries under Alternative 2 -Expand Native Oyster Restoration Program.

| Location | Sanctuary | | Total acres planted with hatchery seed at each location over the 10 year period |
|-------------------|------------|--------------|---|
| | Name | size (acres) | |
| Great Wicomico R. | Shell Bar | 14 | 122 |
| Great Wicomico R. | Haynie Bar | 6 | 53 |
| Piankatank River | Palaces | 53 | 175 |
| | Total | 73 | 350 |

Summary information on hatchery seed planting on Maryland’s harvest reserves is presented in Table 11. In Maryland, 18 harvest reserves, which are all in low salinity waters, will receive 50 to 500 million hatchery seed annually. Over the ten year study period a total of 3,074 acres will be planted with hatchery seed.

Table 11. Summary of hatchery seed planting information for Maryland harvest reserves under Alternative 2 - Expand Native Oyster Restoration Program.

| Location | Reserve | | Total acres planted with hatchery seed at each location over the 10 year period |
|----------------|--------------------|--------------|---|
| | Name | Size (acres) | |
| Chester River | Blunt | 63 | 110 |
| Choptank River | Bolingbroke Sand | 22 | 40 |
| Chester River | Emory Hollow | 120 | 200 |
| Patuxent River | Broadneck | 271 | 220 |
| Choptank River | Howell Point | 71 | 120 |
| Choptank River | Black Buoy | 88 | 174 |
| Upper Bay | Swan Point | 184 | 280 |
| Belvedere | Belvedere | 234 | 220 |
| Chester River | Devil's Playground | 38 | 50 |
| Chester River | Boat House | 58 | 100 |
| Upper Bay | Six foot knoll | 854 | 500 |
| Upper Bay | Hackett Point | 213 | 196 |
| West Wicomico | Lancaster | 250 | 250 |
| Choptank River | Dickinson | 261 | 250 |
| Patuxent River | Holland Point | 67 | 56 |
| Miles River | Persimmon Tree | 233 | 196 |
| Wye River | Middleground | 12 | 12 |
| Choptank River | Lighthouse | 145 | 100 |
| | Total | 3,184 | |

Information on hatchery seed plantings on Potomac River open harvest areas is presented in Table 12. Hatchery seed will be planted on seven bars each year. A total of 975 acres will be planted over the 10 year period

Table 12. Summary of hatchery seed planting information for Potomac River open harvest areas under Alternative 2 - Expand Native Oyster Restoration Program.

| Open Harvest Area | | Total acres planted with hatchery seed at each location over the 10 year period Alternative 2b |
|-------------------|--------------|--|
| Name | Size (acres) | |
| Ragged Point | 165 | 207 |
| Colonial Beach | 99 | 152 |
| Sheepshead | 107 | 161 |
| Cobb Bar | 68 | 117 |
| Swan Point | 113 | 168 |
| Gum Bar | 44 | 86 |
| Lower Cedar Point | 42 | 84 |
| Total | 638 | 975 |

Habitat rehabilitation

Oyster bar habitat rehabilitation activities under Alternative 2 are summarized in Table 13. Maryland will rehabilitate 400 acres of oyster bar habitat each year of the 10 year period. Of this total, approximately 80 acres in open harvest areas and 320 acres in sanctuaries will be rehabilitated. In the Potomac River, habitat rehabilitation activities will be undertaken on 110 acres of oyster bottom in open harvest areas each year. In Virginia 522- 2,850 acres of oyster habitat in sanctuaries will receive shell each year.

Table 13. Total acres of oyster bars rehabilitated within each management area and acres restored in sanctuaries, reserves and open harvest areas under Alternative 2 – Expand Native Oyster Restoration Program.

| Region | Total number of acres of oyster habitat rehabilitated annually | Number of acres rehabilitated annually | | |
|----------|--|--|----------|---------------|
| | | sanctuaries | reserves | harvest areas |
| Maryland | 400 | 320 | | 80 |
| Potomac | 110 | | | 110 |
| Virginia | 522-2,850 | 522-2,850 | | |

Habitat rehabilitation will be undertaken on 11 sanctuaries in Maryland (Table 14). A total of eight open harvest areas in Maryland and 12 open harvest areas in the Potomac River will also be rehabilitated (Tables 15). In Virginia 31 to 78 sanctuaries will receive shell each year (Appendix _). Over the 10 year study period, 3,200 acres of sanctuaries and 800 acres of open harvest areas in Maryland and 1,100 acres in the Potomac River will be rehabilitated. In Virginia a total of 16,899 acres will receive shell over the 10 year period.

Table 14. Summary of habitat rehabilitation information for Maryland sanctuaries under Alternative 2 – Expand Native Oyster Restoration Program.

| Sanctuaries | | | Total acres planted with oyster shell at each location over the 10 year period |
|------------------|-----------------|--------------|--|
| Location | Name | Size (acres) | |
| Choptank River | Black Walnut | 211 | 210 |
| Lower Bay | Point Look-In | 111 | 110 |
| Choptank River | Stone Rock | 1,113 | 1,100 |
| Herring Bay | Holland Point | 394 | 390 |
| Lower Bay | NW Middleground | 103 | 100 |
| Lower Bay | Church Creek | 135 | 130 |
| Little Choptank | Ragged Point | 188 | 180 |
| Little Choptank | Nine Acres | 204 | 200 |
| St. Mary's Shore | Rock Beach | 351 | 350 |
| Eastern Bay | Bald Eagle | 120 | 120 |
| West Wicomico | White Point | 311 | 310 |
| Total | | 3,241 | 3,200 |

Table 15. Summary of habitat rehabilitation information for Maryland and the Potomac River open harvest areas under Alternative 2 - Expand Native Oyster Restoration Program.

| Open harvest areas | | | | Total acres planted with oyster shell at each location over the 10 year period |
|--------------------|------------------|----------------------|--------------|--|
| Region | Location | Name | Size (acres) | |
| Maryland | Tangier Sound | Piney Island West(b) | 204 | 200 |
| | Manokin River | Drum Point(a) | 22 | 20 |
| | Tangier Sound | Harris Addition | 113 | 100 |
| | Little Annessex | Great Rock | 323 | 200 |
| | Pocomoke Sound | Marumsco | 128 | 120 |
| | Eastern Bay | Rich Neck | 82 | 80 |
| | St. Mary's River | Cherry | 36 | 30 |
| | Lower Bay | Butler | 65 | 50 |
| | Total | | 973 | 800 |
| | Potomac | | Jones Shore | 4 |
| | | Poseys Bluff | 110 | 220 |
| | | Great Neck | 98 | 194 |
| | | Kitts Point | 3 | 4 |
| | | Hog Island | 22 | 44 |
| | | Heron Island | 38 | 74 |
| | | St. Georges | 17 | 32 |
| | | Ragged Point | 165 | 280 |
| | | Thickett Point | 11 | 20 |
| | | Bonums | 12 | 20 |
| Potomac | | Tall Timbers | 42 | 80 |
| | | King Copsico | 62 | 124 |
| | Total | | 584 | 1,100 |

2.3. Comparison Of Restoration Alternatives 1 and 2

There are significant increases in the numbers of acres of hatchery seed planted under Alternative 2 compared to Alternative 1 (Table 16). Over the 10 year study period, 200 acres in sanctuaries are planted with hatchery seed under Alternative 1 compared to 4,613 acres under Alternative 2. For harvest reserves in Maryland, 1,600 acres are planted with hatchery seed under Alternative 1 versus 3,074 acres under Alternative 2. In Virginia, the number of acres planted with hatchery seed in sanctuaries increases from 100 for Alternative 1 to 350 for Alternative 2. In the Potomac, the number of open harvest areas seeded increases from 250 in Alternative 1 to 975 in Alternative 2.

In Maryland, the amount of habitat rehabilitated in sanctuaries over the 10 year period increases from 400 acres under Alternative 1 to 3,200 acres under Alternative 2. One result of the increase in the rehabilitation of oyster habitat in sanctuaries is a decline in rehabilitated habitat in open harvest areas in Maryland from 1,600 acres in Alternative 1 to 800 acres rehabilitated in Alternative 2 over the 10 year period. In the Potomac River, 550 acres are restored in Alternative 1 compared to 1100 acres in Alternative 2. In Virginia 9,501 acres of oyster habitat are restored in Alternative 1 and 16,899 acres in Alternative 2.

Table 16. Comparison of number of acres planted under Alternative 1 – No Action and Alternative 2 - Expand Native Oyster Restoration Program.

| Year | Maryland | | | | Virginia | | Potomac | |
|-------|-------------|-------|----------|-------|-------------|-------|---------------|-------|
| | Sanctuaries | | Reserves | | Sanctuaries | | Harvest areas | |
| | Alt 1 | Alt 2 | Alt 1 | Alt 2 | Alt 1 | Alt 2 | Alt 1 | Alt 2 |
| 1 | 20 | 75 | 160 | 50 | 10 | 10 | 25 | 25 |
| 2 | 20 | 131 | 160 | 87 | 10 | 20 | 25 | 45 |
| 3 | 20 | 188 | 160 | 125 | 10 | 40 | 25 | 70 |
| 4 | 20 | 281 | 160 | 187 | 10 | 40 | 25 | 90 |
| 5 | 20 | 375 | 160 | 250 | 10 | 40 | 25 | 120 |
| 6 | 20 | 563 | 160 | 375 | 10 | 40 | 25 | 125 |
| 7 | 20 | 750 | 160 | 500 | 10 | 40 | 25 | 125 |
| 8 | 20 | 750 | 160 | 500 | 10 | 40 | 25 | 125 |
| 9 | 20 | 750 | 160 | 500 | 10 | 40 | 25 | 125 |
| 10 | 20 | 750 | 160 | 500 | 10 | 40 | 25 | 125 |
| Total | 200 | 4,613 | 1,600 | 3,074 | 100 | 350 | 250 | 975 |

References

Appendix _ Location, size and detailed planting information for each sanctuary, harvest reserve and open harvest area to receive hatchery seed, and location and size information for each area to be rehabilitated under Alternative I - No Action and Alternative 2-Expand Native Oyster Restoration Program.




Figure 1. Location of hatchery seed plantings on sanctuaries, harvest reserves and open harvest areas and location of oyster bar rehabilitation activities for Alternative 1 – No Action.




Figure 2. Location of hatchery seed plantings on sanctuaries, harvest reserves and open harvest areas and location of oyster bar rehabilitation activities for Alternative 2a – Expand Native Oyster Restoration.




Figure 3. Location of hatchery seed plantings on sanctuaries, harvest reserves and open harvest areas and location of oyster bar rehabilitation activities for Alternative 2b – Expand Native Oyster Restoration.

Appendix F. Explanation for effects of harvest

Harvest had less effect on modeled abundances than might intuitively be expected for several reasons. The most important cause is that harvest does not simply simulate a large removal of a healthy population, as does an increase in natural mortality due to disease; harvest occurs as an additional removal of remaining larger oysters from a population that has already experienced relatively high natural mortality rates due to disease. Under these conditions, the year class that has just reached market size, but has not yet experienced natural mortality or harvest rates for market sized oysters, composes a large proportion of the total number of market sized oysters.

We use a simplified example to illustrate. Here we assume that a bar has constant salinity, disease level, natural mortality rates, and recruitment of 1,000 spat per year. If the bar is located in medium salinity and experiences tier 2 disease prevalence, then mean natural mortality rates are 0.39 for small oysters and 0.43 for market sized oysters (Table 3). If no harvest occurs, a cohort of 1,000 spat recruited would decline to 610 oysters at age 1 (i.e., $1,000 * 0.61$), 372 oysters at age 2, etc., as listed below. The total number of market sized oysters on the bar would be 527 (i.e., $227 + 129 + \dots + 0$).

| Age | Natural mortality rate | Size | N |
|--------------|------------------------|--------|-----|
| 1 | 0.39 | Small | 610 |
| 2 | 0.39 | Small | 372 |
| 3 | 0.39 | Market | 227 |
| 4 | 0.43 | Market | 129 |
| 5 | 0.43 | Market | 74 |
| 6 | 0.43 | Market | 42 |
| 7 | 0.43 | Market | 24 |
| 8 | 0.43 | Market | 14 |
| 9 | 0.43 | Market | 8 |
| 10 | 0.43 | Market | 4 |
| 11 | 0.43 | Market | 3 |
| 12 | 0.43 | Market | 1 |
| 13 | 0.43 | Market | 1 |
| 14 | 0.43 | Market | 0 |
| Total Market | | | 527 |

If harvest begins on the bar at a constant rate, the numbers will reach a new equilibrium after several years. For example, if the bar experiences a 20% harvest rate, small oysters experience 39% natural mortality, and then an additional 2.2% incidental harvest (10% of the catch of market-size oysters, Section 2.11; i.e., $0.2 / 0.9 = X / 0.1$, $X \approx 0.022$). Market-size oysters experience 43% natural mortality, and then 20% of the remaining oysters are harvested. Note that a model year begins in the fall, after recruitment and natural mortality are calculated because most natural mortality due to disease and predation is believed to occur during summer (Andrews 1996; Burreson and Ragone Calvo 1996; Ford and Tripp 1996; Ford and Haskin 1982; Carriker 1955; Manzi 1970; Gunter 1979; Garton and Stickle 1980; Pearse and Wharton 1938; Landers and Rhodes 1970; citations in Vølstad et al. In Press, Appendix

D). Harvest occurs during winter (Figure 9). Thus harvest corresponds to the exploitation rate (F; Ricker 1975) for the winter, and occurs independently of natural mortality. The population will stabilize after six years to a total of 400 market-size oysters (i.e., $217 + 99 + \dots + 0$). Six-hundred-ten oysters will reach small size each year from the 1,000 spat recruited the previous year, which experience 39% mortality. Other yearly abundances (N_f) are calculated by multiplying the number of oysters alive from the previous year by the appropriate survival rate.

| Age | Natural mortality rate | Size | Year 0, No harvest | Year 1 | | | Year 2 | | | Year 3 | | | Year 4 | | | Year 5 | | | Year 6 | | |
|---------------------------|------------------------|--------|--------------------|---------|---------|---------|--------|-------|-------|--------|-------|-------|--------|-------|-------|--------|-------|-------|--------|-------|-------|
| | | | | N_h^1 | N_r^2 | N_f^3 | N_h | N_r | N_f |
| 1 | 0.39 | Small | 610 | 13 | 597 | 610 | 13 | 597 | 610 | 13 | 597 | 610 | 13 | 597 | 610 | 13 | 597 | 610 | 13 | 597 | 610 |
| 2 | 0.39 | Small | 372 | 8 | 364 | 364 | 8 | 356 | 364 | 8 | 356 | 364 | 8 | 356 | 364 | 8 | 356 | 364 | 8 | 356 | 364 |
| 3 | 0.39 | Market | 227 | 45 | 182 | 222 | 44 | 178 | 217 | 43 | 174 | 217 | 43 | 174 | 217 | 43 | 174 | 217 | 43 | 174 | 217 |
| 4 | 0.43 | Market | 129 | 26 | 104 | 104 | 21 | 83 | 101 | 20 | 81 | 99 | 20 | 79 | 99 | 20 | 79 | 99 | 20 | 79 | 99 |
| 5 | 0.43 | Market | 74 | 15 | 59 | 59 | 12 | 47 | 47 | 9 | 38 | 46 | 9 | 37 | 45 | 9 | 36 | 45 | 9 | 36 | 45 |
| 6 | 0.43 | Market | 42 | 8 | 34 | 34 | 7 | 27 | 27 | 5 | 22 | 22 | 4 | 17 | 21 | 4 | 17 | 21 | 4 | 16 | 21 |
| 7 | 0.43 | Market | 24 | 5 | 19 | 19 | 4 | 15 | 15 | 3 | 12 | 12 | 2 | 10 | 10 | 2 | 8 | 10 | 2 | 8 | 9 |
| 8 | 0.43 | Market | 14 | 3 | 11 | 11 | 2 | 9 | 9 | 2 | 7 | 7 | 1 | 6 | 6 | 1 | 4 | 4 | 1 | 4 | 4 |
| 9 | 0.43 | Market | 8 | 2 | 6 | 6 | 1 | 5 | 5 | 1 | 4 | 4 | 1 | 3 | 3 | 1 | 3 | 3 | 1 | 2 | 2 |
| 10 | 0.43 | Market | 4 | 1 | 4 | 4 | 1 | 3 | 3 | 1 | 2 | 2 | 0 | 2 | 2 | 0 | 1 | 1 | 0 | 1 | 1 |
| 11 | 0.43 | Market | 3 | 1 | 2 | 2 | 0 | 2 | 2 | 0 | 1 | 1 | 0 | 1 | 1 | 0 | 1 | 1 | 0 | 1 | 1 |
| 12 | 0.43 | Market | 1 | 0 | 1 | 1 | 0 | 1 | 1 | 0 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| 13 | 0.43 | Market | 1 | 0 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 14 | 0.43 | Market | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total market-size oysters | | | 527 | 462 | | | 428 | | | 412 | | | 405 | | | 402 | | | 400 | | |

¹ N_h Number harvested

² N_r Number remaining after harvest

³ N_f Final number after recruitment and mortality

We conducted similar calculations for harvest rates of 40%, 60%, and 80% to determine the population structure of the bar at equilibrium. Under these conditions, a 20% harvest rate results in 76% of the number of market-size oysters that would have been present if no harvest occurred (i.e., $100 \times 400 / 527$), but an 80% harvest rate still results in 40% of the number of markets that would have been present if there were no harvest. This is because most of the population of market-size oysters consists of age-3 animals under any harvest rate.

Number of oysters under constant conditions, tier 2 mortality, medium salinity:

| Age | Size Class | Natural mortality rate | No harvest | Harvest rate | | | |
|-----|------------|------------------------|------------|--------------|-------|-------|-------|
| | | | | 20% | 40% | 60% | 80% |
| 0 | Spat | | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 |
| 1 | Small | 0.39 | 610 | 610 | 610 | 610 | 610 |
| 2 | Small | 0.39 | 372 | 364 | 356 | 350 | 339 |
| 3 | Market | 0.39 | 227 | 217 | 207 | 199 | 188 |
| 4 | Market | 0.43 | 129 | 99 | 71 | 45 | 21 |

| | | | | | | | |
|------------------------------------|--------|------|-----|-----|-----|-----|-----|
| 5 | Market | 0.43 | 74 | 45 | 24 | 10 | 2 |
| 6 | Market | 0.43 | 42 | 21 | 8 | 2 | 0 |
| 7 | Market | 0.43 | 24 | 9 | 3 | 1 | 0 |
| 8 | Market | 0.43 | 14 | 4 | 1 | 0 | 0 |
| 9 | Market | 0.43 | 8 | 2 | 0 | 0 | 0 |
| 10 | Market | 0.43 | 4 | 1 | 0 | 0 | 0 |
| 11 | Market | 0.43 | 3 | 1 | 0 | 0 | 0 |
| 12 | Market | 0.43 | 1 | 0 | 0 | 0 | 0 |
| 13 | Market | 0.43 | 1 | 0 | 0 | 0 | 0 |
| 14 | Market | 0.43 | 0 | 0 | 0 | 0 | 0 |
| Total market-size oysters | | | 527 | 400 | 315 | 258 | 213 |
| Percent of mkt pop with no harvest | | | | 76 | 60 | 49 | 40 |

The effect is more pronounced when natural mortality rates are higher. For example, if the same calculations are performed using tier 1 mortality rates in high salinity (Table 3), the population of market-size oysters consists almost entirely of age-3 animals. In this case, an 80% harvest rate still yields 69% of the number of market-sized oysters that would have survived if no harvest had occurred.

Number of oysters under constant conditions, tier 1 mortality, high salinity:

| Age | Size Class | Natural mortality rate | No harvest | Harvest rate | | | |
|------------------------------------|------------|------------------------|------------|--------------|-------|-------|-------|
| | | | | 20% | 40% | 60% | 80% |
| 0 | Spat | | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 |
| 1 | Small | 0.69 | 310 | 310 | 310 | 310 | 310 |
| 2 | Small | 0.69 | 96 | 94 | 92 | 90 | 88 |
| 3 | Market | 0.79 | 20 | 19 | 19 | 18 | 17 |
| 4 | Market | 0.79 | 4 | 3 | 2 | 1 | 1 |
| 5 | Market | 0.79 | 1 | 1 | 0 | 0 | 0 |
| 6 | Market | 0.79 | 0 | 0 | 0 | 0 | 0 |
| Total market-size oysters | | | 25 | 23 | 21 | 19 | 17 |
| Percent of mkt pop with no harvest | | | | 91 | 83 | 75 | 69 |

Jim Wesson of the Virginia Marine Resources Commission recently presented empirical data that support this result. A comparison of oyster population structures on harvested bars and nearby unharvested bars revealed that density of market-size oysters on bars that were unharvested for several years were only slightly greater than densities on harvested bars.

These examples neglect the stock-recruitment relationship, which would increase differences in abundance among scenarios slightly. However, the effect is small given the population on most bars in the model. It is important to note that small oysters provide some reproduction before reaching sizes when they are exposed to the greatest natural mortality and fishing mortality. These oysters provide some protection from driving small populations to extinction in the model, as they do in the Bay.

A second reason that harvest was less important than expected is that a large portion of the habitat layer was designated as sanctuaries, reserves, or closed areas that were not fished. Approximately 22% of the market-size oysters in the starting population were located on bars that were not fished. This means that nominal harvest rates of 20, 40, 60, and 80 percent were actually 16, 31, 47, and 62 percent harvest rates, respectively, for the first year of the simulation. All other conditions being equal, the population on unfished bars grows faster than on similar fished bars in the model, thereby increasing the proportion of the population not exposed to fishing with each yearly iteration. Sensitivity analysis revealed that a 10% poaching rate reduced the total population of the Bay by 13%, and a 25% poaching rate reduced the population by 16%.

Overall, the model produced credible results. Although absolute differences in predicted abundance were not large in relation to the restoration goal, percent changes in abundance varied with harvest rates within the expected range (Figures 11-12, and below).

| | Harvest rate | | | | |
|--|---------------|---------------|---------------|---------------|---------------|
| | No harvest | 20% | 40% | 60% | 80% |
| Alternative 1 abundance | 469,327,386 | 374,725,116 | 330,207,716 | 313,302,325 | 304,097,062 |
| Percent of market-size population in no-harvest scenario | | 0.80 | 0.70 | 0.67 | 0.65 |
| Alternative 2A abundance | 1,761,651,246 | 1,605,625,007 | 1,396,503,468 | 1,278,520,298 | 1,170,811,992 |
| Percent of market-size population in no-harvest scenario | | 0.91 | 0.79 | 0.73 | 0.66 |

As noted in the text, we do not suggest that fishing is unimportant in controlling oyster populations in the Bay. Fishing becomes an increasingly important factor in limiting population growth as natural mortality decreases or the population size increases. This means that managing harvest will likely be necessary to achieve the restoration goal. Further, fishing may reduce the development of disease resistance through natural selection, or have other effects that are not modeled, as described in the text.